



American
Heart
Association.

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN®



PALS

**PEDIATRIC
ADVANCED
LIFE SUPPORT**

PROVIDER MANUAL

Pediatric Advanced Life Support Provider Manual

© 2020 American Heart Association

ISBN 978-1-61669-785-3

Printed in the United States of America

First American Heart Association Printing October 2020

eBook edition © 2020 American Heart Association. ISBN 978-1-61669-804-1. 20-3120

Acknowledgments

The American Heart Association thanks the following people for their contributions to the development of this manual: Kelly D. Kadlec, MD, MEd; Mary E. McBride, MD, MEd; Reylon Meeks, RN, BSN, MS, MSN, EMT, PhD; Sallie Johnson, PharmD, BCPS; Adam Cheng, MD; Ian R. Drennan, ACP, PhD; Susan Fuchs, MD; Mike Helbock, MICP, NR-P, SEI; Elizabeth A. Hunt, MD, MPH, PhD; Garth Meckler, MD, MSHS; Ryan Morgan, MD, MTR; Stephen M. Schexnayder, MD; Sarah Tabbutt, MD, PhD; Janice A. Tijssen, MD, MSc; and the AHA PALS Project Team.



PALS Student Resources can be found at eLearning.heart.org. Contact your Training Center Coordinator for more information about accessing these before your course.

To find out about any updates or corrections to this course, visit www.heart.org/courseupdates.

Contents

Part 1: Course Overview

Course Objectives

Precourse Preparation

Course Materials

Course Completion Requirements

Science Update

The Chain of Survival

Part 2: Review of BLS and AED for Infants and Children

Learning Objectives

BLS for Infants and Children

AED for Infants and Children Younger Than 8 Years

Part 3: High-Performance Teams

High-Performance Team Roles and Dynamics

Part 4: Systematic Approach to the Seriously Ill or Injured Child

Learning Objective

Initial Assessment to Identify a Life-Threatening Condition

Initial Assessment

Evaluate-Identify-Intervene

Primary Assessment

Secondary Assessment

Diagnostic Assessments

Part 5: Recognizing and Managing Cardiac Arrest

Learning Objective

Rapidly Intervene to Prevent Cardiac Arrest

Life-Threatening Problems

Cardiac Arrest in Infants and Children

Definition of Cardiac Arrest

Pathways to Cardiac Arrest

Causes of Cardiac Arrest

Recognizing Cardiac Arrest

Managing Cardiac Arrest

Part 6: Effective High-Performance Team Dynamics

Learning Objective

Roles in a High-Performance Team

Elements of Effective High-Performance Team Dynamics

Part 7: Recognizing Respiratory Distress and Failure

Learning Objective

Fundamental Issues Associated With Respiratory Problems

Identifying Respiratory Problems by Severity

Identifying Respiratory Problems by Type

Summary: Recognizing Respiratory Problems Flowchart

Part 8: Managing Respiratory Distress and Failure

Learning Objective

[Rescue Breathing](#)
[Initial Management of Respiratory Distress and Failure](#)
[Principles of Targeted Management](#)
[Managing Upper Airway Obstruction](#)
[Managing Lower Airway Obstruction](#)
[Managing Lung Tissue Disease](#)
[Managing Disordered Control of Breathing](#)
[Summary: Managing Respiratory Emergencies Flowchart](#)
[Resources for Managing Respiratory Emergencies](#)

Part 9: Recognizing Shock

[Learning Objective](#)
[Defining Shock](#)
[Pathophysiology of Shock](#)
[Identifying Shock by Severity \(Effect on Blood Pressure\)](#)
[Identifying Shock by Type](#)
[Recognizing Shock Flowchart](#)

Part 10: Managing Shock

[Learning Objective](#)
[Goals of Shock Management](#)
[Fundamentals of Shock Management](#)
[General Management of Shock](#)
[Fluid Therapy](#)
[Glucose](#)
[Management According to Type of Shock](#)
[Managing Shock Flowchart](#)
[Resources for Managing Circulatory Emergencies](#)

Part 11: Recognizing Arrhythmias

[Learning Objective](#)
[Bradycardia Definitions](#)
[Recognizing Bradycardia](#)
[Tachyarrhythmias](#)

Part 12: Managing Arrhythmias

[Learning Objective](#)
[Principles of Managing Pediatric Arrhythmias](#)
[Management: Pediatric Bradycardia With a Pulse](#)
[Managing Tachyarrhythmias](#)
[Summary of Emergency Interventions](#)
[Pediatric Tachycardia With a Pulse Algorithm](#)

Part 13: Post-Cardiac Arrest Care

[Learning Objective](#)
[Goals of Therapy](#)
[Respiratory System](#)
[Cardiovascular System](#)
[Neurologic System](#)

Appendix

[BLS Skills Testing Checklists](#)
[Initial Assessment—Pediatric Assessment Triangle](#)

Primary Assessment

Skills Station Competency Checklists

Rhythm Recognition Review

Learning Station Competency Checklists and PALS Case Scenario Testing Checklists

Part 1: Course Overview

Course Objectives

The Pediatric Advanced Life Support (PALS) Provider Course is designed for healthcare providers who manage respiratory and/or cardiovascular emergencies and cardiopulmonary arrest in pediatric patients. By preparing for the course and participating in the skills stations and simulated cases, you will enhance the recognition of and intervention for respiratory emergencies, shock, and cardiopulmonary arrest.

During the course, you will participate in a series of case scenario practices with simulations that reinforce important concepts, including

- Identifying and treating problems that place the child at risk for cardiac arrest
- Applying a systematic approach to pediatric assessment
- Using the evaluate-identify-intervene sequence
- Using the PALS algorithms and flowcharts
- Demonstrating effective resuscitation team dynamics

Goal of the PALS Provider Course

The PALS Provider Course aims to improve outcomes for pediatric patients by preparing healthcare providers to effectively recognize and intervene in patients with respiratory emergencies, shock, and cardiopulmonary arrest by using high-performance team dynamics and high-quality individual skills.

Learning Objectives

After successfully completing this course, you will be able to

- Perform high-quality cardiopulmonary resuscitation (CPR) per American Heart Association (AHA) basic life support (BLS) recommendations
- Perform your role as a high-performance team member
- Differentiate between patients who require immediate intervention and those who don't
- Differentiate between respiratory distress and failure
- Perform early interventions for respiratory distress and failure
- Differentiate between compensated and hypotensive shock
- Perform early interventions for the treatment of shock
- Differentiate between unstable and stable patients with dysrhythmias
- Demonstrate treatment of dysrhythmias
- Implement postarrest management

Course Description

To help you achieve these objectives, the PALS Provider Course includes

- BLS competency testing
- Skills stations
- Case scenario discussions and simulations
- Case scenario testing stations
- An exam

BLS Competency Testing

To receive an AHA PALS Provider course completion card, you must pass 2 BLS tests: the Child CPR and AED Skills Test and the Infant CPR Skills Test.

The PALS Provider Course does not include detailed instructions on how to perform basic CPR or how to use an automated external defibrillator (AED), so you must know this in advance. Consider taking a BLS course to prepare, if necessary.

Before taking the PALS Provider Course, read the *PALS Provider Manual* to prepare for taking the BLS tests.

Skills Stations

The course includes the following skills stations:

- Airway Management
- Rhythm Disturbances/Electrical Therapy
- Vascular Access

In the skills stations, you will use the skills station competency checklists as you practice specific skills and then demonstrate competency. Your instructor will evaluate your skills by using the criteria in these checklists.

To prepare, see the [Appendix](#) for the skills station competency checklists, which list detailed steps for performing each skill.

Airway Management Skills Station

In the Airway Management Skills Station, you will show your understanding of oxygen (O₂) delivery systems and airway adjuncts. You will practice and demonstrate competency in airway and breathing supports, including

- Inserting an oropharyngeal airway
- Providing effective bag-mask ventilation
- Suctioning with an oropharyngeal airway and endotracheal (ET) tube
- Confirming advanced airway device placement by physical examination and an exhaled CO₂ detector device
- Securing the ET tube

If it is within your scope of practice, you may be asked to demonstrate advanced airway skills, including correctly inserting an ET tube.

Review the [Bag-Mask Ventilation](#) and [Endotracheal Intubation](#) sections in [Resources for Managing Respiratory Emergencies](#) in [Part 8](#) of this manual to learn more about airway management skills.

Rhythm Disturbances/Electrical Therapy Skills Station

In the Rhythm Disturbances/Electrical Therapy Skills Station, you will practice and demonstrate competency in identifying rhythms and operating a cardiac monitor and manual defibrillator. Skills include

- Correctly placing electrocardiographic (ECG) leads
- Correctly selecting and placing or positioning the paddles or pads
- Identifying rhythms that require defibrillation
- Identifying rhythms that require synchronized cardioversion
- Operating a cardiac monitor
- Safely performing manual defibrillation and synchronized cardioversion

If necessary, consider taking a BLS course to learn about cardiac monitoring and manual defibrillation. Review the following resources in the *PALS Provider Manual* before the course to help you with rhythm identification, cardiac monitoring, and manual defibrillation

- [Rhythm Recognition Review](#) in the [Appendix](#)
- [Management: Pediatric Bradycardia With a Pulse](#) in [Part 12: Managing Arrhythmias](#)
- [Managing Tachyarrhythmias](#) in [Part 12: Managing Arrhythmias](#)
- [Table 22: Summary of High-Quality CPR Components for BLS Providers](#) in [Part 5: Recognizing and Managing Cardiac Arrest](#)
- [Pediatric Cardiac Arrest Algorithm](#) in [Part 5: Recognizing and Managing Cardiac Arrest](#)

Vascular Access Skills Station

In the Vascular Access Skills Station, you will practice intraosseous (IO) access and other related skills, including

- Inserting an IO needle
- Summarizing how to confirm that the needle has entered the marrow cavity
- Summarizing/demonstrating the method of giving an intravenous (IV)/IO bolus
- Using color-coded length-based resuscitation tape to calculate correct drug doses

Review the [Intraosseous Access](#) and [Color-Coded Length-Based Resuscitation Tape](#) sections in [Resources for Managing Circulatory Emergencies](#) in [Part 10](#) of this manual to learn more about vascular access skills.

PALS Case Scenario Discussions and Case Scenario Practice With Simulations

In the learning stations, you will participate in a variety of learning activities, including

- Case scenario discussions, using a systematic approach for evaluation and decision-making
- Case scenario practice with simulations

In these learning stations, you will practice essential skills both individually and as part of a team. Because this course emphasizes effective team skills as a vital part of the resuscitative effort, you'll practice as a team member and as a Team Leader.

The focus of the PALS Provider Course is to teach you to use a systematic approach when caring for a critically ill or injured child. Read and study the entire *PALS Provider Manual* to understand all the necessary concepts.

PALS Case Scenario Testing Stations

At the end of the course, your team must pass 2 case scenario testing stations to demonstrate the course objectives. You may use the PALS Reference Card and the *2020 Handbook of Emergency Cardiovascular Care for Healthcare Providers* (Handbook of ECC). These simulated clinical scenarios will test the following:

- Ability to evaluate and identify specific clinical conditions covered in the course
- Recognition and management of respiratory and shock emergencies
- Interpretation of core arrhythmias and management by using appropriate medications and electrical therapy
- Performance as an effective Team Leader

This evaluation emphasizes your ability to integrate BLS and PALS skills according to each person's scope of practice.

Review the [Roles in a High-Performance Team](#) section in [Part 6: Effective High-Performance Team Dynamics](#) in this manual to prepare yourself to participate as a Team Leader in the case scenario testing stations.

Exam

The exam measures your mastery of cognitive skills. You must score at least 84% on the exam in the classroom-based course to meet course completion requirements. The AHA has an open-resource policy for exams in eLearning and classroom-based courses.

Open resource means that students may refer to other resources while completing the exam. These resources could include the provider manual, either in printed form or as an eBook on a personal device; any notes the student took during the provider course; the Handbook of ECC; the *2020 AHA Guidelines for CPR and ECC*; posters; etc. *Open resource* does not mean open discussion with other providers or the instructor.

Precourse Preparation

To pass the PALS Provider Course, *you must prepare before the course*. Do the following:

- Take the precourse self-assessment and, if requested by your instructors, complete the online precourse work
- Make sure you are proficient in BLS skills
- Practice identifying and interpreting core ECG rhythms
- Study basic pharmacology and know when to use which drug
- Practice applying your knowledge to clinical scenarios

Precourse Self-Assessment and Precourse Work



Complete the precourse self-assessment at eLearning.heart.org/courses before you take the PALS Provider Course. (Refer to the PALS Student Resources section for details about how to access this resource.) Print your certificate of completion and bring it with you to the course. Because the PALS Provider Course does not teach algorithms, ECG recognition, pharmacology, or BLS skills, use the precourse self-assessment to identify any

deficiencies in your knowledge of these topics. The precourse self-assessment will summarize your strengths and weaknesses. Study the applicable content in the *PALS Provider Manual* or other supplementary resources.

If requested by your instructor, also complete the online precourse work (video lessons) at eLearning.heart.org/courses. You must pass the precourse self-assessment before gaining access to the video lessons. The precourse work will ensure that you are prepared for the course.

BLS Skills

Strong BLS skills are the foundation of advanced life support. Everyone who cares for pediatric patients must be able to perform high-quality CPR, because without it, PALS interventions will fail. For this reason, you must pass the Child CPR and AED and Infant CPR Skills Tests in the PALS Provider Course. *Make sure that you are proficient in BLS skills before attending the course.*

Refer to the [BLS Competency Testing](#) section in the [Appendix](#) for testing requirements and resources.

ECG Rhythm Identification

You must be able to identify and interpret the following core rhythms during case scenario practice with simulations and case scenario tests:

- Normal sinus rhythm
- Sinus bradycardia
- Sinus tachycardia
- Supraventricular tachycardia
- Ventricular tachycardia
- Ventricular fibrillation
- Asystole
- Pulseless electrical activity

The ECG rhythm identification section of the precourse self-assessment will help you evaluate your ability to identify these core rhythms. If you have difficulty with pediatric rhythm identification, study the [Rhythm Recognition Review](#) section in the [Appendix](#).

Basic Pharmacology

You must know basic information about drugs used in the PALS algorithms and flowcharts, including the indications, contraindications, and methods of administration. You will need to know *when* to use *which* drug based on the clinical situation.

The pharmacology section of the precourse self-assessment will help you evaluate and refresh your knowledge of medications used in the course. If you have difficulty with this section of the precourse self-assessment, study the *PALS Provider Manual* and the Handbook of ECC.

Practical Application of Knowledge to Clinical Scenarios



The practical application section of the precourse self-assessment will help you evaluate your ability to apply your knowledge to clinical scenarios. You will need to make decisions based on

- The Systematic Approach Algorithm and the evaluate-identify-intervene sequence
- Identification of core rhythms
- Knowledge of core medications
- Knowledge of PALS flowcharts and algorithms

You should understand the Systematic Approach Algorithm and the evaluate-identify-intervene sequence. Review the core rhythms and medications. Be familiar with the PALS algorithms and flowcharts so that you can apply them to clinical scenarios, but note that the PALS Provider Course does not teach the details of each algorithm. You can find more information in the *PALS Provider Manual*, PALS student resources at eLearning.heart.org/courses, and the Handbook of ECC.

Course Materials



The PALS Provider Course materials consist of the *PALS Provider Manual*, the PALS Reference Card, and the student resources at eLearning.heart.org/courses.

PALS Provider Manual

The *PALS Provider Manual* contains important information that you will use *before, during, and after* the course. This material includes concepts of pediatric evaluation and the recognition and management of respiratory, shock, and cardiac emergencies. *Remember to take this manual with you to class.* Throughout the *PALS Provider Manual*, you will find specific information in the following callout box:



Critical Concepts

These boxes contain the most important information you must know, including specific risks associated with certain interventions and additional background on key topics this course covers.

PALS Reference Card

The PALS Reference Card is a valuable learning aid that contains the following resources:

- Vital Signs in Children
- PALS Systematic Approach Algorithm
- Pediatric Septic Shock Algorithm
- Drugs Used in PALS
- Pediatric Color-Coded Length-Based Resuscitation Tape Chart
- Pediatric Cardiac Arrest Algorithm
- Pediatric Bradycardia With a Pulse Algorithm
- Pediatric Tachycardia With a Pulse Algorithm
- PALS Management of Shock After ROSC Algorithm
- Components of Post-Cardiac Arrest Care

Take your PALS Reference Card with you to the course to use as a reference during the case scenario discussions and the exam.

PALS Student Resources



Go to eLearning.heart.org/courses to access the precourse self-assessment as well as additional information about basic PALS concepts. If you haven't already logged in, the system will ask you to do so. If you haven't visited the site before, you'll be prompted to set up an account. Once logged in, find your course name and select Launch to begin.

The precourse self-assessment is a vital part of your course preparation. Feedback from this assessment will help you identify gaps in your knowledge so that you can target specific material to study.

The precourse self-assessment has 3 parts:

- ECG rhythm identification
- Pharmacology
- Practical application

Complete these assessments *before the course* to identify gaps in and improve your knowledge. Print out your certificate of completion and take it with you to the course.

Student Resources also include PALS videos, precourse work (video lessons) if assigned, a precourse preparation checklist, and supplementary information.

Course Completion Requirements

To successfully complete the PALS Provider Course and obtain your course completion card, you must

- Actively participate in, practice, and complete all skills stations and learning stations
- Pass the Child CPR and AED and Infant CPR Skills Tests
- Pass an exam with a minimum score of 84% (classroom-based course only)
- Pass 2 PALS case scenario tests

Science Update

We have updated the PALS Provider Course to incorporate the recommendations of the *2020 AHA Guidelines for CPR and ECC*. On an ongoing basis, hundreds of international resuscitation scientists and experts evaluate, discuss, and debate thousands of scientific publications. Then, they reach consensus on the best treatments based on the evidence, which form the basis for developing the guidelines. Some recommendations are new, whereas others modify previous recommendations. The following list highlights many of the key recommendations for pediatric basic and advanced life support.

Major Science Changes in 2020

Some of the major science changes in 2020 include the following:

- High-quality CPR should focus on providing adequate chest compression rate and depth, minimizing interruptions in CPR, allowing complete chest recoil between compressions, and avoiding excessive ventilation.
- Target a respiratory rate of 20 to 30 breaths per minute for infants and children who are receiving CPR with an advanced airway in place or are receiving rescue breathing and have a pulse.
- For patients with nonshockable rhythms, the earlier epinephrine is administered after CPR initiation, the more likely the patient is to survive.
- Using a cuffed endotracheal tube decreases the need for endotracheal tube changes.
- The routine use of cricoid pressure does not reduce the risk of regurgitation during bag-mask ventilation and may impede intubation success.
- For out-of-hospital cardiac arrest, bag-mask ventilation is reasonable compared with advanced airway interventions such as endotracheal intubation.
- Resuscitation does not end with return of spontaneous circulation (ROSC). Excellent post-cardiac arrest care is critically important to achieving the best patient outcomes. For children who do not regain consciousness after ROSC, this care includes targeted temperature management and continuous electroencephalography monitoring. For all children, the prevention and/or treatment of hypotension, hyperoxia or hypoxia, and hypercapnia or hypocapnia is important.
- In addition to standard BLS care, naloxone is recommended for all pediatric patients with respiratory arrest due to suspected opioid overdose.
- For septic shock, providers can administer either 10 mL/kg or 20 mL/kg of fluid with frequent reassessment for fluid overload. For fluid-refractory septic shock, use epinephrine or norepinephrine infusions.

Respiratory Rate

Because of lack of pediatric studies, respiratory rates during pediatric CPR have previously been extrapolated from adult data. Data show that higher ventilation rates (at least 30 breaths/min in infants younger than 1 year and at least 25 breaths/min in children 1 year and older) were associated with improved rates of ROSC and survival in cardiac arrest. When performing CPR in infants and children with an advanced airway in place, it may be reasonable to target a respiratory rate range of 1 breath every 2 to 3 seconds (20-30 breaths/min), accounting for age and clinical condition.

Cuffed Endotracheal Tubes

Intubation with a cuffed endotracheal tube can improve capnography and ventilation in patients with poor pulmonary compliance and decrease the need for endotracheal tube changes. Three systematic reviews of the literature show that cuffed tubes decrease the need for reintubation, improve successful ventilation and the accuracy of capnography, and may decrease the risk of aspiration.

Cricoid Pressure

Although cricoid pressure may be useful in certain circumstances, clinical studies show that its routine use reduces the rate of first-attempt intubation success because it can impede visualization during laryngoscopy and chest rise

with bag-mask ventilation. Therefore, routine use of cricoid pressure is not recommended during endotracheal intubation of pediatric patients.

Early Epinephrine

The goal of epinephrine administration during CPR is to optimize coronary perfusion pressure and maintain cerebral perfusion pressure. Earlier administration of epinephrine during CPR may increase survival-to-discharge rates. For pediatric patients in any setting, it is reasonable to administer the initial dose of epinephrine within 5 minutes after the start of chest compressions.

Diastolic Blood Pressure to Guide CPR

For patients with continuous invasive arterial blood pressure monitoring in place at the time of cardiac arrest, it is reasonable for providers to use diastolic blood pressure to assess CPR quality. Although ideal blood pressure targets during CPR are not known, diastolic blood pressure is a marker of adequate blood flow during CPR and may be used if an arterial line is in place.

Seizures After Cardiac Arrest

Post–cardiac arrest seizures are common. Many are nonconvulsive, which can be detected only with electroencephalography monitoring. When resources are available, continuous electroencephalography monitoring is recommended for detecting seizures after cardiac arrest in patients with persistent encephalopathy. It is recommended to treat clinical seizures that follow cardiac arrest.

Septic Shock

Previous AHA guidelines for managing septic shock included aggressive (20 mL/kg) fluid boluses and lacked additional guidance. The *2020 AHA Guidelines for CPR and ECC* suggest a more tailored approach to fluid administration and provide recommendations for vasopressor use. In patients with septic shock, it is reasonable to administer fluid boluses in 10-mL/kg or 20-mL/kg aliquots with frequent reassessment. After every fluid bolus, providers should reassess the patient for fluid responsiveness and for signs of volume overload. Either isotonic crystalloids or colloids can be effective as the initial fluid choice for resuscitation, and either balanced or unbalanced solutions can be effective as the fluid choice for resuscitation. In infants and children with fluid-refractory septic shock, it is reasonable to use either epinephrine or norepinephrine as an initial vasoactive infusion.

Opioid Overdose

Pediatric opioid overdose management is the same as it is for adults. For a patient with suspected opioid overdose who has a definite pulse but no normal breathing or is only gasping (ie, a respiratory arrest), in addition to providing standard BLS care, it is reasonable for responders to administer intramuscular or intranasal naloxone.

The Chain of Survival

For many years, the American Heart Association has adopted, supported, and helped develop the concept of emergency cardiovascular care. The term *Chain of Survival* provides a useful metaphor for the elements of the emergency cardiovascular care systems-of-care concept. The Chain of Survival shows the actions that must take place to give the cardiac arrest victim the best chance of survival. Each link is independent, yet connected, to the links before and after. If any link is broken, the chance for a good outcome decreases.

Overview

Cardiac arrest can happen anywhere—on the street, at home, or in a hospital emergency department (ED), in-patient bed, or intensive care unit. Elements in the system of care and order of actions in the pediatric Chains of Survival differ based on whether the victim has the arrest outside the hospital or inside the hospital.

Actions in the pediatric Chains of Survival differ according to setting (in-hospital vs out-of-hospital). Here are the specific pediatric Chains of Survival ([Figure 1](#)):

- Pediatric in-hospital cardiac arrest
- Pediatric out-of-hospital cardiac arrest



Figure 1A. American Heart Association 2020 pediatric Chains of Survival. Links in the pediatric Chain of Survival will differ based on whether the arrest occurs in or out of the hospital. A, Pediatric In-Hospital Chain of Survival.



Figure 1B. Pediatric Out-of-Hospital Chain of Survival.

Chain of Survival Elements

Although there are slight differences in the pediatric Chains of Survival based on the location of the cardiac arrest, each includes the following elements:

- Prevention and preparedness
- Activating the emergency response system
- High-quality CPR, including early defibrillation
- Advanced resuscitation interventions
- Post–cardiac arrest care
- Recovery

Part 2: Review of BLS and AED for Infants and Children

This Part describes BLS for infants and children and discusses using an AED in infants and children younger than 8 years.

The following age definitions are used in BLS:

- Infants are younger than 1 year (excluding the newly born).
- Children range from 1 year of age to puberty. Signs of puberty include chest or underarm hair on boys and any breast development in girls.

Learning Objectives


After completing this Part, you should be able to

- Perform high-quality CPR for a child
- Perform high-quality CPR for an infant
- Describe the importance of early use of an AED for infants and children younger than 8 years
- Demonstrate the appropriate use of an AED for infants and children younger than 8 years

BLS for Infants and Children

Pediatric BLS Algorithm for Healthcare Providers—Single Rescuer

The Pediatric BLS Algorithm for Healthcare Providers—Single Rescuer outlines steps for a single rescuer of an unresponsive infant or child (Figure 2). Refer to this algorithm as you read the following steps.



Infant and Child 1-Rescuer BLS Sequence

In these sequences, you are the rescuer. If you are alone and encounter an unresponsive infant or child, follow the steps outlined in the [Pediatric BLS Algorithm for Healthcare Providers—Single Rescuer](#) (Figure 2).

Verify scene safety, check for responsiveness, and get help (Steps 1 and 2)

The first rescuer who arrives at the side of an unresponsive infant or child should quickly perform the following steps:

1. Verify scene safety.
2. Check for responsiveness. Tap the child's shoulder or the heel of the infant's foot and shout, "Are you OK?"

3. If the infant or child is not responsive, shout for nearby help. Activate the emergency response system via mobile device (if appropriate).

Assess for breathing and pulse (Step 3)

Next, assess the infant or child for normal breathing and a pulse to help you determine the next appropriate actions.

To minimize delay in starting CPR, take no more than 10 seconds to assess breathing at the same time as you check the pulse.

Breathing

Look for no breathing or only gasping and check pulse (simultaneously). To check for breathing, scan the chest for rise and fall for no more than 10 seconds (Step 3).

- If the infant or child or child has normal breathing and a pulse is felt, monitor until emergency responders arrive (Step 3a).
- If the infant or child is not breathing or is only gasping and pulse is felt, provide rescue breathing (Step 3b).
- If there is no breathing or only gasping and a pulse is not felt, the infant or child has cardiac arrest (Step 5). Gasping is not considered normal breathing and can be a sign of cardiac arrest.

Check Pulse

Infant: To perform a pulse check in an infant, palpate, or feel for, a brachial pulse ([Figure 3A](#)).



Figure 3A. Pulse check. A, In an infant, feel for a brachial pulse.

Child: To perform a pulse check in a child, palpate a carotid or femoral pulse ([Figures 3B and C](#)).




Figure 3B. In a child, feel for a carotid pulse.




Figure 3C. Or, a femoral pulse.

It can be difficult to determine the presence or absence of a pulse, particularly in an infant or child. So, if you do not definitely feel a pulse within 10 seconds, start CPR, beginning with chest compressions.

Infant: Locating the Brachial Artery Pulse

To perform a pulse check in an infant, palpate a brachial pulse. If you do not definitely feel a pulse within 10 seconds, begin high-quality CPR, starting with chest compressions.

Follow these steps to locate the brachial artery pulse:

1. Place 2 or 3 fingers on the inside of the upper arm, midway between the infant's elbow and shoulder.
2. Then press the fingers to attempt to feel for the pulse for at least 5 but no more than 10 seconds ([Figure 3A](#)).

Child: Locating the Femoral Artery Pulse

To perform a pulse check in a child, palpate a carotid or femoral pulse. If you do not definitely feel a pulse within 10 seconds, begin high-quality CPR, starting with chest compressions.

Follow these steps to locate the femoral artery pulse:

1. Place 2 fingers in the inner thigh, midway between the hipbone and the pubic bone and just below the crease where the leg meets the torso ([Figure 3C](#)).
2. Feel for a pulse for *at least 5 but no more than 10 seconds*. If you do not feel a pulse, begin high-quality CPR, starting with chest compressions.

Determine Next Actions

Determine your next actions on the basis of the presence or absence of normal breathing and a pulse:

- If the infant or child has normal breathing and a pulse is felt, monitor the victim until emergency responders arrive (Step 3a).
- If the infant or child does not have normal breathing and a pulse is felt, provide rescue breathing and give 1 breath every 2 to 3 seconds, or about 20 to 30 breaths/min (refer to [Rescue Breathing in Part 8](#)) (Step 3b).
- Assess the pulse for no more than 10 seconds (Step 3b).
 - If heart rate is less than 60/min with signs of poor perfusion (Step 4) start CPR (Step 4a).
 - If heart rate is 60/min or greater (Step 4), continue rescue breathing and check the pulse every 2 minutes. If no definite pulse is felt, start CPR (Step 4b).
- If the infant or child is not breathing or is only gasping and a pulse is not felt (Step 5),
 - If you are alone and witness the sudden collapse of an infant or child, activate the emergency response system (if not already done), and retrieve an AED/defibrillator (Step 5a). For example, phone 9-1-1 from your phone, mobilize the code team, or notify advanced life support.
 - If you are alone and did not witness the sudden collapse of an infant or child, continue to the next step: Start high-quality CPR for 2 minutes (Step 6).



Critical Concepts: Signs of Poor Perfusion

- Temperature: Cool extremities
 - Altered mental state: Continued decline in consciousness/responsiveness
 - Pulses: Weak pulses
 - Skin: Paleness, mottling (patchy appearance), and later cyanosis (turning blue)
-

Was the Collapse Sudden?

If the collapse was sudden and witnessed, leave the infant or child and activate the emergency response system (unless you have already done so by mobile device) and retrieve the AED. If others arrive, send them to activate the emergency response system (if not already done) and retrieve the AED while you remain with the child to start CPR.

Begin High-Quality CPR, Starting With Chest Compressions

If the infant or child is not breathing normally or is only gasping and no pulse is felt, begin high-quality CPR, starting with chest compressions (refer to details in [Part 5: Recognizing and Managing Cardiac Arrest](#)). Quickly move bulky clothing covering the child or infant's chest out of the way so that you can locate appropriate hand or finger placement for compressions. If clothes are difficult to remove, you can still provide compressions over clothing. If an AED becomes available, remove all clothes that cover the chest. AED pads must not be placed over any clothing.

Single rescuers should use the following compression techniques (refer to [Infant/Child Chest Compressions](#) later in this Part for complete details):

- Infant: 2-finger chest compressions or 2 thumb–encircling hands technique

- Child: 1 or 2 hands (whatever is needed to provide compressions of adequate depth)

After about 2 minutes of CPR, if you are still alone and were unable to activate the emergency response system (have no mobile phone), leave the infant or child and activate the emergency response system and get the AED (Step 7).

Attempt Defibrillation With the AED

Use the AED as soon as it is available and follow the prompts (Step 8).

Resume High-Quality CPR

After shock delivery or if no shock is advised, immediately resume high-quality CPR for 2 minutes (until prompted by the AED to allow for a rhythm check). Continue until ALS providers take over or the child begins to breathe, move, or otherwise react (Steps 9 and 10).

Infant/Child Chest Compressions

Compression Rate and Compression-to-Ventilation Ratio

The *universal* rate for compressions in all cardiac arrest victims is 100 to 120/min. The compression-to-ventilation ratio for single rescuers is the same (30:2) in adults, children, and infants.

If 2 rescuers are present for the resuscitation attempt of an infant or child, use a compression-to-ventilation ratio of 15:2.

Chest Compression Technique

For most children, use either 1 or 2 hands to compress the chest. For most children, the compression technique will be the same as for an adult: 2 hands (heel of one hand with heel of other hand on top of the first hand). For a very small child, you may be able to use 1-handed compressions to achieve the desired compression depth. Compress the chest at least one third the anteroposterior (AP) diameter of the chest (approximately 2 inches, or 5 cm) with each compression.

For infants, single rescuers should use 2 fingers or 2 thumbs placed just below the nipple line. If multiple rescuers are present, the 2 thumb–encircling hands technique is preferred because it produces better blood supply to the heart muscle, helps ensure consistent depth and force of chest compressions, and may generate higher blood pressures. These techniques are described in the next sections.

Infant (1 Rescuer): 2-Finger Technique

Follow these steps to use the 2-finger technique on an infant:

1. Place the infant on a firm, flat surface.
2. Place 2 fingers in the center of the infant's chest, just below the nipple line, on the lower half of the sternum. Do not press the tip of the sternum ([Figure 4](#)).
3. Give compressions at a rate of 100 to 120/min.
4. Compress at least one third the AP diameter of the infant's chest (approximately 1½ inches [4 cm]).
5. At the end of each compression, make sure you allow the chest to fully recoil (reexpand); do not lean on the chest. Chest compression and chest recoil/relaxation times should be about equal. Minimize interruptions in compressions (eg, to give breaths) to less than 10 seconds.
6. After every 30 compressions, open the airway with a head tilt–chin lift and give 2 breaths, each over 1 second. The chest should rise with each breath.
7. After about 5 cycles or 2 minutes of CPR, if you are alone and the emergency response system has not been activated, leave the infant (or carry the infant with you) and activate the emergency response system and retrieve the AED.
8. Continue compressions and breaths in a ratio of 30:2, and use the AED as soon as it is available. Continue until advanced providers take over or the infant begins to breathe, move, or otherwise react.



Figure 4. Two-finger chest compression technique for an infant.

Infant: 2 Thumb–Encircling Hands Technique

The 2 thumb–encircling hands technique is the preferred 2-rescuer chest compression technique because it improves blood flow. Follow these steps to use the 2 thumb–encircling hands technique on an infant:

1. Place the infant on a firm, flat surface.
2. Place both thumbs side by side in the center of the infant's chest, on the lower half of the sternum. The thumbs may overlap in very small infants. Encircle the infant's chest and support the infant's back with the fingers of both hands.
3. With your hands encircling the chest, use both thumbs to depress the sternum ([Figure 5](#)) at a rate of 100 to 120/min.
4. Compress at least one third the AP diameter of the infant's chest (approximately 1½ inches [4 cm]).
5. After each compression, completely release the pressure on the sternum and allow the chest to recoil completely.
6. After every 15 compressions, pause briefly for the second rescuer to open the airway with a head tilt–chin lift and give 2 breaths, each over 1 second. The chest should rise with each breath. Minimize interruptions in compressions (eg, to give breaths) to less than 10 seconds.
7. Continue compressions and breaths in a ratio of 15:2 (for 2 rescuers). For chest compressions to remain effective, the rescuer providing chest compressions should switch roles with another provider at least every 5 cycles or 2 minutes to avoid fatigue. Continue CPR until the AED arrives, advanced providers take over, or the infant begins to breathe, move, or otherwise respond.



Figure 5. Two thumb–encircling hands technique for an infant (2 rescuers).



Critical Concepts: Compression Depth in Adults vs Children and Infants

- Adults and adolescents: At least 2 inches (5 cm)
- Children: At least one third the AP diameter of the chest or approximately 2 inches (5 cm)
- Infants: At least one third the AP diameter of the chest or approximately 1½ inches (4 cm)

Infant/Child Breaths

Opening the Airway

Open the airway to make rescue breaths effective. Two methods for opening the airway are the head tilt–chin lift and jaw-thrust maneuvers. As with adults, if you suspect a head or neck injury, use the jaw-thrust maneuver. If the jaw thrust does not open the airway, use the head tilt–chin lift. If you tilt (extend) an infant's head beyond the neutral (sniffing) position, the infant's airway may become blocked. Maximize airway patency by positioning the infant with the neck in a neutral position so that the external ear canal is level with the top of the infant's shoulder.

Why Breaths Are Important for Infants and Children in Cardiac Arrest

When *sudden* cardiac arrest occurs, the oxygen content of the blood can typically meet oxygen demands of the body for the first few minutes after arrest. So delivering chest compressions can effectively distribute oxygen to the heart and brain.

In contrast, infants and children who develop cardiac arrest often have respiratory failure or shock that reduces the oxygen content in the blood even before the onset of arrest. As a result, for most infants and children in cardiac

arrest, chest compressions alone do not deliver oxygenated blood to the heart and brain as effectively as compressions and breaths do. *For this reason, it is very important to give both compressions and breaths to infants and children during high-quality CPR.*

Ventilation for an Infant or Child With a Barrier Device


Use a barrier device (eg, pocket mask) or a bag-mask device to deliver breaths to an infant or child.

When providing bag-mask ventilation for an infant or child, do the following:

- Select a bag and mask that covers the mouth and nose completely without covering the eyes or overlapping the chin.
- Perform a head tilt–chin lift to open the airway. Press the mask to the face as you lift the jaw, making a seal between the child’s face and the mask.
- Connect supplemental oxygen when available.

Pediatric BLS Algorithm for Healthcare Providers—2 or More Rescuers

Refer to the [Pediatric BLS Algorithm for Healthcare Providers—2 or More Rescuers](#) as you read the following steps (Figure 6).



Infant and Child 2-Rescuer BLS Sequence

If the rescuer encounters an unresponsive infant or child and other rescuers are available, follow the steps outlined in the [Pediatric BLS Algorithm for Healthcare Providers—2 or More Rescuers](#) (Figure 6).

Verify scene safety, check for responsiveness, and get help (Steps 1 and 2)

In these sequences, you are the rescuer. If you arrive first at the side of an unresponsive infant or child, quickly perform the following steps. As more rescuers arrive, assign roles and responsibilities. When more rescuers are available for a resuscitation attempt, they can perform more tasks simultaneously.

1. Verify that the scene is safe.
2. Check for responsiveness. Tap the child’s shoulder or the heel of the infant’s foot and shout, “Are you OK?”
3. If the infant or child is not responsive, shout for nearby help. The first rescuer remains with the infant or child while the second rescuer activates the emergency response system (Figure 7) and retrieves the AED and emergency equipment.



Figure 7A. If the arrest of an infant or child was sudden

and witnessed, activate the emergency response system in your setting. A, In-facility setting.



Figure 7B. Prehospital setting.

Assess for breathing and pulse (Step 3)

For details on assessing the infant or child for normal breathing and a pulse, refer to the [Infant and Child 1-Rescuer BLS Sequence](#) section earlier in this Part.

Determine Next Actions

For details on determining next actions on the basis of the presence or absence of breathing and pulse, refer to the [Infant and Child 1-Rescuer BLS Sequence](#) earlier in this Part (Steps 3a and 3b, 4, 4a, and 4b). If CPR is indicated when the second rescuer can assist, use a compression-to-ventilation ratio of 15:2.

Begin High-Quality CPR, Starting With Chest Compressions

If the child or infant is not breathing normally or is only gasping and has no pulse, immediately do the following (Step 5):

Start high-quality CPR, starting with chest compressions (refer to [Infant/Child Chest Compressions](#) earlier in this Part for complete details). Quickly move bulky clothing covering the chest out of the way so that you can locate appropriate hand or finger placement for compressions. If the clothes are difficult to remove, you can still provide compressions over clothing. If an AED becomes available, remove all clothes that cover the chest. AED pads must not be placed over any clothing.

- For an infant, single rescuers can use either the 2-finger technique for chest compressions or the 2 thumb–encircling hands technique until the second rescuer returns. When the second rescuer arrives, use the 2 thumb–encircling hands technique.
- For a child, use 1 or 2 hands (1 hand for a very small child) to perform compressions.
- When the second rescuer returns, that rescuer gives breaths.
- Switch compressors about every 5 cycles or 2 minutes (or earlier if needed) to avoid reducing CPR quality because of fatigue.

Attempt defibrillation with the AED (Step 6)

Use the AED as soon as it is available and follow the prompts.

Resume high-quality CPR (Steps 7 and 8)

After shock delivery or if no shock is advised, immediately resume high-quality CPR, starting with chest compressions, when advised by the AED. Continue providing CPR and following the AED prompts until advanced life support providers take over or the infant or child starts to move.

AED for Infants and Children Younger Than 8 Years

Be Familiar With the AED Equipment in Your Setting

Although all AEDs operate in basically the same way, AED equipment varies according to model and manufacturer. For instance, some AED models called *pediatric-capable AEDs*, are designed for both pediatric and adult use. These AEDs deliver a reduced shock dose when pediatric pads are used. Be familiar with the AED in your particular setting.

Delivering a Pediatric Shock Dose

You may reduce the AED shock dose by using pediatric cables, an attenuator, or preprogramming in the device. One commonly used method for reducing a shock dose is a pediatric dose attenuator ([Figure 8](#)). When attached to an AED, it reduces the shock dose by about two thirds. Typically, you can use child pads to deliver the reduced shock dose.




Figure 8. Example of a pediatric dose attenuator, which reduces the shock dose delivered by an AED. Child pads are also used with this attenuator.

Choosing and Placing the AED Pads

Use child pads ([Figure 9](#)), if available, for infants and for children younger than 8 years. If child pads are not available, use adult pads. Adult pads deliver a higher shock dose, but that is preferred to no shock at all. However, make sure the pads do not touch each other or overlap. Also, if the AED has a key or switch that will deliver a child shock dose, turn the key or switch. For victims 8 years and older, use adult pads ([Figure 10](#))—child pads will likely give a shock dose that is too low.



Figure 9. Child AED pads.



Figure 10. Adult AED pads.

Follow the AED manufacturer's instructions for pad placement and the illustrations on the AED pads. Some AEDs require that child pads be placed in a front and back (anteroposterior) position ([Figure 11](#)), while others require right-left (anterolateral) placement. Anteroposterior pad placement is commonly used for infants.




Figure 11. Anteroposterior AED pad placement on a child victim.

Using an AED for Infants

For infants, a manual defibrillator is preferred to an AED for defibrillation because it has more capabilities than an AED and can provide lower energy doses needed in infants. Operating a manual defibrillator requires advanced training that will not be covered in this course.

- If a manual defibrillator is not available, an AED equipped with a pediatric dose attenuator is preferred.
- If neither is available, you may use an AED without a pediatric dose attenuator.


Part 3: High-Performance Teams

High-performance teams are essential to successful resuscitation attempts. High-performance teams carry out their roles in highly effective manners, resulting in superior performance and timing, which can translate to improved survival for patients in cardiac arrest. What distinguishes high-performance teams from others is that each team member is committed to ensuring the highest-quality performance of the team rather than simply following orders. To function effectively, a high-performance team needs to focus on

- **Timing** (immediate CPR and defibrillation, limiting pauses to achieve a chest compression fraction (CCF) of ideally greater than 80%_ preshock pause, EMS response time)
- **Quality** (best possible performance of each team member, eg, rate, depth, recoil)
- **Coordination** (team dynamics: team members working together seamlessly toward a common goal, proficient in their roles)
- **Administration** (leadership, planning ahead, obtaining additional resources, tailoring the interventions, CQI, measurement, assigned number of members working an arrest)

_High-performing systems target at least 60%, with 80% or higher being a frequent goal.

High-performance teams (Figure 12) will need to incorporate timing, quality, coordination, and administration of the appropriate procedures during a cardiac arrest. The team will need to consider their overall purpose and goals, skills each team member possesses, appropriate motivation and efficacy, as well as appropriate conflict resolution and communication needs of the team. In addition, high-performance teams measure their performance, evaluate the data, and look for ways to improve performance and implement the revised strategy.



Critical Concepts: Ways to Increase CCF

Whether you are a team member or the Team Leader during a resuscitation attempt, you should understand how a high-performance team can maximize CCF when performing CPR during a cardiac arrest. The team can achieve key metrics and increase CCF by doing the following:

- **Precharge the defibrillator** 15 seconds before a 2-minute rhythm analysis (deliver shock immediately if VF or pVT on the monitor). This makes it possible to conduct a rhythm analysis and give a shock (if needed) within 10 seconds or less.
- Perform a pulse check during the precharge phase in anticipation of an organized rhythm during analysis (a pulse check during compressions is not a reliable indicator of CPR quality).
- Compressor **hovers over the chest** (not touching it), ready to start chest compressions immediately after a shock, a rhythm analysis, or other necessary pauses in compressions.
- **Have the next compressor ready to take over immediately.**
- Intubate without pausing compressions.
- Deliver medications during compressions.
- Consider CPR protocols that deliver fewer pauses (eg, continuous compressions with asynchronous ventilation using a bag-mask device).

High-Performance Team Roles and Dynamics

Successful resuscitation attempts often require healthcare providers to simultaneously perform a variety of interventions. Although a CPR-trained bystander working alone can resuscitate a patient within the first moments after collapse, most attempts require multiple healthcare providers. Effective teamwork divides the tasks while multiplying the chances of a successful outcome.

Successful high-performance teams not only have medical expertise and mastery of resuscitation skills but also demonstrate effective communication and team dynamics. This section discusses the importance of team roles, behaviors of effective Team Leaders and team members, and elements of effective high-performance team dynamics.



Critical Concepts: Understanding Team Roles

Whether you are a team member or a Team Leader during a resuscitation attempt, you should understand your role and the roles of other members. This awareness will help you anticipate

- What actions will be performed next
 - How to communicate and work as a member or as a leader of a high-performance team
-

Roles in a High-Performance Team

Team Leader Role

Every high-performance team needs a leader to organize the efforts of the group. The Team Leader

- Organizes the group
- Monitors individual performance of team members
- Backs up team members
- Models excellent team behavior
- Trains and coaches
- Facilitates understanding
- Focuses on comprehensive patient care
- Temporarily designates another team member to take over as Team Leader if an advanced procedure is required (eg, advanced airway placement)

The Team Leader is responsible for making sure everything is done at the right time in the right way by monitoring and integrating individual performance of team members. The Team Leader should also help train future Team Leaders and improve team effectiveness. After resuscitation, the Team Leader can help analyze, critique, and practice for the next resuscitation attempt.

The Team Leader also helps team members understand why they must perform certain tasks in a specific way. The Team Leader should be able to explain why it is essential to

- Push hard and fast in the center of the chest
- Ensure complete chest recoil
- Minimize interruptions in chest compressions
- Avoid excessive ventilation

Whereas members of a high-performance team should focus on their individual tasks, the Team Leader must focus on comprehensive patient care.

Team Member Roles

For a successful resuscitation attempt, high-performance team members must be

- Proficient in performing the skills in their scope of practice
- Clear about role assignments
- Prepared to fulfill their role responsibilities
- Well-practiced in resuscitation skills
- Knowledgeable about the algorithms

- Committed to success

Team Member Role: CPR Coach

Many resuscitation teams now include the role of CPR Coach. The CPR Coach supports performance of high-quality BLS skills, allowing the Team Leader to focus on other aspects of clinical care. Studies have shown that resuscitation teams with a CPR Coach perform higher-quality CPR with higher CCF and shorter pause durations compared with teams that do not use a CPR Coach.

The CPR Coach does not need to be a separate role; they can be most effectively blended into the current responsibilities of the Monitor/Defibrillator. The CPR Coach's main responsibilities are to help team members provide high-quality CPR and minimize pauses in compressions. The CPR Coach needs a direct line of sight to the Compressor, so they should stand next to the Defibrillator. Below is a description of the CPR Coach's actions:

Coordinate the start of CPR: As soon as a patient is identified as having no pulse, the CPR Coach says, "I am the CPR Coach," and tells rescuers to begin chest compressions. The CPR Coach can adjust the environment to help ensure high-quality CPR. They can lower the bedrails or the bed, get a step stool, or roll the victim to place a backboard and defibrillator pads to better facilitate high-quality CPR.

Coach to improve the quality of chest compressions: The CPR Coach gives feedback about performance of compression depth, rate, and chest recoil. They state the CPR feedback device's data to help the Compressor improve performance. This is useful because visual assessment of CPR quality is often inaccurate.

State the midrange targets: The CPR Coach states the specific midrange targets so that compressions and ventilation are within the recommended range. For example, they should tell the Compressor to compress at a rate of 110/min instead of a rate between 100 and 120/min.

Coach to the midrange targets: The CPR Coach gives team members feedback about their ventilation rate and volume. If needed, they also remind the team about compression-to-ventilation ratio.

Help minimize the length of pauses in compressions: The CPR Coach communicates with the team to help minimize the length of pauses in compressions. Pauses happen when the team defibrillates, switches Compressors, and places an advanced airway.



Critical Concepts: CPR Coach Role

The CPR Coach role is designed to help a high-performance team achieve the key metrics of high-quality CPR by providing feedback about

- The Compressor's rate, depth, and recoil
- Delivery of ventilations (rate and volume)
- Compression pauses

Working closely with the Team Leader, the CPR Coach should facilitate all compression pauses, including intubation. The CPR Coach should be integrated into the existing role of Monitor/Defibrillator on a high-performance team.

Elements of Effective Team Dynamics as Part of a High-Performance Team

Roles

Clear Roles and Responsibilities

Every member of the team should know his or her role and responsibilities because each team member's role is important to the performance of the team. [Figure 13](#) identifies 6 team roles for resuscitation. When fewer than 6 people are present, Team Leaders must prioritize these tasks and assign them to the healthcare providers present.




Figure 13. Suggested locations for the Team Leader and team members during case simulations and clinical events.

When roles are unclear, team performance suffers. Signs of unclear roles include

- Performing the same task more than once
- Missing essential tasks
- Assigning team members multiple roles when additional providers are available

For efficiency, the Team Leader must clearly delegate tasks. Team members should communicate when they can handle additional responsibilities. The Team Leader should encourage team members to participate actively and not simply follow directions. [Table 1](#) lists some additional information about roles.

Table 1. Clear Roles and Responsibilities

Team member	Task
Team Leader	<ul style="list-style-type: none"> • Clearly define all team member roles in the clinical setting • Distribute tasks evenly to all available team members who are sure of their responsibilities
Team members	<ul style="list-style-type: none"> • Seek out and perform clearly defined tasks appropriate to their abilities • Ask for a new task or role if an assignment is beyond their level of expertise • Accept only assignments that are within their level of expertise

Knowing Your Limitations

Everyone on the team should know his or her own limitations and capabilities, including the Team Leader. This allows the Team Leader to evaluate resources and call for backup when necessary. High-performance team members should anticipate situations in which they need help and inform the Team Leader.

During the stress of an attempted resuscitation, do not practice or explore a new skill, especially without seeking advice from more experienced personnel. If you need extra help, request it early rather than waiting until the patient deteriorates further. Asking for help is not a sign of weakness or incompetence; it is better to have more help than needed rather than not enough help, which might negatively affect patient outcome. [Table 2](#) lists some additional information about knowing your limitations.

Table 2. Knowing Your Limitations

Team member	Task
Team Leader and team members	<ul style="list-style-type: none"> • Call for assistance early rather than waiting until the patient deteriorates • Seek advice from more experienced personnel when the patient's condition worsens despite primary treatment • Allow others to carry out assigned tasks, especially if the task is essential to treatment
Team members	<ul style="list-style-type: none"> • Seek advice from more experienced personnel before starting an unfamiliar treatment or therapy • Accept assistance from others when it is readily available

Constructive Interventions

During a resuscitation attempt, anyone on a high-performance team may need to intervene tactfully if a team member is about to take an inappropriate action. Team Leaders should avoid confrontation with team members and instead debrief afterward if needed. [Table 3](#) lists some additional information about constructive interventions.

Table 3. Constructive Interventions

Team member	Task
Team Leader	<ul style="list-style-type: none"> • Ask that a different intervention be started if it has a higher priority • Reassign a team member who is trying to function beyond his or her level of skill
Team members	<ul style="list-style-type: none"> • Suggest an alternative drug or dose confidently • Question a colleague who is about to make a mistake

- Intervene if a team member is about to administer a drug incorrectly

What to Communicate

Knowledge Sharing

Sharing information is critical to effective team performance. Team Leaders may become fixated on a specific treatment or diagnostic approach. Examples of these types of *fixation errors* are

- “Everything is OK.”
- “This and only this is the correct path.”
- “Do anything but this.”

When resuscitative efforts are ineffective, go back to the basics and talk as a team. Have conversations like, “Well, we’ve observed the following on the Primary Assessment... Have we missed something?” High-performance team members should provide all available information about changes in the patient’s condition to ensure that the Team Leader makes appropriate decisions. [Table 4](#) lists some additional information about knowledge sharing.

Table 4. Knowledge Sharing

Team member	Task
Team Leader	<ul style="list-style-type: none"> • Encourage information sharing • Ask for suggestions about interventions, differential diagnoses, and possible overlooked treatments (eg, intravenous access or drug treatments) • Look for clinical signs that are relevant to the treatment
Team members	<ul style="list-style-type: none"> • Share information with each other • Accept information that will improve their roles

Summarizing and Reevaluating

An essential role of the Team Leader is monitoring and reevaluating interventions, assessment findings, and the patient’s status.

Team Leaders should periodically state this information to the team and announce the plan for the next few steps. Remember that the patient’s condition can change. Be flexible to changing treatment plans, and ask for information and summaries from the timer/recorder as well. [Table 5](#) lists some additional information about summarizing and reevaluating.

Table 5. Summarizing and Reevaluating

Team member	Task
Team Leader	<ul style="list-style-type: none"> • Continuously revisit decisions about differential diagnoses • Maintain an ongoing record of treatments and the patient’s response • Change a treatment strategy when new information supports it • Inform arriving personnel of the current status and plans for further action
Team Leader and team members	<ul style="list-style-type: none"> • Note significant changes in the patient’s clinical condition • Increase monitoring if patient’s condition deteriorates (eg, frequency of respirations and blood pressure)

How to Communicate

Closed-Loop Communications

When communicating with high-performance team members, the Team Leader should use these closed-loop communication steps:

1. Give a message, order, or assignment to a team member.
2. Request a clear response and eye contact from the team member to ensure that he or she understood the message.
3. Confirm that the team member completed the task before you assign him or her another task.

[Table 6](#) lists some additional information about closed-loop communications.

Table 6. Closed-Loop Communications

--	--

Team member	Task
Team members	<ul style="list-style-type: none"> After receiving a task, close the loop by informing the Team Leader when the task begins or ends, such as, "The IV is in" Only give drugs after verbally confirming the order with the Team Leader
Team Leader	<ul style="list-style-type: none"> Always assign tasks by using closed-loop communication such as, "Give 1 milligram of epinephrine and let me know when it has been given" Assign additional tasks to a team member <i>only</i> after receiving confirmation of a completed assignment

Clear Messages

Clear messages means concise communication spoken with distinctive speech in a controlled voice. All healthcare providers should deliver clear messages calmly and directly, without yelling or shouting. Distinct, concise messages are crucial for clear communication because unclear communication can delay treatment or cause medication errors. Yelling or shouting can also impair effective high-performance team interaction. [Table 7](#) lists some additional information about clear messages.

Table 7. Clear Messages

Team member	Task
Team Leader	<ul style="list-style-type: none"> Encourage all team members to speak clearly and use complete sentences
Team Leader and team members	<ul style="list-style-type: none"> Repeat orders, and question them if the slightest doubt exists Be careful not to mumble, yell, scream, or shout Ensure that only 1 person talks at a time

Mutual Respect

The best high-performance team members mutually respect each other and work together in a collegial, supportive manner. Everyone in a high-performance team must abandon ego and show respect during the resuscitation attempt, regardless of any additional training or experience that specific team members may have. [Table 8](#) lists some additional information about mutual respect.

Table 8. Mutual Respect

Team member	Task
Team Leader	<ul style="list-style-type: none"> Acknowledge correctly completed assignments by saying, "Thanks—good job!"
Team Leader and team members	<ul style="list-style-type: none"> Show interest and listen to what others say Speak in a friendly, controlled tone of voice Avoid displaying aggression if teammates do not initially understand each other Understand that when one person raises his voice, others will respond similarly Try not to confuse directive behavior with aggression

Part 4: Systematic Approach to the Seriously Ill or Injured Child

You should use a systematic approach when caring for a seriously ill or injured child so that you can quickly recognize signs of respiratory distress, respiratory failure, and shock and immediately provide lifesaving interventions.

Learning Objective

After completing this Part, you should be able to differentiate between patients who do and do not require immediate intervention.

You should know all of the concepts presented in this Part to be able to identify the child's clinical condition and target appropriate management in case simulations. The ongoing process of evaluate-identify-intervene is a core component of systematic evaluation and care of a seriously ill or injured child.

Initial Assessment to Identify a Life-Threatening Condition



The PALS Systematic Approach Algorithm (Figure 14) outlines the approach to caring for a critically ill or injured child. Use the Pediatric Assessment Triangle (PAT) to make your initial assessment during your first quick “from the doorway” observation of the child’s appearance, breathing, and color, within the *first few seconds* after encountering the child. You can use the PAT immediately on entering the scene to help identify the general type of physiologic problem (ie, respiratory, circulatory, or neurologic) and urgency for treatment and transport.




Figure 14. PALS Systematic Approach Algorithm.

Identify a Life-Threatening Condition and Act

The initial assessment helps you to quickly identify a life-threatening condition. If at any time you identify a life-threatening problem, immediately begin appropriate interventions and activate the emergency response system as indicated in your practice setting. If the child's condition is not life threatening, continue with the systematic approach.

Child Who Is Unresponsive and Not Breathing or Is Only Gasping

If the child is unresponsive, shout for help. In the text that follows, numbers refer to the corresponding steps in the PALS Systematic Approach Algorithm.

Activate emergency response system (Step 3)

If the child is unresponsive with no normal breathing or only gasping and no pulse is felt, activate the emergency response system as appropriate for your practice setting (Step 3), and immediately start CPR, beginning with chest compressions (Step 7).

Proceed according to the Pediatric Cardiac Arrest Algorithm. If ROSC, go to the Post–Cardiac Arrest Care checklist (Step 9) and begin the evaluate-identify-intervene sequence (Step 12).

Check Breathing and Pulse (Step 4)

Next, check for breathing and a pulse simultaneously (Step 4).

If No Normal Breathing, But a Pulse Is Felt, Provide Rescue Breaths (Step 5)

If a pulse is felt, open and maintain a patent airway and provide rescue breathing (Step 5). Administer oxygen as soon as it is available. (Refer to [Rescue Breathing](#) in [Part 8](#) for more information.) For infants and children, give 1 breath every 2 to 3 seconds (about 20-30 breaths/min), and give each breath over 1 second. Each breath should result in visible chest rise. Monitor pulse and oximetry.

Check the heart rate. If the heart rate is less than 60/min with signs of poor perfusion despite adequate oxygenation and ventilation (Step 6), provide chest compressions and breaths (Step 7). Proceed according to the Pediatric Cardiac Arrest Algorithm.

If the heart rate is 60/min or greater, continue rescue breathing as needed. Begin the evaluate-identify-intervene sequence (Steps 10 through 12). Check the pulse about every 2 minutes. Be prepared to intervene according to the Pediatric Cardiac Arrest Algorithm if needed.

If There Is Normal Breathing and a Pulse Is Felt

If unresponsiveness is a new finding and breathing and pulse are adequate, activate the emergency response system as appropriate for your setting. Continue your systematic assessment (follow Steps 10 through 12).

Initial Assessment



If you do not immediately detect a life-threatening emergency, you can continue forming your initial assessment of the child's condition. Use the PAT to form your initial impression.

The PAT uses A-B-C, which stands for **a**ppearance, **w**ork of **b**reathing, and **c**irculatory status. It begins with evaluating appearance (A) as an indicator of overall physiologic status, including degree of interactivity, muscle tone, and verbal response or cry. Use the TICLS (**t**one, **i**nteractiveness, **c**onsolability, **l**ook/gaze, **s**peech/cry) mnemonic as an adjunct. The second component of the PAT is breathing (B), which determines whether a child has increased work of breathing by assessing the patients' position (ie, tripod or sniffing position), work of breathing (ie, retractions), and adventitious breath sounds (eg, stridor, sonorous respirations). The final component of the PAT evaluates the child's overall circulatory status (C) based on general color (eg, pale, mottled, cyanotic). A child with abnormal PAT findings requires prompt evaluation and management. The findings of the PAT may indicate need for immediate intervention (eg, CPR for a patient who is apneic and pulseless, tourniquet use for exsanguinating hemorrhage of an extremity).

Appearance

Appearance



The first part of the PAT is the child's appearance, including level of consciousness and ability to interact. Carefully, but quickly, observe the child's appearance to evaluate the level of consciousness by observing the child's TICLS. If the child is unresponsive, shout for nearby help, assess breathing and pulse, and then activate the rapid response or the emergency response system as appropriate for your clinical setting.

If the child is crying or upset, it can be difficult to know if she is responding appropriately. Try to keep the child as calm as possible by letting her remain with her parent or caregiver if practical, and use distractions such as toys.

Breathing

Work of Breathing



The next part of the PAT is evaluating the child's work of breathing, position, and any audible breath sounds (ie, sounds that can be heard without a stethoscope) ([Table 9](#)). Look for signs of absent or increased respiratory effort. Listen for obvious sounds of abnormal breathing, such as grunting, stridor, or wheezing. Note whether the patient's position suggests respiratory distress, such as the tripod position.

Table 9. Evaluating Work of Breathing

Evaluate	Normal	Abnormal
Respiratory effort*	<ul style="list-style-type: none">Regular breathing, no increased effortPassive expiration	<ul style="list-style-type: none">Nasal flaringRetractions or use of accessory musclesIncreased, inadequate, or absent respiratory effort
Lung and airway sounds*	No abnormal respiratory sounds audible	Noisy breathing (eg, wheezing, grunting, stridor)

*Refer to detailed discussion of respiratory effort and lung and airway sounds in the [Primary Assessment](#) section in this Part.

Circulation (Color)



Circulation (Color)

To complete the third part of the PAT, assess the child's overall circulatory status. Assess the child's color, including skin color and pattern or obvious significant bleeding, which may help you assess how well the child is perfusing. You can often identify important information about circulatory status just by looking at a child.

Pallor (paleness), mottling (an irregular skin color), or cyanosis (bluish/gray skin color) suggests poor perfusion, poor oxygenation, or both. Cyanosis of the lips and fingernails may appear if the child cannot adequately oxygenate the blood.

Observe the exposed parts of the child, such as the face, arms, and legs. A flushed appearance may suggest fever or distributive shock such as from sepsis, toxins, or anaphylaxis. A skin inspection may reveal bruising that suggests

injury. You may also see evidence of bleeding within the skin, called *petechiae* or *purpura*. This purplish discoloration of the skin may be a sign of a life-threatening infection.

Evaluate the skin and mucous membranes ([Table 10](#)). *Are they normal or abnormal?*


Table 10. Evaluating Skin and Mucous Membranes

Evaluate	Normal	Abnormal
Skin color*	Appears normal	<ul style="list-style-type: none"> • Pallor • Mottling • Cyanosis
Petechiae or purpura or visible bleeding wounds	Not normal	<ul style="list-style-type: none"> • Obvious significant bleeding • Bleeding within the skin (eg, purpura)

*Refer to the discussion of skin color in the [Primary Assessment](#) section in this Part.

Evaluate-Identify-Intervene

Use the evaluate-identify-intervene sequence ([Figure 15](#)) when caring for a seriously ill or injured child to help you determine the best treatment or intervention at any point. From the information gathered during your evaluation, identify the child's clinical condition by type and severity. Intervene with appropriate actions, and then repeat the sequence. This process is ongoing.



If at any point you identify a life-threatening problem, immediately activate the emergency response system (or send someone to do so) while you begin lifesaving interventions. Sometimes a child's condition may seem stable despite the presence of a life-threatening problem. Examples include a child who has ingested a toxin but is not yet showing effects and a trauma victim with internal bleeding who may initially maintain blood pressure by increasing heart rate and systemic vascular resistance. Reassess frequently.

Evaluate

Always verify the scene is safe and assess for potential environmental dangers before you evaluate the child, especially in out-of-hospital settings.

If no life-threatening condition is present, evaluate the child's condition by using these clinical assessment tools:

- Initial assessment: a first, quick, "from the doorway" observation of the child's appearance, breathing, and color, performed within the *first few seconds* after encountering the child
- Primary assessment: a rapid, hands-on ABCDE approach to evaluate respiratory, cardiac, and neurologic function; includes assessment of vital signs and pulse oximetry
- Secondary assessment: a focused medical history and a focused physical exam

Identify

Identify the type and severity of the child's problem ([Table 11](#)).

Table 11. Type and Severity of Potential Problems

Type	Severity
Respiratory	<ul style="list-style-type: none"> • Upper airway obstruction • Lower airway obstruction
	<ul style="list-style-type: none"> • Respiratory distress • Respiratory failure

	<ul style="list-style-type: none"> • Lung tissue disease • Disordered control of breathing 	
Circulatory	<ul style="list-style-type: none"> • Hypovolemic shock • Distributive shock • Cardiogenic shock • Obstructive shock 	<ul style="list-style-type: none"> • Compensated shock • Hypotensive shock
Cardiopulmonary failure		
Cardiac arrest		

The child's clinical condition can result from a combination of respiratory and circulatory problems. As a seriously ill or injured child deteriorates, one problem may lead to others, such as cardiopulmonary failure and cardiac arrest. Note that in the initial phase of your identification, you may be uncertain about the type or severity of problems.

Identifying the problem will help you determine the best initial interventions. Recognition and management are discussed in detail later in this manual.

Intervene

Once you identify the child's clinical condition, intervene with appropriate actions within your scope of practice. PALS interventions may include

- Positioning the child to maintain an open/patent airway
- Activating the emergency response system
- Starting CPR
- Obtaining the code cart and monitor
- Placing the child on a cardiac monitor and pulse oximeter
- Administering O₂
- Supporting ventilation
- Starting medications and fluids (eg, nebulizer treatment, IV/IO fluid bolus)

Continuous Sequence

Continue the evaluate-identify-intervene sequence until the child is stable. Use this sequence before and after each intervention to look for trends in the child's condition. For example, after you give O₂, reevaluate the child. Is the child breathing a little easier? Are color and mental status improving? After you give a fluid bolus to a child in hypovolemic shock, do heart rate and perfusion improve? Is another bolus needed? Use the evaluate-identify-intervene sequence whenever the child's condition changes.



Critical Concepts: The Evaluate-Identify-Intervene Sequence Is Continuous

Remember to repeat the evaluate-identify-intervene sequence until the child is stable:

- After each intervention
- When the child's condition changes or deteriorates

Primary Assessment



The primary assessment uses a hands-on ABCDE approach and includes assessment of the patient's vital signs (including oxygen saturation by pulse oximetry).

- Airway

- Breathing
- Circulation
- Disability
- Exposure

As you proceed through each component of the primary assessment, treat life-threatening abnormalities in real time before you complete the remainder of the primary assessment. In patients with life-threatening conditions evident in the primary assessment, correcting those conditions takes precedence over establishing baseline vital sign measures such as blood pressure or pulse oximetry. When you complete the primary assessment and after you address any life-threatening problems, proceed to the secondary assessment.



When you assess the airway, you determine if it is open/patent. To assess upper airway openness/patency:

- Look for chest or abdomen movement
- Listen for air movement and breath sounds
- Feel for air movement at the nose and mouth

To check for breathing, scan the victim's chest for rise and fall for no more than 10 seconds.

- If the victim is breathing, monitor the victim until additional help arrives.
- If the victim is not breathing or is only gasping, this is not considered normal breathing and is a sign of cardiac arrest.

To minimize delay in starting CPR, you may assess breathing at the same time as you check the pulse. This should take no more than 10 seconds.

The following signs suggest that the upper airway is obstructed:

- Increased inspiratory effort with retractions
- Abnormal inspiratory sounds (snoring or high-pitched stridor)
- Episodes where no airway or breath sounds are present despite respiratory effort (ie, complete upper airway obstruction)

If the upper airway is obstructed, determine if you can open and maintain the airway with *simple measures* or if you need *advanced interventions*.

Simple Measures to Maintain the Airway

Simple measures to open and maintain an open/patent upper airway may include one or more of the following:

Positioning

For a responsive child, allow the child to assume a position of comfort or elevate the head of the bed. For an unresponsive child, turn the child on her side if you do not suspect cervical injury or use a head tilt–chin lift or jaw thrust to improve airway patency.

Head Tilt–Chin Lift or Jaw Thrust

If you do not suspect cervical spine injury, use the head tilt–chin lift maneuver to open the airway. Avoid overextending the head/neck in infants because this may occlude, or block, the airway. If you suspect cervical spine injury (eg, the child has a head or neck injury), open the airway by using a jaw thrust without neck extension. If this does not open the airway, use a head tilt–chin lift or jaw thrust with neck extension because opening the airway is a priority. During CPR, stabilize the head and neck manually rather than with immobilization devices. Note that the jaw thrust may be used in children without trauma as well.

Suctioning

Suction the nose and oropharynx. Avoid overextending the head/neck in infants because this may occlude the airway.

Relief Techniques for Foreign-Body Airway Obstruction

If you suspect a child has aspirated a foreign body, has complete airway obstruction (is unable to make any sound), and is *responsive*, repeat the following as needed:

- Younger than 1 year: give 5 back slaps and 5 chest compressions
- 1 year or older: give abdominal thrusts

If at any time the child becomes unresponsive, activate (if you have a mobile device) or send someone to activate the emergency response system and begin CPR.


Airway Adjuncts

Use airway adjuncts to keep the tongue from falling back and obstructing the airway, but don't rely on an adjunct alone to maintain an open airway. You may still need to use a head tilt–chin lift. Assess the patient.

Advanced Interventions

Advanced interventions to maintain airway openness/patency may include one or more of the following:

- Endotracheal (ET) intubation or placing a laryngeal mask airway
- Applying continuous positive airway pressure or noninvasive ventilation
- Removing a foreign body; this intervention may require direct laryngoscopy (ie, visualizing the larynx with a laryngoscope)
- Cricothyrotomy (a needle puncture or surgical opening through the skin and cricothyroid membrane and into the trachea below the vocal cords)



Breathing

Assessing breathing includes evaluating

- Respiratory rate and pattern
- Respiratory effort
- Chest expansion and air movement
- Lung and airway sounds
- O₂ saturation by pulse oximetry

Normal Respiratory Rate

You can accomplish normal spontaneous breathing with minimal work, resulting in quiet breathing with unlabored inspiration and passive expiration. The normal respiratory rate is inversely related to age ([Table 12](#)): rapid in the neonate and decreased as the child gets older.

Table 12. Normal Respiratory Rates by Age*

Age	Rate (breaths/min)
Infant	30-53
Toddler	22-37
Preschooler	20-28
School-age child	18-25
Adolescent	12-20

*Consider the patient's normal range. The child's respiratory rate is expected to increase in the presence of fever or stress. Reproduced from Hazinski MF. Children are different. In: Hazinski MF, ed. *Nursing Care of the Critically Ill Child*. 3rd ed. Mosby; 2013:1-18, copyright Elsevier. Data from Fleming S, Thompson M, Stevens R, et al. *Lancet*.2011;377(9770):1011-1018.



Critical Concepts: Very Slow or Very Fast Respiratory Rate Is a Warning Sign

A consistent respiratory rate of less than 10 or more than 60 breaths per minute in a child of any age is often abnormal and warrants further assessment for the presence of a potentially serious condition.

Evaluate respiratory rate before your hands-on assessment because anxiety and agitation commonly alter the baseline rate. If the child has any condition that causes increased metabolic demand (eg, excitement, anxiety, exercise, pain, or fever), it may be appropriate for the respiratory rate to be higher than normal.

Determine the respiratory rate by counting the number of times the chest rises in 30 seconds and multiplying by 2. Normal sleeping infants may have irregular (periodic) breathing with pauses lasting up to 10 or even 15 seconds. If you count the number of times the chest rises for less than 30 seconds, you may estimate the respiratory rate inaccurately. Count the respiratory rate several times as you assess and reassess the child to detect changes. Alternatively, the respiratory rate may be displayed continuously on a monitor.

A decreasing respiratory rate from a rapid to a more “normal” rate may indicate overall improvement if it is associated with an improved level of consciousness and reduced signs of dyspnea and work of breathing. A decreasing or irregular respiratory rate in a child with a deteriorating level of consciousness, however, often indicates a worsening of the child’s clinical condition.

Abnormal Respiratory Rate and Pattern

Abnormal respirations include

- Irregular respiratory pattern
- Fast respiratory rate (tachypnea)
- Slow respiratory rate (bradypnea)
- Apnea

Irregular Respiratory Pattern

Children with neurologic problems may have irregular respiratory patterns. Examples of such irregular patterns are

- A deep gasping breath, followed by a period of apnea (no breathing or only gasping)
- A rapid respiratory rate, followed by periods of apnea or very shallow breaths

Irregular patterns such as these are serious and require urgent evaluation.

Fast Respiratory Rate

A *fast respiratory rate* (tachypnea) is a breathing rate that is faster than normal for age. It is often the first sign of respiratory distress in infants and can develop during periods of stress.

A fast respiratory rate may or may not be accompanied by signs of increased respiratory effort. A fast respiratory rate *without* signs of increased respiratory effort (ie, *quiet tachypnea*) may result from conditions that are not primarily respiratory in origin, such as

- High fever
- Pain
- Anemia
- Cyanotic congenital heart disease
- Metabolic acidosis
- Dehydration
- Sepsis (serious infection)

Slow Respiratory Rate

A slower than normal respiratory rate (bradypnea) may be caused by

- Respiratory muscle fatigue
- A central nervous system injury or problem that affects the respiratory control center
- Severe hypoxia

- Severe shock
- Hypothermia
- Drugs that depress the respiratory drive
- Some muscle diseases that cause muscle weakness



Critical Concepts: Bradypnea or Irregular Respiratory Rate Often Signal Impending Arrest

Bradypnea or an irregular respiratory rate in an acutely ill infant or child is an ominous clinical sign and often signals impending arrest.

Apnea

Apnea is when breathing stops, typically defined as longer than 20 seconds. Apnea may be further classified as **central** or **obstructive**, depending on whether inspiratory muscle activity is present. Central apnea indicates that the child is making no respiratory effort because of an abnormality or suppression of the brain or spinal cord, whereas obstructive apnea occurs when the passage of air is impeded, resulting in hypoxemia, hypercapnia, or both. Furthermore, in mixed apnea, periods of obstructive apnea and periods of central apnea occur.

Agonal gasps commonly happen in adults after sudden cardiac arrest and may be confused with normal breathing. They also appear late in the deterioration of a very sick child. Agonal gasps will not produce effective oxygenation and ventilation.

Increased Respiratory Effort

Increased respiratory effort results from conditions that increase resistance to airflow (eg, asthma or bronchiolitis) or that cause the lungs to be stiffer and difficult to inflate (eg, pneumonia, pulmonary edema, or pleural effusion). Nonpulmonary conditions that result in severe metabolic acidosis (eg, shock, DKA, salicylate ingestion, inborn errors of metabolism) can also cause increased respiratory rate and effort. Signs of increased respiratory effort reflect the child's attempt to improve oxygenation, ventilation, or both, so use these signs to assess the severity of the condition and determine the urgency of intervention. Signs of increased respiratory effort include

- Nasal flaring
- Retractions
- Head bobbing or seesaw respirations

Other signs are prolonged inspiratory or expiratory times, open-mouth breathing, gasping, and use of accessory muscles. Grunting is a serious sign and may indicate respiratory distress or respiratory failure. (Refer to [Grunting](#) later in this Part.)

Nasal Flaring

Nasal flaring is dilation of the nostrils with each inhalation to maximize airflow. It is most commonly observed in infants and younger children and is usually a sign of respiratory distress.

Retractions

Retractions are inward movements of the chest wall or tissues or sternum during inspiration. Chest retractions signify that the child is trying to move air into the lungs by using the chest muscles, but air movement is impaired by increased airway resistance or stiff lungs. Retractions may occur in several areas of the chest. The severity of the retractions generally corresponds with the severity of the child's breathing difficulty.

[Table 13](#) describes the location of retractions commonly associated with each level of breathing difficulty.

Table 13. Location of Retractions Commonly Associated With Each Level of Breathing Difficulty

Breathing difficulty	Location of retraction	Description
Mild to moderate	Subcostal	Retraction of the abdomen, just below the rib cage
	Substernal	Retraction of the abdomen at the bottom of the sternum
	Intercostal	Retraction between the ribs

Severe (may include the same retractions as seen with mild to moderate breathing difficulty)	Supraclavicular	Retraction in the tissues just above the collarbone
	Suprasternal	Retraction in the chest, just above the sternum
	Sternal	Retraction of the sternum toward the spine

Retractions accompanied by stridor or an inspiratory snoring sound suggest upper airway obstruction, while retractions accompanied by expiratory wheezing suggest marked lower airway obstruction (asthma or bronchiolitis), causing obstruction during both inspiration and expiration. Retractions accompanied by grunting or labored respirations suggest lung tissue disease. Severe retraction also may be accompanied by head bobbing or seesaw respirations.

Head Bobbing or Seesaw Respirations

Head bobbing and seesaw respirations often indicate that the child has increased risk for deterioration.

- Head bobbing is caused by using neck muscles to assist breathing. The child lifts the chin and extends the neck during inspiration and allows the chin to fall forward during expiration. It is most frequently seen in infants and can be a sign of respiratory failure.
- Seesaw respirations are present when the chest retracts and the abdomen expands during inspiration, and then the chest expands and the abdomen moves inward during expiration. In most children with neuromuscular disease, seesaw breathing is caused by weakness of the abdominal and chest wall muscles and by strong contraction of the diaphragm that dominates the weaker abdominal and chest wall muscles. Seesaw respirations usually indicate upper airway obstruction, but they also may be observed in severe lower airway obstruction, lung tissue disease, and states of disordered control of breathing. Seesaw respirations are characteristic of infants and children with neuromuscular weakness. This inefficient form of ventilation can quickly lead to fatigue.

Inadequate Respiratory Effort

When evaluating respiratory effort, look for signs that respiratory effort is inadequate and be prepared to support airway, oxygenation, and ventilation. These signs include

- Apnea
- Weak cry or cough
- Bradypnea
- Agonal gasps

Chest Expansion and Air Movement

Evaluate magnitude of chest wall expansion and air movement to assess adequacy of the child's tidal volume, the volume of air inspired with each breath. Normal tidal volume is approximately 5 to 7 mL/kg of ideal body weight throughout life. It is difficult to measure unless a child is mechanically ventilated, so your clinical assessment is very important.

Chest Wall Expansion

Chest expansion (chest rise) during inspiration should be symmetrical. It may be subtle during spontaneous quiet breathing, especially when clothing covers the chest, but chest expansion should be readily visible when the chest is uncovered. In normal infants, the abdomen may move more than the chest. Decreased or asymmetrical chest expansion may result from inadequate effort, airway obstruction, atelectasis, pneumothorax, hemothorax, pleural effusion, mucous plug, or foreign-body aspiration.

Air Movement

Auscultation for air movement is critical. Listen for the intensity of breath sounds and quality of air movement in the following areas:

- Anterior, located at mid-chest (just to the left and right of the sternum)
- Lateral, located under the armpits (the best location for evaluating air movement into the lower parts of the lungs)
- Posterior, located on both sides of the back

Because the chest is small and the chest wall is thin in infants and children, breath sounds readily transmit from one side of the chest to the other and from the upper airway. To evaluate distal air entry, listen below both axillae. Because

these areas are farther from the larger conducting airways, upper airway sounds are less likely to be transmitted.

Listen to the loudness of the air movement:

- Typical inspiratory sounds can be heard distally as soft, quiet noises occurring simultaneously with observed inspiratory effort.
- Normal expiratory breath sounds are often short and quieter; sometimes you may not even hear them.

Decreased chest excursion or decreased air movement during auscultation often accompanies poor respiratory effort. In the child with apparently normal or increased respiratory effort, diminished distal air entry suggests airflow obstruction or lung tissue disease. If the child's work of breathing and coughing suggest lower airway obstruction, but you cannot hear wheezes, the amount and rate of airflow may be insufficient to cause wheezing.

Distal air entry may be difficult to hear in the obese child, so it may be difficult to identify significant airway abnormalities in this population.

Minute Ventilation

Minute ventilation is the volume of air that moves into or out of the lungs each minute. It is the product of the number of breaths per minute (respiratory rate) and the volume of each breath (tidal volume).

Minute Ventilation = Respiratory Rate × Tidal Volume

Low minute ventilation (hypoventilation) may result from

- Slow respiratory rate
- Small tidal volume (ie, shallow breathing, high airway resistance, stiff lungs)
- Extremely rapid respiratory rate (resulting in very small tidal volumes)

Lung and Airway Sounds

During the primary assessment, listen for lung and airway sounds. Abnormal sounds include stridor, snoring, grunting, gurgling, wheezing, crackles, and change in cry/phonation/cough (including barking cough).

Stridor

Stridor is a coarse, usually higher-pitched breathing sound typically heard on inspiration, though you may hear it during both inspiration and expiration. Stridor is a sign of upper airway (extrathoracic) obstruction and may indicate that the obstruction is critical and requires immediate intervention.

Stridor has many causes of stridor, including

- A foreign body in the airway
- Infection (eg, croup)
- Congenital airway abnormalities (eg, laryngomalacia)
- Acquired airway abnormalities (eg, tumor or cyst)
- Upper airway edema (eg, allergic reaction or swelling after a medical procedure)

Snoring

Although *snoring* may be common in children during sleep, it also can be a sign of airway obstruction. Soft tissue swelling or decreased level of consciousness may cause airway obstruction and snoring.

Grunting

Grunting is typically a short, low-pitched sound heard during expiration that can sometimes be misinterpreted as a soft cry. It occurs as the child exhales against a partially closed glottis. Although grunting may accompany the response to pain or fever, infants and young children often grunt to help keep the small airways and alveolar sacs in the lungs open in an attempt to optimize oxygenation and ventilation.

Grunting often signifies lung tissue disease resulting from small airway collapse, alveolar collapse, or both. It may indicate progression of respiratory distress to respiratory failure. Pulmonary conditions that cause grunting include

- Pneumonia
- Pulmonary contusion
- Acute respiratory distress syndrome
- Cardiac conditions, such as congestive heart failure, that result in pulmonary edema

Grunting may be a sign of pain resulting from abdominal pathology (eg, bowel obstruction, perforated viscus, appendicitis, or peritonitis). It is typically a sign of severe respiratory distress or failure from lung tissue disease. Identify and treat the cause as quickly as possible, and be prepared to quickly intervene if the child's condition worsens.

Gurgling

Gurgling is a bubbling sound heard during inspiration or expiration that results from upper airway obstruction due to airway secretions, vomit, or blood.

Wheezing

Wheezing is a high-pitched or low-pitched whistling or sighing sound heard most often during expiration and less frequently during inspiration. This sound typically indicates lower (intrathoracic) airway obstruction, especially of the smaller airways. Common causes of wheezing are bronchiolitis and asthma. Isolated inspiratory wheezing suggests a foreign body or other cause of partial obstruction of the trachea or upper airway.

Crackles

Crackles, also known as *rales*, are sharp, crackling inspiratory sounds. Dry crackles sound similar to the sound made when you rub several hairs together close to your ear. Dry crackles are more often heard with atelectasis (small airway collapse) and interstitial lung disease. Moist crackles indicate accumulation of alveolar fluid and are typically associated with lung tissue disease (eg, pneumonia and pulmonary edema) or interstitial lung disease. Note that you may not hear crackles despite the presence of pulmonary edema.

Change in Cry/Phonation/Cough (Including Barking Cough)

If an infant's cry becomes very soft with only short sounds during expiration (ie, so the cry sounds more like the soft "mewing" of a cat) or an older child begins to talk in short phrases or single words instead of sentences, this may indicate severe respiratory distress and shortness of breath. If an infant or child develops a "barking" cough or change in pitch of cry or voice, this may indicate upper airway obstruction.

Oxygen Saturation by Pulse Oximetry

The O_2 saturation is the percent of total hemoglobin that is fully saturated with oxygen (ie, the oxyhemoglobin saturation). This oxyhemoglobin saturation does not indicate the amount of O_2 delivered to the tissues. O_2 delivery is the product of arterial O_2 content (oxygen bound to hemoglobin plus dissolved O_2) and cardiac output. It is also important to note that O_2 saturation does not provide information about effectiveness of ventilation (CO_2 elimination).

Pulse oximetry monitors the percentage of the child's hemoglobin that is fully saturated with oxygen. The pulse oximeter consists of a probe that is linked to a monitor and attached to the child's finger, toe, or earlobe. It must detect a consistent pulsatile signal. The unit displays the calculated percentage of saturated hemoglobin. Most units make an audible sound for each pulse beat and display the heart rate, while some models display the quality of the pulse signal as a waveform or with bars.

Because pulse oximetry can indicate low O_2 saturation (hypoxemia) before it causes cyanosis or bradycardia, providers can use pulse oximetry to monitor trends in O_2 saturation in response to treatment. If available, continuously monitor pulse oximetry for a child in respiratory distress or failure during stabilization, transport, and post-cardiac arrest care.

An O_2 saturation (SpO_2) of 94% or more while a child is breathing room air usually indicates that oxygenation is adequate; conversely, an SpO_2 less than 94% when the child is breathing room air indicates hypoxemia. Consider administering supplemental O_2 if the O_2 saturation is below this value in a critically ill or injured child. An SpO_2 of less than 90% in a child receiving 100% O_2 is an indication for additional intervention. The pulse oximeter probe has 2 parts that must be placed opposite each other, so they are located on either side of a pulsatile tissue bed. Lights of different wavelengths originate from one side of the probe, and the light is captured on the other side of the tissue by the other side of the probe. A processor in the oximeter calculates the percent of each light that has been absorbed by the tissues. Hemoglobin that is fully saturated with oxygen absorbs light differently than hemoglobin that is not fully saturated with oxygen. By determining the absorption of the different wavelengths of light, the pulse oximeter can estimate the percent of hemoglobin that is fully saturated with blood.

Errors in pulse oximetry can occur if the probe is not placed across an area of pulsatile blood flow (ie, the pulses are very weak or poorly perfused, or the probe is placed so the side emitting light is not directly across from the side that captures the light). Bright light in the room can interfere with accurate light absorption detection. In addition, a pulse oximeter will report falsely high values in the presence of carbon monoxide poisoning because hemoglobin bound by

carbon monoxide (as in carbon monoxide poisoning) absorbs light in a way that is very similar to hemoglobin that is fully saturated with oxygen (refer to [Caution in Interpreting Oximetry Readings](#) section).

Caution in Interpreting Pulse Oximetry Readings

Be careful to interpret pulse oximetry readings in conjunction with your clinical assessment, including considering signs such as respiratory rate, respiratory effort, and level of consciousness. A child may be in respiratory distress yet maintain normal O₂ saturation by increasing respiratory rate and effort, especially if receiving supplemental O₂.

If the pulse oximeter displays a different heart rate from the one determined by electrocardiographic (ECG) monitoring, the displayed O₂ saturation is not reliable. *When the pulse oximeter does not detect a consistent pulse or there is an irregular or poor waveform, the child may have poor distal perfusion and the pulse oximeter may not be accurate—check the child and intervene as needed.* The pulse oximeter may not be accurate if the child develops severe shock and won't be accurate during cardiac arrest.

Pulse oximetry indicates only O₂ saturation and not O₂ delivery. For example, if the child is profoundly anemic (hemoglobin is very low), the saturation may be 100%, but O₂ content in the blood and O₂ delivery may be low.

The pulse oximeter does not accurately recognize methemoglobin or carboxyhemoglobin (hemoglobin bound to carbon monoxide). If carboxyhemoglobin (from carbon monoxide poisoning) is present, the pulse oximeter will reflect a falsely high O₂ saturation because it counts the carboxyhemoglobin as fully saturated hemoglobin. If methemoglobin concentrations are above 5%, the pulse oximeter will read approximately 85% regardless of the degree of methemoglobinemia. If you suspect either of these conditions, obtain a blood gas and send it for laboratory analysis of O₂ saturation measurement by co-oximeter.

Signs of probable respiratory failure include

- Very rapid or inadequate respiratory rate; possible apnea
- Significant, inadequate, or absent respiratory effort
- Absent distal air movement
- Extreme tachycardia; bradycardia often indicates life-threatening deterioration
- Low oxygen saturation (hypoxemia) despite high-flow supplemental oxygen
- Decreased level of consciousness
- Cyanosis
- Seesaw breathing
- Head bobbing



Circulation is assessed by evaluating

- Heart rate and rhythm
- Pulses (both peripheral and central)
- Capillary refill time
- Skin color and temperature
- Blood pressure

Adequate circulation is also reflected by urine output and level of consciousness. Urine output can indirectly indicate kidney perfusion. Normal urine output requires adequate blood flow and hydration. Normal values for urine output are age dependent:

- Infants and young children: 1.5 to 2 mL/kg per hour normal urine output
- Older children and adolescents: 1 mL/kg per hour normal urine output
- Children with shock: decreased urine output

Accurately measuring urine output in all critically ill or injured children requires an indwelling catheter. Initial urine output is not a reliable indicator of the child's clinical condition because much of the urine may have been produced before the onset of symptoms. An increase in urine output is a good indicator of positive responses to therapy.

For more information on assessing level of consciousness, refer to the [Disability](#) section later in this Part.

Heart Rate and Rhythm

To determine heart rate, check the pulse rate, listen to the heart, or view a monitor display of the ECG or pulse oximeter waveform. The heart rate should be appropriate for the child's age, level of activity, and clinical condition (Table 14). Note the wide range of normal heart rates. For example, a child who is sleeping or is athletic may have a heart rate lower than the normal range for age.

Table 14. Normal Heart Rates*

Age	Awake rate (/min)	Sleeping rate (/min)
Neonate	100-205	90-160
Infant	100-180	90-160
Toddler	98-140	80-120
Preschooler	80-120	65-100
School-age child	75-118	58-90
Adolescent	60-100	50-90

*Always consider the patient's normal range and clinical condition. Heart rate will normally increase with fever or stress.

Reproduced from Hazinski MF. Children are different. In: Hazinski MF, ed. *Nursing Care of the Critically Ill Child*. 3rd ed. Mosby; 2013:1-18, copyright Elsevier.

Consider the following when evaluating the heart rate and rhythm in any seriously ill or injured child:

- The child's typical heart rate and baseline rhythm
- The child's level of activity and clinical condition
- The child's cardiac function and perfusion

Children with congenital heart disease may have conduction abnormalities. Consider the child's baseline ECG when interpreting heart rate and rhythm. Children with poor cardiac function are more likely to be symptomatic from cardiac rhythm disturbances (arrhythmias) than are children with normal cardiac function.

The heart rhythm is typically regular with only small fluctuations in rate. When checking the heart rate, assess for abnormalities in the monitored ECG. Arrhythmias result from abnormalities in, or insults to, the cardiac conduction system or heart tissue. Arrhythmias also can result from shock or hypoxia. In the advanced life support setting, an arrhythmia in a child can be broadly classified according to the observed heart rate or effect on perfusion, with slow heart rate classified as bradycardia, fast heart rate classified as tachycardia, and absent heart rate classified as cardiac arrest.

Bradycardia is a heart rate slower than normal for a child's age and clinical condition. Slight bradycardia may be normal in athletic children, but a very slow rate in a child with other symptoms is worrisome and may indicate that cardiac arrest is imminent. Hypoxia is the most common cause of bradycardia in children. If a child with bradycardia has signs of cardiopulmonary compromise, immediately support oxygenation and ventilation with a bag and mask and administer supplemental O₂. Start chest compressions if the heart rate remains less than 60/min with signs of cardiopulmonary compromise despite adequate oxygenation and ventilation. If the child with bradycardia is alert and has no signs of cardiopulmonary compromise, consider other causes of a slow heart rate, such as heart block or drug overdose.

Tachycardia is a resting heart rate that is faster than the normal range for a child's age and clinical condition. Sinus tachycardia, a common, nonspecific response to a variety of conditions, is often appropriate when the child is anxious, crying, febrile, or seriously ill or injured. To determine if the tachycardia is a sinus tachycardia or represents a cardiac rhythm disturbance, evaluate the child's history, clinical condition, and ECG. Any tachycardia associated with signs of circulatory compromise, including hypotension, altered mental status, or signs of shock, requires immediate evaluation and intervention.



Critical Concepts: Tachycardia Can Be a Sign of a Serious Condition

A heart rate that is greater than 180/min in an infant or toddler and greater than 160/min in a child older than 2 years warrants further assessment and may be a serious condition.

For more information, refer to [Part 11: Recognizing Arrhythmias](#).

Relationship of Breathing to Heart Rhythm

In healthy children, the heart rate may fluctuate with the respiratory cycle, increasing with inspiration and slowing down with expiration. This condition is called *sinus arrhythmia*. Note if the child has an irregular rhythm that is not related to breathing. An irregular rhythm may indicate an underlying rhythm disturbance, such as premature ventricular or atrial contractions or an atrioventricular block.

Pulses

Evaluating central and peripheral pulses is critical to the assessment of systemic perfusion in an ill or injured child. Central pulses are ordinarily stronger than peripheral pulses because they are present in vessels of larger size that are located closer to the heart. An exaggerated difference in quality between central and peripheral pulses occurs when peripheral vasoconstriction is associated with shock. Central pulses include femoral, brachial (in infants), carotid (in older children), and axillary. Peripheral pulses include radial, dorsalis pedis, and posterior tibial. These pulses are easily palpable in healthy infants and children (unless the child is obese or the ambient temperature is cold).



Critical Concepts: Weakening of Pulses as Perfusion Decreases

Weak central pulses are worrisome and indicate the need for rapid intervention to prevent cardiac arrest. When cardiac output decreases in shock, systemic perfusion decreases incrementally. The decrease in perfusion starts in the extremities with a decrease in intensity of pulses and then an absence of peripheral pulses. As cardiac output and perfusion decrease further, central pulses eventually weaken.

A cold environment can cause vasoconstriction and a discrepancy between peripheral and central pulses. However, if cardiac output remains adequate, central pulses should remain strong.

Beat-to-beat fluctuation in pulse volume may occur in children with arrhythmias (eg, premature atrial or ventricular contractions). Fluctuation in pulse volume with the respiratory cycle (pulsus paradoxus) can occur in children with severe asthma or pericardial tamponade. In an intubated child receiving positive-pressure ventilatory support, a reduction in pulse volume with each positive-pressure breath may indicate hypovolemia.

Capillary Refill Time

Capillary refill time is the time it takes for blood to return to tissue blanched by pressure. It increases as skin perfusion decreases. Note that normal capillary refill time is 2 seconds or less, and a prolonged capillary refill time may indicate low cardiac output.

Evaluate capillary refill in a neutral thermal environment (ie, room temperature) by lifting the extremity slightly above the level of the heart, pressing on the skin, and rapidly releasing the pressure. Note how many seconds it takes for the area to return to its baseline color.

Common causes of sluggish, delayed, or prolonged capillary refill (a refill time of greater than 2 seconds) are dehydration, shock, and hypothermia. Note that shock can be present despite a normal (or even brisk) capillary refill time. Children with septic shock (refer to [Part 9: Recognizing Shock](#)) may have warm skin and extremities with very rapid (ie, less than 2 seconds) capillary refill time (often called *flash capillary refill*) despite the presence of shock.

Skin Color and Temperature

Monitor changes in skin color, temperature, and capillary refill time to assess a child's perfusion and response to therapy. Normal skin color and temperature should be consistent over the trunk and extremities. The mucous membranes, nail beds, palms of the hands, and soles of the feet should be pink. When perfusion deteriorates and O₂ delivery to the tissues becomes inadequate, the hands and feet are typically affected first and become cool, pale, dusky, or mottled. If perfusion becomes worse, the skin over the trunk and extremities may undergo similar changes.

Consider the temperature of the child's surroundings (ie, ambient temperature) when evaluating skin color and temperature. If the ambient temperature is cool, peripheral vasoconstriction may produce mottling or pallor with cool skin and delayed capillary refill, particularly in the extremities. These changes develop despite normal cardiovascular function.

To assess skin temperature, use the back of your hand because it is more sensitive to temperature changes than the palm, which has thicker skin. Slide the back of your hand up the extremity to determine a line where the skin changes

from cool to warm, and monitor this line of demarcation between warm and cool skin over time to determine the child's response to therapy. The line should move distally as the child improves.

Carefully monitor for the following skin findings ([Table 15](#)), which may indicate inadequate O₂ delivery to the tissues:

- Pallor
- Mottling
- Cyanosis

Table 15. Skin Findings, Location, and Causes

Skin color	Location	Causes
Pallor (paleness; lack of normal color)	Skin or mucous membranes	<ul style="list-style-type: none"> • Normal skin color • Decreased blood supply to the skin (cold; stress; shock, especially hypovolemic and cardiogenic) • Decreased number of red blood cells (anemia) • Decreased skin pigmentation • Poor perfusion
Mottling (irregular or patchy discoloration)	Skin	<ul style="list-style-type: none"> • Normal distribution of skin melanin • Intense vasoconstriction from irregular supply of oxygenated blood to the skin due to hypoxemia, hypovolemia, or shock
Cyanosis (blue discoloration)	Skin or mucous membranes	
Acrocyanosis (bluish discoloration)	Hands and feet and around the mouth (ie, the skin around the lips)	Normal in the newly born
Peripheral cyanosis (bluish discoloration)	Hands and feet (beyond newborn period)	<ul style="list-style-type: none"> • Shock • Congestive heart failure • Peripheral vascular disease • Conditions causing venous stasis
Central cyanosis (blue color)	Lips and other mucous membranes	<ul style="list-style-type: none"> • Low ambient O₂ tension (eg, high altitude) • Alveolar hypoventilation (eg, traumatic brain injury, drug overdose) • Diffusion defect (eg, pneumonia) • Ventilation/perfusion imbalance (eg, asthma, bronchiolitis, acute respiratory distress syndromes) • Intracardiac shunt (eg, cyanotic congenital heart disease)

You must interpret pallor within the context of other signs and symptoms because it is not necessarily abnormal and can result from lack of exposure to sunlight or inherited paleness. Pallor is often difficult to detect in a child with dark skin and in those with thick skin and variation in the vascularity of subcutaneous tissue. However, family members often can tell you if a child's color is abnormal. Central pallor (ie, pale color of the lips and mucous membranes) strongly suggests anemia or poor perfusion. Pallor of the mucous membranes (the lips, lining of the mouth, tongue, lining of the eyes) and pale palms and soles are more likely to be clinically significant.

- Areas that are mottled may appear as an uneven combination of pink, bluish gray, or pale skin tones.
- Blood saturated with O₂ is bright red, whereas unoxygenated blood is dark bluish-red. The location of cyanosis is important.
- Acrocyanosis is a common normal neonatal finding in the first 24 to 48 hours after delivery.
- Peripheral cyanosis can be caused by diminished O₂ delivery to the tissues.
- Central cyanosis is a blue color of the lips and other mucous membranes.



Critical Concepts: Variability in Appearance of Central Cyanosis

Cyanosis is not apparent until at least 5 g/dL of hemoglobin are desaturated (not bound to O₂). The O₂ saturation at which a child will appear cyanotic depends on the child's hemoglobin concentration. For example, in a child with a hemoglobin concentration of 16 g/dL, cyanosis will appear at an O₂ saturation of approximately 70% (ie, 30% of the hemoglobin, or 5 g/dL, is desaturated). If the hemoglobin concentration is low (eg, 8 g/dL), a very low O₂ saturation (eg, less than 40%) is required to produce cyanosis. Thus, cyanosis may be apparent with a milder degree of

hypoxemia in a child with polycythemia (increased amount of hemoglobin and red blood cells, eg, in cyanotic heart disease) but may not be apparent despite significant hypoxemia if the child is anemic.

Cyanosis may be more obvious in the mucous membranes and nail beds than in the skin, particularly if the skin is dark. It also can be seen on the soles of the feet, tip of the nose, and earlobes. Children with different hemoglobin levels will be cyanotic at different levels of O₂ saturation; cyanosis is more readily detected at higher O₂ saturations if the hemoglobin level is high. The development of central cyanosis typically indicates the need for emergency intervention, such as O₂ administration and ventilatory support.

Blood Pressure

Accurate blood pressure measurement requires a cuff that extends at least 50% to 75% of the length of the upper arm (from the axilla to the antecubital fossa) and a cuff bladder that covers about 40% of the mid–upper arm circumference.

Normal Blood Pressures

Table 16 summarizes the normal range of systolic and diastolic blood pressures according to age from 1 standard deviation below to 1 standard deviation above the mean in the first year of life and from the 50th to 95th percentile (assuming the 50th percentile for height) for children 1 year or older. As with heart rates, there is a wide range of values within the normal range, and normal blood pressures may fall outside the ranges listed here.

Table 16. Normal Blood Pressures

Age	Systolic pressure (mm Hg)*	Diastolic pressure (mm Hg)*	Mean arterial pressure (mm Hg) [‡]
Birth (12 hours, <1000 g)	39-59	16-36	28-42 [‡]
Birth (12 hours, 3 kg)	60-76	31-45	48-57
Neonate (96 hours)	67-84	35-53	45-60
Infant (1-12 months)	72-104	37-56	50-62
Toddler (1-2 years)	86-106	42-63	49-62
Preschooler (3-5 years)	89-112	46-72	58-69
School-age child (6-9 years)	97-115	57-76	66-72
Preadolescent (10-12 years)	102-120	61-80	71-79
Adolescent (12-15 years)	110-131	64-83	73-84

*Systolic and diastolic blood pressure ranges assume 50th percentile for height for children 1 year and older.

[‡]Mean arterial pressures (diastolic pressure + [difference between systolic and diastolic pressure/3]) for 1 year and older, assuming 50th percentile for height.

[‡]Approximately equal to postconception age in weeks (may add 5 mm Hg).

Modified from Hazinski MF. Children are different. In: Hazinski MF, ed. *Nursing Care of the Critically Ill Child*. 3rd ed. Mosby; 2013:1-18, copyright Elsevier. Data from Gemelli M, Manganaro R, Mami C, De Luca F. Longitudinal study of blood pressure during the 1st year of life. *Eur J Pediatr*. 1990;149(5):318-320; Versmold HT, Kitterman JA, Phibbs RH, Gregory GA, Tooley WH. Aortic blood pressure during the first 12 hours of life in infants with birth weight 610 to 4,220 grams. *Pediatrics*. 1981;67(5):607-613; Haque IU, Zaritsky AL. Analysis of the evidence for the lower limit of systolic and mean arterial pressure in children. *Pediatr Crit Care Med*. 2007;8(2):138-144; and National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. *The Fourth Report on the Diagnosis, Evaluation, and Treatment of High Blood Pressure in Children and Adolescents*. National Heart, Lung, and Blood Institute; 2005. NIH publication 05-5267.

Hypotension

Hypotension is defined by the thresholds of systolic blood pressure shown in Table 17.

Table 17. Definition of Hypotension by Systolic Blood Pressure and Age

Age	Systolic blood pressure (mm Hg)
Term neonates (0-28 days)	<60
Infants (1-12 months)	<70
Children 1-10 years	<70 + (age in years × 2) (this estimates systolic blood pressure that is less than the fifth blood pressure percentile for age)*

*This fifth percentile is a systolic blood pressure that is lower than all but 5% of normal children (ie, it will be hypotensive for 95% of normal children).

Note that these blood pressure thresholds approximate just above the fifth percentile systolic blood pressures for age, so they will overlap with normal blood pressure values for 5% of healthy children. An observed decrease in systolic blood pressure of 10 mm Hg from baseline should prompt serial evaluations for additional signs of shock. In addition, remember that these threshold values were established in normal, resting children. Children with injury and stress typically have increased blood pressure, and a blood pressure in the low-normal range may be abnormal in a seriously ill child.

When hypotension develops in a child with shock, physiologic compensatory mechanisms (eg, tachycardia and vasoconstriction) have failed. Hypotension in acute hemorrhage may be present with an acute loss of 20% of the circulating blood volume, but it is almost always present when the loss exceeds 30%. Hypotension in septic shock can occur from loss of intravascular volume and inappropriate vasodilation or severe vasoconstriction and inadequate cardiac output/cardiac index.

The development of bradycardia in a child with hypotension and poor perfusion is an ominous sign. To prevent cardiac arrest, manage airway and breathing and support adequate intravascular volume, cardiac function, and perfusion are needed to prevent cardiac arrest.



Disability

Perform the disability assessment to quickly evaluate neurologic function, using one of several tools to evaluate responsiveness and level of consciousness. Perform this evaluation at the end of the primary assessment, and repeat it during the secondary assessment to monitor for changes in the child's neurologic status.

Clinical factors that reflect brain perfusion can provide indirect evidence of circulatory function in the ill or injured pediatric patient. These signs include level of consciousness and TICLS. Signs of inadequate O₂ delivery to the brain correlate with the severity and duration of cerebral hypoxia.

Sudden and severe cerebral hypoxia may cause the following neurologic signs:

- Decreased level of consciousness
- Loss of muscular tone
- Generalized seizures
- Pupil dilation

You may observe other neurologic signs when cerebral hypoxia develops gradually. Because these signs can be subtle, repeat measurements over time to best detect them:

- Decreased level of consciousness with or without confusion
- Irritability
- Lethargy
- Agitation alternating with lethargy

Standard evaluations include

- AVPU (Alert, Responsive to Voice, Responsive to Pain, Unresponsive) Pediatric Response Scale
- Glasgow Coma Scale (GCS)
- Pupil response to light
- Blood glucose test

AVPU Pediatric Response Scale

To rapidly evaluate cerebral cortex function, use the AVPU Pediatric Response Scale. This scale is a system for rating a child's level of consciousness, an indicator of cerebral cortex function. The scale consists of 4 ratings:

- **Alert:** The child is awake, active, and appropriately responsive to caregivers and external stimuli. Assess “appropriate response” in terms of the anticipated response based on the child’s age and/or developmental level and the setting or situation.
- **Voice:** The child responds only to voice (eg, calling the child’s name or speaking loudly).
- **Painful:** The child responds only to a painful stimulus, such as a sternal rub or pinching the trapezius.
- **Unresponsive:** The child does not respond to any stimulus.

Decreased level of consciousness in children can be caused by

- Poor cerebral perfusion
- Severe shock
- Traumatic brain injury
- Seizure activity
- Encephalitis, meningitis
- Hypoglycemia
- Drugs
- Hypoxemia
- Hypercarbia

Altered mental status refers to the range of mental states from agitation to coma. For a pediatric patient with altered mental status, consider hypoglycemia and evaluate blood glucose as soon as possible.

If an ill or injured child has decreased responsiveness, immediately assess oxygenation, ventilation, perfusion, and blood glucose.

Glasgow Coma Scale Overview

The GCS is the most widely used method of evaluating a child’s level of consciousness and neurologic status. The *best* eye opening (E), verbal (V), and motor (M) responses are individually scored ([Table 18](#)) and then added together to produce the GCS score.

Table 18. Glasgow Coma Scale

Eye opening		Best motor response		Best verbal response	
4	Spontaneous	6	Obeys commands	5	Oriented
3	To speech	5	Localizes pain	4	Confused
2	To pain	4	Withdraws from pain	3	Inappropriate words
1	No response	3	Abnormal flexion	2	Incomprehensible words
		2	Abnormal extension	1	No response
		1	No response		

Teasdale G, Jennett B. Assessment of coma and impaired consciousness: a practical scale. *Lancet*. 1974;2(7872):81-84.

For example, a person who has spontaneous eye opening (E=4), is fully oriented (V=5), and is able to follow commands (M=6) is assigned a GCS score of 15, the highest possible score. A person with no eye opening (E=1), no verbal response (V=1), and no motor response (M=1) to a painful stimulus is assigned a GCS score of 3, the lowest possible score.

Severity of head injury is categorized into 3 levels based on GCS score after initial resuscitation:

- Mild head injury: GCS score 13 to 15
- Moderate head injury: GCS score 9 to 12
- Severe head injury: GCS score 3 to 8

Glasgow Coma Scale Scoring

The GCS has been modified for preverbal or nonverbal children ([Table 19](#)). While scores for eye opening are essentially the same as in the standard GCS, the best motor and verbal response scores were adapted to accommodate age-appropriate responses.

Important: When using the GCS or its pediatric modification, record the individual components of the score. If the patient is intubated, unconscious, or preverbal, the most important part of this scale is motor response. Providers

should carefully evaluate this component.

Table 19. Pediatric Glasgow Coma Scale*

Score	Child	Infant
Eye opening		
4	Spontaneously	Spontaneously
3	To verbal command	To shout, speech
2	To pain	To pain
1	No response	No response
Best motor response		
6	Obeys commands	Spontaneous movements
5	Localizes pain	Withdraws to touch
4	Flexion-appropriate withdraw	Flexion-appropriate withdraw
3	Flexion-abnormal (decorticate rigidity)	Flexion-abnormal (decorticate rigidity)
2	Extension (decerebrate rigidity)	Extension (decerebrate rigidity)
1	No response	No response
Best verbal response		
5	Oriented and converses	Smiles, coos, and babbles
4	Disoriented, confused	Cries but is consolable
3	Inappropriate words	Persistent, inappropriate crying and/or screaming
2	Incomprehensible sounds	Moans, grunts to pain
1	No response	No response
Total = 3 to 15		

*Score is the sum of the individual scores from eye opening, best motor response, and best verbal response, using age-specific criteria. GCS score of 13 to 15 indicates mild head injury; GCS score of 9 to 12 indicates moderate head injury; and GCS score of ≤ 8 indicates severe head injury.

Modified from James HE, Trauner DA. The Glasgow Coma Score and Modified Coma Score for Infants. In: James HE, Anas NG, Perkin RM, eds. *Brain Injuries in Infants and Children: Pathophysiology and Management*. Grune & Stratton Inc; 1985:179-182, copyright Elsevier.

The best disability scale for an individual child may be site specific. For example, the AVPU scale may be appropriate in the prehospital setting, whereas the GCS (particularly the motor component) or pediatric GCS may be better in the emergency department and hospital. The differences between the AVPU scale and GCS or pediatric GCS do not appear significant when associated with neurologic outcome. Each component of the AVPU scale generally correlates with the GCS scores shown in [Table 20](#).

Table 20. AVPU Scale and Glasgow Coma Scale Equivalents





Response	GCS score
Alert	15
Verbal	13
Painful stimulation	8
Unresponsive to noxious stimulation	6

Pupil Response to Light

Healthcare providers also should assess and record the pupillary size and response to light for each eye in any patient with altered level of consciousness. Pupil response to light is a useful indicator of brainstem function. Normally, pupils constrict in response to light and dilate in a dark environment, so if the pupils don't constrict in response to direct light (eg, flashlight directed at the eyes), suspect a brainstem injury. The pupils are typically equal in size, although slight variations are normal. Irregularities in pupil size or response to light may occur as a result of ocular

trauma or other conditions, such as increased intracranial pressure. Refer to [Table 21](#) for examples of abnormal pupil responses and their possible causes.

Table 21. Abnormal Pupil Responses and Possible Causes

Abnormal pupil response	Possible cause
Pinpoint pupils 	<ul style="list-style-type: none"> Narcotic ingestion (eg, opioid)
Dilated pupils 	<ul style="list-style-type: none"> Predominant sympathetic autonomic activity Sympathomimetic ingestion (eg, cocaine) Anticholinergic ingestion (eg, local or systemic atropine) Increased intracranial pressure
Unilaterally dilated pupils 	<ul style="list-style-type: none"> Inadvertent topical absorption of a breathing treatment (eg, ipratropium) Dilating eye drops
Unilaterally dilated pupils with altered mental status 	<ul style="list-style-type: none"> Ipsilateral (same side) uncal herniation (lateral herniation of the temporal lobe, caused by increased intracranial pressure)

During the disability assessment, assess and record the following for each eye:

- Size of pupils (in millimeters)
- Equality of pupil size
- Constriction of pupils to light (ie, the magnitude and rapidity of the response to light)

The acronym PERRL (**P**upils **E**qual, **R**ound, **R**eactive to **L**ight) describes the normal pupil responses to light.

Blood Glucose Test

Hypoglycemia refers to blood glucose 40 mg/dL or less in the newly born and 60 mg/dL or less in a child. It may result in brain injury if not recognized and effectively treated. Base treatment decisions on patient symptoms, and potentially include oral glucose. Monitor the blood glucose concentration of any seriously ill infant or child. A low blood glucose concentration may cause altered level of consciousness or even brain injury if it is not quickly identified and adequately treated. Measure the blood glucose concentration with a point-of-care glucose test.

For more information about the recognizing and treating hypoglycemia, refer to the [Glucose](#) section in [Part 10](#).



Exposure

Exposure, or undressing the seriously ill or injured child to perform a focused physical examination, is the final component of the primary assessment. Remove clothing one area at a time to carefully observe the child's face and head, trunk (front and back), extremities, and skin. Maintain cervical spine precautions when turning any child with a suspected neck or spine injury. If necessary, use blankets to keep the child comfortable and warm and, if available, heating lamps to prevent cold stress or hypothermia. Assess core temperature, noting any difference in warmth between trunk and extremities. Identify the presence of fever, which may indicate infection and early need for antibiotics (eg, sepsis).

During this part of the examination, look for evidence of trauma, such as bleeding, burns, or unusual markings that suggest nonaccidental trauma. Such signs include bruises in different stages of healing, injuries that don't correlate with the child's history, and delay from time of injury until the child receives medical attention.

Look for the presence and progression of petechiae and purpura (nonblanching purple discolorations in the skin caused by bleeding from capillaries and small vessels). Petechiae appear as tiny red dots and suggest a low platelet count, whereas purpura appears as larger areas. Both petechiae and purpura may signify septic shock. Also look for other rashes that may suggest shock (eg, hives in anaphylactic shock).

Look for signs of injury to the extremities, including deformities or bruising. Palpate the extremities and note the child's response. If tender, suspect injury; if necessary, immobilize the extremity.

Secondary Assessment

Secondary assessment consists of a focused history and detailed physical examination with ongoing reassessment of physiologic status and response to treatment.

H's and T's

The most common causes of cardiac arrest are presented as H's and T's:

- *Hypovolemia*
- *Hypoxia*
- *Hydrogen ion (acidosis)*
- *Hypoglycemia*
- *Hypo-/hyperkalemia*
- *Hypothermia*
- *Tension pneumothorax*
- *Tamponade (cardiac)*
- *Toxins*
- *Thrombosis (pulmonary)*
- *Thrombosis (coronary)*

Focused History

Obtain a focused history to gather information about the patient and the incident, particularly information that might help explain impaired respiratory or cardiovascular function. You can use the SAMPLE mnemonic, a systematic method for gathering information on a sick child, as a memory aid for obtaining a focused history. Obtain an accurate timeline for all signs, symptoms, and events leading up to the current presentation. Gather the following information for each category:

- Signs and symptoms at onset of illness, such as
 - Breathing difficulty (eg, cough, rapid breathing, increased respiratory effort, breathlessness, abnormal breathing pattern, chest pain on deep inhalation), wheezing
 - Tachypnea
 - Tachycardia
 - Diaphoresis
 - Decreased level of consciousness, fatigue
 - Agitation, anxiety
 - Fever, headache
 - Decreased oral intake
 - Diarrhea, vomiting, abdominal pain
 - Bleeding
 - Duration of symptoms
- Allergies
 - Medications, foods, latex, etc
 - Associated reactions
- Medications
 - Patient medications (eg, over-the-counter, vitamins, inhalers, herbal supplements), medications that can be found in the child's environment
 - Last dose and time of recent medications
- Past medical history

- Health history (eg, premature birth, previous illnesses, hospitalizations)
- Significant underlying medical problems (eg, asthma, chronic lung disease, congenital heart disease, arrhythmia, congenital airway abnormality, seizures, head injury, brain tumor, diabetes, hydrocephalus, neuromuscular disease)
- Past surgeries
- Immunization status
- *Last meal*
 - Time and nature of last intake of liquid or food (including breast or bottle feeding in infants)
 - Elapsed time between last meal and presentation of current illness can affect treatment and management of the condition (eg, possible anesthesia, possible intubation)
- *Events*
 - Events leading to current illness or injury (eg, onset sudden or gradual, type of injury)
 - Hazards at scene

Treatment during interval from onset of disease or injury until evaluation, estimated time of onset (if out-of-hospital onset)

Focused Physical Examination

Next, perform a focused physical examination based on the severity of the child's illness or injury. Carefully assess the primary area of concern of the illness or injury (ie, respiratory assessment with respiratory distress) and perform a brief head-to-toe evaluation. Some examples of areas to assess for certain illnesses include

- The nose/mouth (signs of obstruction, nasal congestion, stridor, mucosal edema), chest/lungs, heart (tachycardia, gallop, or murmur), and level of consciousness (somnolence secondary to hypercarbia, anxiety secondary to hypoxia) for respiratory distress
- The heart (gallop or murmur), lungs (crackles, difficulty breathing, intolerance of supine position), abdomen (evidence of hepatomegaly consistent with right heart failure), and extremities (peripheral edema) for suspected heart failure or arrhythmias
- The abdomen and back for trauma

Ongoing Reassessment

Ongoing reassessment of all patients is essential to evaluate the response to treatment and to track the progression of identified physiologic and anatomic problems. Apply this reassessment in real time as needed based on the child's clinical condition through all phases of assessment. Do not limit it to the last part of the assessment sequence. You may also identify new problems on reassessment. Data from the reassessment will guide ongoing treatment. The elements of ongoing reassessment involve continuous application of the initial, primary, and secondary assessments to determine the effectiveness of interventions.

Diagnostic Assessments

Diagnostic assessments help detect and identify the presence and severity of respiratory and circulatory problems. You may use some of these assessments (such as rapid bedside glucose or point-of-care laboratory testing) early in your evaluation. The clinical situation dictates the timing of the following diagnostic assessments:

- Arterial blood gas (ABG)
- Venous blood gas (VBG)
- Capillary blood gas
- Hemoglobin concentration
- Central venous O₂ saturation
- Arterial lactate
- Central venous pressure monitoring
- Invasive arterial pressure monitoring
- Chest x-ray
- ECG
- Echocardiogram
- Point-of-care ultrasound sonography

- Peak expiratory flow rate (PEFR)

Arterial Blood Gas

An ABG analysis measures the partial pressure of arterial O_2 (P_{aO_2}) and CO_2 (P_{aCO_2}) dissolved in the blood plasma (ie, the liquid component of blood). An additional tool to assess the adequacy of arterial oxygenation is the pulse oximeter, a device that estimates hemoglobin saturation with O_2 . It may also be calculated from the P_{aO_2} and pH (by using an oxyhemoglobin dissociation curve) or directly measured by using a co-oximeter. Obtain co-oximeter measurement if there is uncertainty about the calculated O_2 saturation and to rule out the presence of carbon monoxide intoxication or methemoglobinemia.



Critical Concepts: Normal P_{aO_2} Does Not Confirm Adequate Blood O_2 Content

P_{aO_2} reflects only the O_2 dissolved in the arterial blood plasma. If the child's hemoglobin is only 3 g/dL, the P_{aO_2} may be normal or high, but O_2 delivery to the tissues will be inadequate. Because most O_2 is carried by hemoglobin in the red blood cells rather than the plasma, a hemoglobin of 3 g/dL is inadequate to carry sufficient O_2 . In this case, the pulse oximeter may reflect 100% saturation despite inadequate O_2 content and delivery.

Identify respiratory failure on the basis of inadequate oxygenation (hypoxemia) or inadequate ventilation (hypercarbia). Use an ABG analysis to confirm your clinical impression and evaluate the child's response to therapy. An ABG analysis, however, is not required to initiate therapy or to make the diagnosis of respiratory failure. The following list will help you interpret an ABG:

- Hypoxemia: low P_{aO_2}
- Hypercarbia: High P_{aCO_2}
- Acidosis: pH less than 7.35
- Alkalosis: pH greater than 7.45



Critical Concepts: ABG and Treatment Decisions in Seriously Ill or Injured Children

Don't wait for an ABG so you can start treatment or determine therapy. Limitations of ABG analysis in pediatric critical care include the following:

- An ABG analysis may not be available (eg, during transport); do not delay therapy initiation.
- A single ABG analysis provides only information at the time the sample was obtained. It does not provide information about trends in the child's condition. Monitoring clinical response to therapy by using serial ABG analyses is often more valuable than any single ABG analysis.

To interpret ABG results, consider the child's clinical appearance and condition. For example, an infant with bronchopulmonary dysplasia (a form of chronic lung disease) is likely to have chronic hypoxemia and hypercarbia. Diagnosing acute respiratory failure in this infant requires clinical examination and evaluating the arterial pH. The infant will compensate for chronic hypercarbia by renal retention of bicarbonate, and, as a result, the child's arterial pH is likely to be normal or nearly normal at baseline. Deterioration will be apparent if the child's respiratory status (ie, hypercarbia) is significantly worse than the baseline status and acidosis develops.

Hyperoxia is an increased arterial oxygen saturation detected by directly measuring oxygen saturation in an ABG sample. This has been associated with worse outcomes, such as after ROSC, in the newly born and in patients with some forms of cyanotic heart disease. Because of the uncertainty of measuring hyperoxia without ABGs, administering oxygen to achieve a displayed pulse oximetry of 100% is not recommended in these conditions.

The arterial pH and bicarbonate (HCO_3) concentrations obtained with ABG analysis may be useful in diagnosing acid-base imbalances. Note that ABG values do not reliably reflect O_2 , CO_2 , or acid-base status in the tissues. However, it is useful to monitor these values over time as an index of improving or worsening tissue oxygenation, as reflected by an increasing base deficit (accumulation of acid in the blood).

Venous Blood Gas

A venous blood pH, measured by VBG analysis, typically correlates well with the pH on ABG analysis. A VBG analysis is not as useful for monitoring ABG status (P_{aO_2} and P_{aCO_2}) in acutely ill children. If the child is well

perfused, the venous PCO_2 is usually within 4 to 6 mm Hg of the arterial PCO_2 . If the child is poorly perfused, however, the gradient between arterial and venous PCO_2 increases. In general, venous PO_2 is not useful in assessing arterial oxygenation.

When interpreting a VBG, also consider the source of the venous specimen. A peripheral specimen that is free flowing from a well-perfused extremity may give results similar to the ABG, but if a tourniquet is used and the specimen is from a poorly perfused extremity, it often shows a much higher PCO_2 and lower pH than an arterial specimen. For this reason, a central venous specimen, if available, is preferred to a peripheral venous specimen. A VBG may be used if an arterial sample is unavailable. There is generally adequate correlation with ABG samples to make venous pH useful in detecting acid-base imbalance.

Capillary Blood Gas

If arterial collection is not practical, CBG analysis can be used. Arterialization of the capillary bed yields pH and PaCO_2 comparable to arterial blood. A CBG analysis is not as useful for estimating arterial oxygenation (PaO_2).

Hemoglobin Concentration

Hemoglobin concentration determines the O_2 -carrying capacity of the blood. O_2 content is the total amount of O_2 bound to hemoglobin plus the unbound (dissolved) O_2 in arterial blood and is determined chiefly by the hemoglobin concentration (in grams per deciliter) and its saturation with O_2 (SaO_2). The amount of O_2 dissolved in the blood is determined by the partial pressure of arterial O_2 (PaO_2), also referred to as the *arterial oxygen tension*, and represents a very small part of total O_2 content at normal hemoglobin concentrations. For more information on the determining the arterial oxygen content, refer to [Impaired Oxygenation and Ventilation in Respiratory Problems](#) in [Part 7](#).

Central Venous Oxygen Saturation

VBGs may indicate changes in the balance between O_2 delivery to the tissues and tissue O_2 consumption. Trends in venous O_2 saturation (SvO_2) may be used as a surrogate for monitoring trends in O_2 delivery (ie, the product of cardiac output and arterial O_2 content). Such trending assumes that O_2 consumption remains stable (an assumption that is not always correct, so remain diligent).

Normal SvO_2 is about 70% to 75%, assuming arterial O_2 saturation is 100%. If the arterial O_2 saturation is not normal, the SvO_2 should be about 25% to 30% below the arterial O_2 saturation. For example, if the child has cyanotic heart disease and the arterial O_2 saturation is 80%, the SvO_2 should be about 55%.

When O_2 delivery to the tissues is low, the tissues consume proportionately more O_2 , so the difference between arterial O_2 saturation and SvO_2 is more substantial when shock is present. For more information about SvO_2 , refer to [Part 10: Managing Shock](#).

Arterial Lactate

The arterial concentration of lactate reflects the balance between lactate production and metabolism or breakdown. In a seriously ill or injured child, the arterial lactate can rise as a result of increased production of lactate (metabolic acidosis) associated with tissue hypoxia and resultant anaerobic metabolism. Arterial lactate is easy to measure, and it's a good prognostic indicator that you can follow sequentially to assess the child's response to therapy. However, arterial lactate can be inaccurate if not drawn with a free-flowing sample or if testing is delayed.

An elevated lactate concentration does not always represent tissue ischemia, especially without accompanying metabolic acidosis. For example, lactate concentration also can be elevated in conditions associated with increased glucose production, such as stress hyperglycemia. In general, it is more helpful to monitor trends in lactate concentration over time than any single measurement. If treatment of shock is effective, the lactate concentration should decrease, but the trend in concentration over time is more predictive than the initial concentration. Lack of response to therapy (ie, the lactate concentration does not decrease) is more predictive of poor outcome than the initial elevated lactate concentration. Monitor central venous lactate concentration if arterial blood samples are not readily available.

Invasive Arterial Pressure Monitoring

Invasive arterial pressure monitoring enables you to continually evaluate and view the systolic and diastolic blood pressures. The arterial waveform pattern may provide information about systemic vascular resistance and compromised cardiac output (eg, *pulsus paradoxus*, an exaggerated decrease in the systolic blood pressure during inspiration). This type of monitoring requires an arterial catheter, monitoring (noncompliant) tubing, a transducer, and a monitoring system. Accurate measurements require you to appropriately zero, level, and calibrate the transducer.

Near-Infrared Spectroscopy

Near-infrared spectroscopy is a noninvasive optical technique to monitor tissue oxygenation in the brain and other tissues. The monitors measure the concentrations of oxyhemoglobin and desaturated hemoglobin to determine cerebral oxygen saturation, which you can follow to assess central venous oxygenation trends. With the cerebral oximeter electrode on the forehead below the hairline, its 2 light-emitting diodes and receivers detect light from shallow and deep tissue, and a computer analyzes the data to provide continuous measurements. Near-infrared spectroscopy monitoring has wide interpatient variability but is used to trend cerebral and other tissue oxygenation in the critical care settings.

Chest X-Ray

A chest x-ray is useful in respiratory illness to aid in diagnosing the following conditions:

- Airway obstruction (upper airway or lower airway)
- Lung tissue disease
- Barotrauma (pneumothorax, pneumomediastinum)
- Pleural disease (pleural effusion/pneumothorax)

A chest x-ray will show the depth of ET tube placement, but it will *not* help determine tracheal vs esophageal placement.

Use the chest x-ray and clinical assessment to evaluate circulatory abnormalities. The chest x-ray may be helpful to assess heart size and presence or absence of congestive heart failure (pulmonary edema).

- A small heart is often present with reduced cardiac preload or severe lung hyperinflation.
- A large heart may be associated with normal or increased cardiac preload, pericardial effusion, congestive heart failure, or when the patient is unable to take a deep breath (eg, with severe abdominal distension).
- An x-ray taken from the front (anteroposterior view) of the heart will appear larger as compared with the heart size when taken from a posterior-anterior view.

Electrocardiogram

Obtain a 12-lead ECG to assess for cardiac arrhythmias. For more information, refer to [Part 12: Managing Arrhythmias](#).

Echocardiogram

Echocardiography is a valuable noninvasive tool for determining

- Cardiac chamber size
- Ventricular wall thickness
- Ventricular wall motion (contractility)
- Valve configuration and motion
- Pericardial space
- Estimated ventricular pressures
- Interventricular septal position
- Congenital anomalies

It can be useful in diagnosing and evaluating cardiac disease. Technical expertise in performing and interpreting the echocardiogram is essential.

Part 5: Recognizing and Managing Cardiac Arrest

This section discusses how to recognize and promptly intervene in life-threatening emergencies and cardiac arrest in infants and children. It covers reversible causes, treatments, and the Pediatric Cardiac Arrest Algorithm.


Learning Objective

After completing this Part, you should be able to recognize cardiopulmonary arrest immediately and begin CPR within 10 seconds.

During the course, you will practice and be tested on CPR skills. Your performance will also be tested in 2 case scenarios.

Rapidly Intervene to Prevent Cardiac Arrest

If not appropriately treated, children with respiratory failure and shock can quickly develop cardiopulmonary failure and even cardiac arrest ([Figure 16](#)). In infants and children, most cardiac arrests result from progressive respiratory failure, shock, or both. Less commonly, pediatric cardiac arrests can occur without warning (ie, with sudden collapse) secondary to an arrhythmia (ventricular fibrillation [VF] or ventricular tachycardia [VT]).



Once cardiac arrest occurs, even with optimal resuscitation efforts, the outcome is generally poor. Only about 8% of children who experience cardiac arrest outside the hospital survive to hospital discharge. The outcome is better for children who experience cardiac arrest in the hospital, although only about 43% of those children survive to hospital discharge. The concepts in this course will help you identify signs of respiratory failure and shock and rapidly intervene to prevent progression to cardiac arrest.

Respiratory conditions may progress to respiratory failure with or without signs of respiratory distress when the child fails to maintain an open airway or adequate respiratory effort, typically because of a decreased level of consciousness. Sudden cardiac arrest in children is less common than in adults and typically results from arrhythmias such as VF or VT. During sports activities, sudden cardiac arrest can occur in children with underlying cardiac conditions that may or may not have been previously recognized.

Life-Threatening Problems

Signs of a life-threatening condition include

- Airway: complete or severe airway obstruction
- Breathing: apnea, significant increased work of breathing, bradypnea
- Circulation: weak or absent pulses, poor perfusion, hypotension, bradycardia

- Disability: unresponsiveness, decreased level of consciousness
- Exposure: significant hypothermia or hyperthermia, significant bleeding, petechiae, or purpura consistent with septic shock or coagulation problem

Interventions

Activate the emergency response system as indicated in your practice setting and begin lifesaving interventions in the following circumstances:

- If the child has a life-threatening condition
- If you are uncertain or “something feels wrong”

If the child does not have a life-threatening condition, begin the secondary assessment and diagnostic assessments.

Cardiac Arrest in Infants and Children

In contrast to adult cardiac arrest, cardiac arrest in infants and children usually results from progressive respiratory failure or shock rather than a primary cardiac cause (ie, sudden cardiac arrest). This *hypoxic/asphyxial arrest* occurs most often in infants and young children, especially those with underlying disease.

It is important to identify and treat respiratory distress, respiratory failure, and shock before progression to cardiopulmonary failure and cardiac arrest. Early identification and treatment are crucial to saving the lives of seriously ill or injured children.

Sudden cardiac arrest from ventricular arrhythmia occurs in about 5% to 15% of all pediatric in-hospital cardiac arrests (IHCA) and out-of-hospital cardiac arrests (OHCA). Although a shockable rhythm (ie, VF or pulseless ventricular tachycardia [pVT]) is the presenting rhythm in only about 14% of pediatric in-hospital arrests, it is present in up to 27% of such arrests at some point during the resuscitation. The incidence of cardiac arrest from VF/pVT increases with age and should be suspected in any patient with a sudden collapse. Increasing evidence suggests that sudden unexpected death in young people is often associated with underlying cardiac conditions.

Despite the improved outcome of in-hospital CPR, most children with IHCA, and an even larger percentage of children with OHCA, do not survive or survive with significant neurologic impairment. Because outcome from cardiac arrest is so poor, focus on preventing cardiac arrest through

- Preventing disease processes and injuries that can lead to cardiac arrest
- Recognizing and managing respiratory distress, respiratory failure, and shock before they cause cardiac arrest

Survival Rates From Pediatric Cardiac Arrest

Information from Get With The Guidelines[®]-Resuscitation shows that survival rates from pediatric cardiac arrest vary according to the location of the arrest and the presenting rhythm. Rate of survival to hospital discharge is higher if the arrest occurs in-hospital (43%) compared with out-of-hospital (8%). Intact neurologic survival is also much higher if the arrest occurs in-hospital. In addition, survival is higher (about 25% to 34%) when the presenting rhythm is shockable (VF or pVT) compared with asystole (about 7% to 24%). Survival for a presenting rhythm of pulseless electrical activity (PEA) is about 38% for in-hospital arrest. Conversely, when VF/pVT develops as a secondary rhythm during a resuscitation attempt (ie, not as the initial arrest rhythm) in hospitalized children, survival is lower than that observed in cardiac arrests with nonshockable rhythms (11% vs 27% survival to discharge). The highest survival rate (64%) occurs when there is bradycardia and poor perfusion, and chest compressions and ventilation are provided before pulseless arrest develops.



For additional information about in-hospital resuscitation outcomes, refer to the Get With The Guidelines-Resuscitation website: www.heart.org/resuscitation.

Definition of Cardiac Arrest

Cardiac arrest occurs when blood circulation ceases because of absent or ineffective cardiac mechanical activity. Clinically, the child is unresponsive and is not breathing or is only gasping, and you cannot detect a pulse. Cerebral hypoxia causes the child to lose consciousness and stop breathing, although you may observe agonal gasps during

the first minutes after sudden arrest. When circulation stops, the resulting organ and tissue ischemia can cause cell, organ, and patient death if not rapidly reversed.

Pathways to Cardiac Arrest

The 2 pathways to cardiac arrest in children are

- Hypoxic/asphyxial
- Sudden cardiac arrest

Hypoxic/Asphyxial Arrest

Hypoxic/asphyxial arrest is the most common cause of cardiac arrest in infants, children, and adolescents and is the end result of progressive tissue hypoxia and acidosis caused by respiratory failure or hypotensive shock. Regardless of the initiating event or disease process, the final common pathway preceding cardiac arrest is cardiopulmonary failure ([Figure 16](#)).



Critical Concepts: Prioritize Airway and Breathing in Children in Cardiac Arrest

In contrast to adults, infants and children usually arrest as the result of progression of respiratory failure or shock. As a result, the child's arterial oxygen content and oxygen delivery to the tissues are often low at the time of cardiac arrest. Although C-A-B sequence for CPR is used for both adults and children, establishing an adequate airway, along with adequate oxygenation and ventilation, should be a particularly high priority in children during CPR.

Sudden Cardiac Arrest

Sudden cardiac arrest, less common in children than it is in adults, is most often caused by the sudden development of VF or pVT. Predisposing conditions or causes for sudden cardiac arrest may include

- Hypertrophic cardiomyopathy
- Anomalous coronary artery
- Long QT syndrome or other channelopathies
- Myocarditis
- Drug intoxication (eg, digoxin, ephedra, cocaine)
- Commotio cordis (ie, sharp blow to the chest)

For primary prevention of pediatric cardiac arrest, providers may use cardiovascular screening (eg, for hypertrophic cardiomyopathy or long QT syndrome) and treat predisposing problems (eg, myocarditis or anomalous coronary artery). Some cases of sudden cardiac arrest in children and young adults are associated with genetic mutations that cause cardiac ion channelopathies. A *channelopathy* is a disorder of the ion channels in myocardial cells that predisposes the heart to arrhythmias. These genetic mutations are known as *familial channelopathies*, so conduct a careful family history to identify syncopal episodes, seizures, and sudden and unexplained death (including sudden infant death syndrome/sudden unexplained infant death, drowning, and even a motor vehicle crash).

For secondary prevention of death from sudden cardiac arrest, providers must perform prompt and effective resuscitation, including timely defibrillation. Most episodes of sudden cardiac arrest in children occur during athletic activity. Coaches, trainers, parents, and the general public must know about sudden cardiac arrest in children so that they can ensure prompt treatment. In addition, bystanders must be prepared and willing to activate the emergency response system, provide high-quality CPR, and use an AED as soon as one is available.

Causes of Cardiac Arrest

Causes of cardiac arrest in children vary based on the child's age and underlying health as well as on event location (in-hospital vs out-of-hospital). Most OHCA in infants and children occurs at or near the home.

Co-bedding is an increasing cause of death in the infant population. Therefore, instruct parents that infants should have their own sleep space. Trauma is the predominant cause of death in children 6 months of age through young adulthood. Causes of traumatic cardiac arrest include airway compromise, tension pneumothorax, hemorrhagic shock, and brain injury.

The most common causes of pediatric cardiac arrest are respiratory failure and hypotensive shock. Arrhythmia is a less common cause of arrest. Cardiac arrest in children may be associated with a reversible condition. If you don't think about reversible causes or complicating factors, you are likely to miss them. Review the following H's and T's to help you identify potentially reversible causes of cardiac arrest or factors that may be complicating resuscitative efforts.

H's

- **Hypovolemia**
- **Hypoxia**
- **Hydrogen ion (acidosis)**
- **Hypoglycemia**
- **Hypo-/hyperkalemia**
- **Hypothermia**

T's

- **Tension pneumothorax**
- **Tamponade (cardiac)**
- **Toxins**
- **Thrombosis, pulmonary**
- **Thrombosis, coronary**

Also consider unrecognized trauma (eg, abdominal injury and hemorrhage) as a cause of cardiac arrest, especially in infants and young children.

Identifying a Child at Risk for Cardiac Arrest



Children with the combination of severe respiratory failure and shock will likely develop cardiac arrest within minutes unless you intervene immediately. Be alert for signs of inadequate oxygenation, ventilation, and tissue perfusion.

Recognizing Cardiac Arrest

Signs of cardiac arrest are

- Unresponsiveness
- No normal breathing or only gasping (agonal gasps aren't effective breathing)
- No pulse felt (assess for no more than 10 seconds)

Arrest rhythm may be noted on the cardiac monitor, but monitoring is not mandatory for recognizing cardiac arrest.

If a child is unresponsive and does not have normal breathing (or only gasping), try to palpate a central pulse (brachial in an infant, carotid or femoral in a child). Because even healthcare providers cannot always detect a pulse, take no more than 10 seconds to try to palpate the pulse. If there is no pulse felt or you are not sure if a pulse is present, start CPR, beginning with chest compressions.

Arrest Rhythms

Cardiac arrest is associated with one of the following rhythms, also known as *arrest rhythms* or *states*:

- Asystole
- PEA
- VF
- pVT, including torsades de pointes

Asystole and PEA are the most common initial rhythms seen in both in-hospital and out-of-hospital pediatric cardiac arrest, especially in children younger than 12 years. Slow wide QRS complex rhythms that immediately precede asystole are often referred to as *agonal rhythms* ([Figure 17](#)). VF and pVT are more likely terminal rhythms in older children with sudden collapse or in children with underlying cardiovascular conditions.




Figure 17. Agonal rhythm (slow ventricular rhythm progressing to asystole).

Asystole

Asystole is cardiac standstill without discernable electrical activity represented by a straight (flat) line on the electrocardiogram (ECG). Do not rely on the ECG to diagnose cardiac arrest; always confirm it clinically because a “flat line” on the ECG also can be caused by a loose ECG lead.

Recall potentially reversible causes of asystole by remembering the H's and T's.

Pulseless Electrical Activity

PEA is not a specific rhythm but a term describing any organized electrical activity (ie, not VF/pVT or asystole) on an ECG or cardiac monitor that is associated with no *palpable* pulses; an arterial waveform or Doppler study may detect pulsations, but these pulses are not palpable. The rate of electrical activity may be slow (most common), normal, or fast. You may refer to very slow PEA as *agonal*.

In PEA, the ECG may display normal or wide QRS complexes or other abnormalities, including

- Low- or high-amplitude T waves
- Prolonged PR and QT intervals
- Atrioventricular dissociation, complete heart block, or ventricular complexes without P waves

Assess the monitored rhythm and note the rate and width of the QRS complexes.

Unless you can quickly identify and treat the cause of PEA (recall the H's and T's), the rhythm will likely deteriorate to asystole.

Ventricular Fibrillation

When VF is present, the heart has no organized rhythm and no coordinated contractions ([Figure 18](#)). Electrical activity is chaotic, and the heart quivers and does not pump blood. Therefore, pulses are not palpable. VF may be preceded by a brief period of VT with or without pulses.




Figure 18A. Ventricular fibrillation. A, Coarse VF. High-amplitude electrical activity varies in size and shape, representing chaotic ventricular electrical activity with no identifiable P, QRS, or T waves.




Figure 18B. Fine VF. Electrical activity is reduced

as compared with previous
(A) rhythm strip.

Primary VF is uncommon in children. In studies of pediatric cardiac arrest, VF was the initial rhythm in 5% to 15% of both OHCA and IHCA, but overall prevalence may be higher because VF may occur early during an arrest and quickly deteriorate to asystole. It has been reported in up to 27% of pediatric in-hospital arrests at some point during the resuscitation.

VF without a previously known underlying cause may occasionally occur in otherwise healthy teens during sports activities because of an undiagnosed cardiac abnormality or channelopathy, such as long QT syndrome. In addition, sudden impact to the chest from a collision or moving object may result in commotio cordis that leads to VF. Consider the H's and T's for other potential reversible causes.

Survival and outcome of patients with VF or pVT as the *initial* arrest rhythm are generally better than those of patients presenting with asystole or PEA. Outcome may be improved by promptly recognizing and providing CPR and defibrillation.

Pulseless Ventricular Tachycardia

VT may produce pulses or may be a form of pulseless arrest of ventricular origin. Because the treatment of pVT differs from the treatment of VT with a pulse, you must assess the pulse to determine appropriate treatment. Almost any cause of VT can present without detectable pulses. Unlike VF, pVT is characterized by organized, wide QRS complexes. As shown in [Figure 19A](#), the ventricular rhythm is rapid and regular at a rate of 158/min (greater than the minimum 120/min characteristic of VT). The QRS is wide (greater than 0.09 second), and there is no evidence of atrial depolarization. The complexes are uniform in appearance, so the VT is monomorphic. This form of pulseless arrest is usually brief before it deteriorates into VF. Refer to [Part 11: Recognizing Arrhythmias](#) for more information.



Figure 19A. Ventricular tachycardia (VT). A, VT in a child with muscular dystrophy and known cardiomyopathy.

Treat pVT exactly the same as VF. Refer to the [Pediatric Cardiac Arrest Algorithm](#) in the [Managing Cardiac Arrest](#) section ([Figure 21](#)).

Torsades de Pointes (Turning on a Point)

pVT may be monomorphic (ventricular complexes appear uniform) or polymorphic (ventricular complexes do not look alike). Torsades de pointes is a distinctive form of polymorphic VT ([Figure 19B](#)), in which the complexes appear to be “turning on a point.” This arrhythmia is seen in conditions distinguished by a prolonged QT interval, including congenital long QT syndrome, drug toxicity, and electrolyte abnormalities (eg, hypomagnesemia). Refer to [Part 11: Recognizing Arrhythmias](#) for more information.



Figure 19B. Torsades de pointes in a child with hypomagnesemia.

Managing Cardiac Arrest

The success of any resuscitation is built on high-quality CPR, timely shock delivery for any shockable rhythm, and good teamwork.

High-Quality CPR

High-quality CPR is the foundation of basic and advanced life support for managing cardiac arrest. Study and practice the characteristics of high-quality CPR, including performing compressions and breaths, so that you can perform each skill effectively to improve a victim's chances of survival.

Consistency in the order of compressions, airway, and breathing for CPR in victims of all ages may be easiest for rescuers who treat people of all ages to remember and perform. Maintaining the same sequence for adults and children offers consistency in teaching.



Critical Concepts: CPR in Hypoxic/Asphyxial Arrest

Conventional CPR (chest compressions and rescue breaths) should be provided for pediatric cardiac arrests by all healthcare providers and trained lay rescuers. The asphyxial nature of the majority of pediatric cardiac arrests, as well as with drowning and drug overdose, requires ventilation as part of effective CPR. However, because Hands-Only (compression-only) CPR is effective in patients with sudden, witnessed cardiac arrest, if rescuers are unwilling or unable to deliver breaths, we recommend rescuers perform compression-only CPR.

Review the fundamentals of BLS in [Table 22](#). These recommendations are based on the *2020 AHA Guidelines for CPR and ECC*.

Table 22. Summary of High-Quality CPR Components for BLS Providers

Component	Adults and adolescents	Children (age 1 year to puberty)	Infants (age less than 1 year, excluding newborns)
Verifying scene safety	Make sure the environment is safe for rescuers and victim		
Recognizing cardiac arrest	Check for responsiveness No breathing or only gasping (ie, no normal breathing) No definite pulse felt within 10 seconds (Breathing and pulse check can be performed simultaneously in less than 10 seconds)		
Activating emergency response system	If a mobile device is available, phone emergency services (9-1-1)		
	If you are alone with no mobile phone, leave the victim to activate the emergency response system and get the AED before beginning CPR Otherwise, send someone and begin CPR immediately; use the AED as soon as it is available	Witnessed collapse Follow steps for adults and adolescents on the left Unwitnessed collapse Give 2 minutes of CPR Leave the victim to activate the emergency response system and get the AED Return to the child or infant and resume CPR; use the AED as soon as it is available	
Compression-to-ventilation ratio without advanced airway	1 or 2 rescuers 30:2	1 rescuer 30:2 2 or more rescuers 15:2	
Compression-to-ventilation ratio with advanced airway	Continuous compressions at a rate of 100-120/min Give 1 breath every 6 seconds (10 breaths/min)	Continuous compressions at a rate of 100-120/min Give 1 breath every 2-3 seconds (20-30 breaths/min)	
Compression rate	100-120/min		
Compression depth	At least 2 inches (5 cm)*	At least one third AP diameter of chest Approximately 2 inches (5 cm)	At least one third AP diameter of chest Approximately 1½ inches (4 cm)
Hand placement	2 hands on the lower half of the breastbone (sternum)	2 hands or 1 hand (optional for very small child) on the lower half of the breastbone (sternum)	1 rescuer 2 fingers or 2 thumbs in the center of the chest, just below

			<p>the nipple line</p> <p>2 or more rescuers</p> <p>2 thumb–encircling hands in the center of the chest, just below the nipple line</p> <p>If the rescuer is unable to achieve the recommended depth, it may be reasonable to use the heel of one hand</p>
Chest recoil	Allow complete recoil of chest after each compression; do not lean on the chest after each compression		
Minimizing interruptions	Limit interruptions in chest compressions to less than 10 seconds		

*Compression depth should be no more than 2.4 inches (6 cm).


Abbreviations: AED, automated external defibrillator; AP, anteroposterior; CPR, cardiopulmonary resuscitation.

Monitoring for CPR Quality

During the resuscitation attempt, the Team Leader as well as team members should monitor CPR quality. Use good team communication to ensure that chest compressions are the appropriate depth and rate, that the chest is allowed to fully recoil after each compression, and that ventilation is not excessive. Elements of high-quality CPR include the following:

- **Push fast:** Push at a rate of 100 to 120 compressions per minute for infants, children, and adults
- **Push hard:** Push with enough force to depress the chest at least one third the anteroposterior diameter of the chest in pediatric patients (infants [younger than 1 year] to children up to the onset of puberty). This equates to approximately 1½ inches (4 cm) in infants and 2 inches (5 cm) in children. Once children have reached puberty (ie, adolescents), use the recommended adult compression depth of at least 2 inches (5 cm), but no greater than 2.4 inches (6 cm).
- **Allow complete chest recoil:** Allow complete chest recoil after each compression; this allows the heart to refill with blood.
- **Minimize interruptions:** Try to limit interruptions in chest compressions to 10 seconds or less or as needed for interventions (eg, defibrillation). Ideally, compressions are interrupted only for ventilation (until an advanced airway is placed), rhythm check, and actual shock delivery. Once an advanced airway is in place, provide continuous chest compressions and asynchronous ventilation (ie, without pausing compressions for ventilation).
- **Avoid excessive ventilation:** Each rescue breath should be given over about 1 second and should result in visible chest rise. After an advanced airway is in place, deliver 1 breath every 2 to 3 seconds or about 20 to 30 breaths per minute, being careful to avoid excessive ventilation.

Many in-hospital patients, especially those in an intensive care unit, have advanced monitoring in place; some have advanced airways and receive mechanical ventilation. Continuously monitoring the child's end-tidal CO₂ (PETCO₂) can provide indirect evidence of chest compression quality (Figure 20). If PETCO₂ is less than 10 to 15 mm Hg, cardiac output during CPR is probably low, meaning not much blood is delivered to the lungs. Verify that cardiac compressions are effective, with a goal of increasing the PETCO₂ to greater than 10 to 15 mm Hg. If the child has an indwelling arterial catheter, use the waveform as feedback to evaluate hand position and chest compression depth. Minorly adjusting hand position or compression depth can significantly improve the amplitude of the arterial waveform, reflecting better chest compression–induced stroke volume and cardiac output. Verify that ventilation is not excessive. Both the PETCO₂ and arterial waveform may be useful in identifying ROSC.



This capnography tracing displays the PETCO₂ in mm Hg on the vertical axis over time (depicted horizontally). This patient is intubated and receiving CPR. Note that the ventilation rate is approximately 8 to 10/min. Chest compressions in this patient were given continuously at a rate slightly faster than 100/min but are not visible with this tracing. The initial PETCO₂ is less than 12.5 mm Hg during the first minute, indicating very low blood flow. In this example, the PETCO₂ increases to between 12.5 and 25 mm Hg during the second and third minutes, consistent with the increase in blood flow with ongoing resuscitation. ROSC occurs during the fourth minute and is identifiable by the

abrupt increase in the PETCO₂ (visible just after the fourth vertical line) to over 40 mm Hg, which is consistent with a substantial improvement in blood flow with ROSC.

PALS in Cardiac Arrest

The immediate goal of therapeutic interventions for cardiac arrest is ROSC, which occurs when an organized cardiac electrical rhythm on the monitor and palpable central pulses resume. Corresponding clinical evidence of perfusion will also be apparent (eg, sudden increase in PETCO₂, measurable blood pressure, improved color).

PALS treatment of cardiac arrest may include the following:

- Rhythm analysis (shockable vs nonshockable)
- Establishment of vascular access
- Defibrillation
- Medication therapy
- Advanced airway management

Arrest Rhythm Assessment

The Pediatric Cardiac Arrest Algorithm outlines the recommended sequence of CPR, shocks, and medication administration for both shockable and nonshockable pediatric cardiac arrest rhythms, so identifying the arrest rhythm as shockable or nonshockable determines which pathway to follow. Although the algorithm depicts actions sequentially, many actions (eg, compressions and medication administration) are typically performed simultaneously when multiple rescuers are present.

Vascular Access

Priorities for drug delivery routes during PALS are (in order of preference)

1. Intravenous
2. Intraosseous
3. Endotracheal

When the critically ill child develops cardiac arrest, vascular access may already be established. If vascular access is not present, establish it immediately. Peripheral IV access is the first choice during resuscitation if you can place it rapidly, but placement may be difficult in critically ill or injured children. Limit the time you spend trying to obtain IV access in a seriously ill or injured child. If IV access is not already present and you cannot achieve it immediately, establish IO access. *IO access is useful as the initial vascular access in cases of cardiac arrest.* If neither IV nor IO access is available for medication delivery, the endotracheal (ET) route is a third option.

Intravenous Route

Although a central venous catheter provides a more secure route of vascular access than a peripheral catheter, central venous access is not needed during most resuscitation attempts, and its placement requires interruptions in chest compressions. This is because trying to place the central catheter during chest compressions may cause vascular lacerations, hematomas, pneumothorax, and bleeding. If a central venous catheter is already in place, it is the preferred route for drug and fluid administration. Central venous administration of medications provides more rapid onset of action and higher peak concentration than peripheral venous delivery.

Establishing peripheral venous access does not require interrupting CPR, but it may delay drug delivery to the central circulation. To improve drug delivery, do the following when administering drugs into a peripheral IV catheter infusion system:

- Give the drug by bolus injection
- Give the drug while chest compressions are being performed
- Follow with a 5-mL flush of normal saline to move the drug from the peripheral to the central circulation

Intraosseous Route

If IV access is not available, drugs and fluids can be delivered safely and effectively via the IO route. In fact, the IO route is useful as the initial route of vascular access in cases of cardiac arrest. Important points about IO access are

- IO access can be established in all age groups
- IO access can often be achieved in 30 to 60 seconds
- The IO route is preferred over the ET route

- Any drug or fluid that can be administered IV can be administered IO

IO cannulation provides access to a noncollapsible marrow venous plexus, which serves as a rapid, safe, and reliable route for administering resuscitation drugs and fluids. The technique uses a rigid needle, preferably a specially designed IO or bone marrow needle. Although an IO needle with a stylet is preferred to prevent obstruction of the needle with cortical bone during insertion, butterfly needles, standard hypodermic needles, and spinal needles can be inserted and used effectively. Powered IO insertion devices are commercially available and are widely used in the United States. Refer to the [Resources for Managing Circulatory Emergencies](#) in [Part 10](#) of this manual for more information on IO access.

Endotracheal Route

The IV and IO routes are preferable to the ET route for administering drugs, but lipid-soluble drugs can be given by the ET route. These include lidocaine, epinephrine, atropine, and naloxone (LEAN) and vasopressin. However, there are limited human studies about ET vasopressin administration and limited studies to provide dosing guidelines for most drugs given by the ET route.

When considering drug administration via the ET route during CPR, keep these concepts in mind:

- Drug absorption from the tracheobronchial tree is unpredictable, so drug concentrations and drug effects will be unpredictable.
- The optimal drug dose given by the ET route is unknown.
 - Drug administration into the trachea results in lower blood concentrations than the same dose given via IV or IO routes.
 - Animal data suggest that the lower epinephrine concentrations achieved when the drug is delivered by the ET route may produce transient but detrimental β -adrenergic-mediated vasodilation.
- Recommended drug doses administered by the ET route are higher than for the IV/IO route.
 - The recommended ET dose of epinephrine is 10 times the IV/IO dose.
 - The typical ET dose of other drugs is 2 to 3 times the IV/IO dose.

To give a drug by the ET route, administer it as follows:

- Instill the drug into the ET tube (briefly pause compressions during instillation).
- Follow with a minimum of 5 mL normal saline flush; a smaller volume may be used in neonates.
- Provide 5 rapid positive-pressure breaths after the drug is instilled.

Defibrillation

A defibrillation shock “stuns” the heart by depolarizing a critical mass of the myocardium. If a shock is successful, it terminates VF, which allows the heart’s natural pacemaker cells to resume an organized rhythm. The return of an organized rhythm alone, however, does not ensure survival. The organized rhythm must ultimately produce effective cardiac mechanical activity that results in ROSC, defined by the presence of palpable central pulses. If the child’s PETCO₂ or intra-arterial pressure is being monitored, it also can indicate ROSC ([Figure 20](#)).

When attempting defibrillation, provide compressions until the defibrillator is charged, deliver 1 shock, and immediately resume CPR, starting with chest compressions. Chest compressions are needed to maintain blood flow to the heart (the coronary circulation) and brain until effective cardiac contractility resumes. No evidence indicates that performing chest compressions in a child with spontaneous cardiac activity is harmful. If the shock does not eliminate VF, the heart is probably ischemic. Resuming chest compressions is likely to be of greater value to the child than immediate delivery of a second shock.

In an out-of-hospital or unmonitored setting, do not waste time looking for a shockable rhythm or palpating a pulse immediately after shock delivery; neither is likely to be present. Resume high-quality CPR, beginning with chest compressions. Providers may modify this sequence in hospital units with invasive arterial monitoring. In in-hospital settings with invasive monitoring in place, return of an arterial waveform or a sudden increase in PETCO₂ suggests ROSC. When monitored parameters indicate ROSC, confirm by palpating a central pulse.

For more information about the manual defibrillation procedure, refer to the [Critical Concepts box Manual Defibrillation \(for VF or pVT\)](#), later in this Part.



If a shock eliminates VF, continue CPR because most victims have asystole or PEA immediately after shock delivery.

Medication Therapy

The objectives for medication administration during cardiac arrest are to

- Increase coronary and cerebral perfusion pressures and blood flow
- Stimulate spontaneous or more forceful myocardial contractility
- Accelerate heart rate
- Correct and treat the possible cause of cardiac arrest
- Suppress or treat arrhythmias

Medications that may be used while treating pediatric cardiac arrest are listed in [Table 23](#).

Table 23. Pediatric Cardiac Arrest Medications

Agent	Indication	Mechanism(s) of action	Clinical data
Vasopressors			
Epinephrine	<ul style="list-style-type: none"> • Used for cardiac arrest associated with VF/pVT as well as asystole/PEA • High doses may be considered for special resuscitation circumstances, such as β-blocker overdose 	<ul style="list-style-type: none"> • Causes α-adrenergic–mediated vasoconstriction that increases aortic diastolic pressure and coronary perfusion pressure, a critical determinant of successful resuscitation 	<ul style="list-style-type: none"> • Both beneficial and toxic physiologic effects during CPR have been demonstrated in animal and human studies • No pediatric studies have demonstrated improved survival with the use of epinephrine • High doses may be harmful, particularly in hypoxic/asphyxial arrest • High-dose IV/IO epinephrine is not recommended because it offers no survival advantage
Antiarrhythmics			
Amiodarone	<ul style="list-style-type: none"> • May be used for shock-refractory VF or pVT 	<ul style="list-style-type: none"> • α-Adrenergic and β-adrenergic blocking activity • Affects sodium, potassium, and calcium channels • Slows atrioventricular conduction • Prolongs the atrioventricular refractory period and QT interval • Slows ventricular conduction (widens the QRS) 	<ul style="list-style-type: none"> • Increased survival to hospital admission but not to hospital discharge compared with placebo or lidocaine for shock-resistant VF in adults • No association between amiodarone use and ROSC, 24-hour survival, or survival to hospital discharge in pediatric observational study
Lidocaine	<ul style="list-style-type: none"> • May be used to treat shock-refractory VF or pVT in children 	<ul style="list-style-type: none"> • Decreases automaticity and suppresses ventricular arrhythmias 	<ul style="list-style-type: none"> • Pediatric observational data showed lidocaine improved ROSC as compared with no lidocaine • No association between lidocaine use and survival to hospital discharge
Magnesium sulfate	<ul style="list-style-type: none"> • Used to treat torsades de pointes • Used for hypomagnesemia 	<ul style="list-style-type: none"> • Used to treat arrhythmias associated with hypomagnesemia 	<ul style="list-style-type: none"> • Insufficient evidence to recommend for or against routine use in pediatric cardiac arrest not associated with torsades de pointes or hypomagnesemia
Other Agents			
Atropine	<ul style="list-style-type: none"> • Indicated to treat bradycardia, especially if it results from excessive vagal tone, cholinergic drug toxicity (eg, 	<ul style="list-style-type: none"> • Increases heart rate 	<ul style="list-style-type: none"> • No published studies suggesting efficacy for treating cardiac arrest in pediatric patients • Refer to complete discussion in Part 12: Managing Arrhythmias

	organophosphates), or complete atrioventricular block		
Calcium	<ul style="list-style-type: none"> • Routine use in cardiac arrest is not recommended • Indicated for documented ionized hypocalcemia (relatively common in critically ill children, particularly during sepsis or after cardiopulmonary bypass) and hyperkalemia, particularly in children with hemodynamic compromise • May also be considered for hypermagnesemia or calcium channel blocker overdose 	<ul style="list-style-type: none"> • Restores calcium • Helps maintain the cell membrane action potential threshold • Helps maintain the gradient between intracellular potassium and extracellular sodium 	<ul style="list-style-type: none"> • Does not improve survival in cardiac arrest and may be harmful
Sodium bicarbonate	<ul style="list-style-type: none"> • Routine administration in cardiac arrest is not recommended • Recommended for symptomatic patients with hyperkalemia, tricyclic antidepressant overdose, or an overdose of other sodium channel blocking agents 	<ul style="list-style-type: none"> • Helpful in treating arrhythmias in tricyclic overdose • Rapidly reduces potassium concentrations in hyperkalemia 	<ul style="list-style-type: none"> • Does not improve survival in cardiac arrest

Advanced Airway Management

When managing the airway and ventilation in pediatric victims of cardiac arrest, consider the following:

- Avoid excessive ventilation during resuscitation.
 - Excessive ventilation can be harmful because it impedes venous return and decreases cardiac output.
 - Increased intrathoracic pressure from positive-pressure ventilation also elevates right atrial pressure and thus reduces coronary perfusion pressure.
 - When providing ventilation with a bag and mask (in cycles of 15 compressions and 2 breaths for 2 rescuers), give each breath over 1 second and provide just enough volume to make the chest rise.
 - Excessive tidal volume or pressure during bag-mask ventilation may distend the stomach; gastric distention impedes ventilation and increases the risk of regurgitation and aspiration.
- Avoid routine use of cricoid pressure if it interferes with intubation or ventilation.
- Use waveform capnography or capnometry with clinical examination to confirm and monitor ET tube placement.
- Colorimetric exhaled CO₂ devices may fail to detect the presence of exhaled CO₂ (ie, lack of a color change indicates no CO₂ detected) during cardiac arrest despite correctly placing the ET tube. Use direct laryngoscopy to confirm tube placement if exhaled CO₂ is not detected and there is evidence that the tube is in the trachea (eg, chest rise and bilateral breath sounds).
- When providing ventilation via an advanced airway during CPR, provide 1 breath every 2 to 3 seconds (20-30 breaths/min) without pausing chest compressions. Chest compressions are delivered without interruption at a rate of 100 to 120/min. For more details, refer to [Inserting an Advanced Airway During CPR](#) section later in this Part.

An advanced airway (eg, ET tube) can be placed during CPR. However, in a study of OHCA, ET intubation demonstrated no survival advantage over effective bag-mask ventilation when emergency medical services transport time was short, and providers had limited ongoing experience in pediatric intubation. This study does not address ET intubation in the in-hospital setting but suggests that immediate intubation may not be necessary.

Pediatric Cardiac Arrest Algorithm

The **Pediatric Cardiac Arrest Algorithm** (Figure 21) outlines steps for assessing and managing an infant or child in cardiac arrest who does not respond to BLS interventions. The expert consensus–based Pediatric Cardiac Arrest Algorithm maximizes uninterrupted periods of CPR while enabling efficient delivery of electrical therapy and medications as appropriate. Although the actions appear sequentially, when several rescuers are involved, some actions will occur simultaneously.




Figure 21. Pediatric Cardiac Arrest Algorithm.



Critical Concepts: Coordinating Team Members During Resuscitation

Using the Pediatric Cardiac Arrest Algorithm, providers should structure assessments and interventions around 2-minute periods of uninterrupted high-quality CPR, rotating chest compressors after every 2-minute period. This requires organization so that all team members know their responsibilities. When all team members are familiar with the algorithm, they can anticipate next steps and prepare the equipment and proper doses of medications.

The algorithm consists of 2 pathways, depending on the cardiac rhythm as identified on a monitor or interpreted by an AED. A shockable rhythm (VF/pVT) pathway is displayed on one side of the algorithm while a nonshockable rhythm (asystole/PEA) pathway is displayed on the other side of the algorithm.

Step numbers in the following sections refer to the corresponding steps in the algorithm.

Start CPR (Step 1)

- **Start CPR**
 - As soon as the child is found to be unresponsive with no breathing (or only gasping), shout for nearby help and activate the emergency response system, send for a defibrillator (manual or AED), check for a pulse, and start CPR, beginning with chest compressions. Attach the ECG monitor or AED pads as soon as they are available. Throughout the resuscitation attempt, provide high-quality CPR (give chest compressions of adequate rate and depth, allow complete chest recoil after each compression, minimize interruptions in compressions, and avoid excessive ventilation). Use a compression-to-ventilation ratio of 30:2 for 1 rescuer and 15:2 for 2 rescuers.
- **Begin bag-mask ventilation and give oxygen.**
- **Attach monitor/defibrillator.** Once the monitor/defibrillator is attached, check the rhythm. Determine whether the rhythm is shockable (VF/pVT) or nonshockable (asystole/PEA). If the rhythm is shockable, follow the VF/pVT pathway of the algorithm.

Shockable rhythm: VF/pVT (Step 2)

If the rhythm is shockable, deliver 1 unsynchronized shock (Step 3). Perform CPR while the defibrillator is charging, if possible, because the shorter the interval is between the last compression and shock delivery, the higher the potential shock success (elimination of VF). Therefore, try to keep that interval as short as possible, ideally less than 10 seconds. Immediately after shock delivery, resume high-quality CPR, beginning with chest compressions. Providers may use their discretion to modify this approach in a monitored setting.

If the resuscitation attempt takes place in a critical care setting and the child has continuous intra-arterial monitoring in place, the diastolic blood pressure can be used to guide CPR quality. Invasive arterial blood pressure monitoring during CPR provides insight to blood pressures generated with compressions and medications. End-tidal CO₂ reflects

both the cardiac output produced and ventilation efficacy and may provide feedback on the quality of CPR. A sudden rise in the End-tidal CO₂ may be an early sign of ROSC. In other settings, you can identify ROSC after about 2 minutes of CPR during the next rhythm check.

Defibrillation devices for children are the

- AED (able to distinguish pediatric shockable from nonshockable rhythms and ideally equipped with a pediatric dose attenuator)
- Manual cardioverter/defibrillator (capable of variable shock doses)

Institutions that care for children at risk for arrhythmias and cardiac arrest (eg, hospitals, emergency departments) ideally should have defibrillators available that can adjust the energy dose for children.

Manual Defibrillator

The optimal electrical energy dose for pediatric defibrillation is unknown, but an initial dose of 2 to 4 J/kg is acceptable. For ease of teaching, consider a 2 J/kg dose (biphasic or monophasic waveform). If VF or pVT persist at the next rhythm check, deliver a 4 J/kg dose for the second shock. If VF persists after the second shock, use at least 4 J/kg or higher, but do not exceed 10 J/kg or the maximum adult dose. Successful resuscitation using shock doses up to 9 J/kg have been reported in children.

Refer to the Critical Concepts box Manual Defibrillation (for VF or pVT), later in this Part for the universal steps for operating a manual defibrillator.

Pads/Paddles

Use either self-adhesive electrode pads or paddles to deliver shocks with a manual defibrillator. Self-adhesive pads are preferred because they are easy to apply and reduce the risk of current arcing, but they also can be used to monitor the heart rhythm. If you use paddles, apply a conducting gel, cream, or paste, or place an electrode pad between the paddle and the child's chest to reduce transthoracic impedance. Do not use saline-soaked gauze pads, sonographic gels, or alcohol pads because alcohol pads may pose a fire hazard and cause chest burns.

Use the largest self-adhering electrode pads that will fit on the chest wall while ensuring that the pads don't touch. Recommended paddle sizes are based on the child's weight/age:

- Pediatric patients heavier than 10 kg (or approximately 1 year or older) should use large "adult" paddles (8-13 cm).
- Pediatric patients lighter than 10 kg (or younger than 1 year) should use small "infant" paddles (4.5 cm).

The selection of pediatric pads vs adult pads is manufacturer specific. Refer to package instructions to determine the appropriate size.

Place the electrode pads/paddles so that the heart is between them. Place 1 electrode pad/paddle on the upper right side of the victim's chest below the right clavicle and the other to the left of the left nipple in the anterior axillary line directly over the heart, making sure they don't touch. Allow at least 3 cm between paddles, and apply firm pressure to create good contact with the skin.

However, some defibrillator manufacturers recommend placing self-adhesive electrode pads in an anteroposterior position, with one over the victim's heart and the other over the back. This placement may be necessary in an infant, particularly if only large electrode pads are available. Place the electrode pads according to the recommendations of the defibrillator manufacturer that are typically illustrated on the pads themselves.

Modifications may be required in special situations (eg, if the child has an implanted defibrillator).

Clearing for Defibrillation

To ensure rescuer safety during defibrillation, visually check the child and the resuscitation team just before you deliver a shock. Make sure that high-flow O₂ is not directed across the child's chest. Warn others that you are about to deliver a shock and that everyone must stand clear. (This entire sequence should take less than 5 seconds.) Refer to the Critical Concepts box Manual Defibrillation (for VF or pVT).



Critical Concepts: Manual Defibrillation (for VF or pVT) (Step 3)

Continue CPR without interruption during all steps until the defibrillator is fully charged. Minimize interval between the last compression and shock delivery (do not deliver breaths between last compression and shock delivery).

1. Turn on the defibrillator.
2. Set lead switch to paddles (or lead I, II, or III if monitor leads are used).
3. Select adhesive pads or paddles; use the largest pads or paddles that can fit on the patient's chest without touching each other.
4. If using paddles, apply conductive gel or paste. Be sure cables are attached to the defibrillator.
5. Position the adhesive pads on the patient's chest: right anterior chest wall and left axillary positions. If using paddles, apply firm pressure. If the patient has an implanted pacemaker, do not place the pads/paddles directly over the device. Be sure that oxygen is not directed over the patient's chest.
6. Select energy dose:
Initial dose: 2 J/kg (acceptable range 2-4 J/kg)
Subsequent doses: 4 J/kg or higher (not to exceed 10 J/kg or standard adult dose)
7. Announce "Charging defibrillator," and press Charge on the defibrillator controls or apex paddle. Continue chest compressions during charging unless charging occurs immediately.
8. When the defibrillator is fully charged, state a firm chant, such as "I am going to shock on 3." Then count. This chant can be shortened to "Clear for shock." (Continue chest compressions until this announcement.)
9. After confirming all personnel are clear of the patient, press the Shock button on the defibrillator or press the 2 paddle discharge buttons simultaneously.
10. Immediately after shock delivery, resume CPR beginning with compressions for 5 cycles (about 2 minutes), and then recheck the rhythm. Minimize interruptions to compressions.

Resume CPR, establish IV/IO access, check rhythm (Step 4)

- Perform CPR for 2 minutes
 - Resume CPR immediately after the shock, beginning with chest compressions. Give about 2 minutes of CPR. (For 2 rescuers, this will be about 10 cycles of 15 compressions followed by 2 breaths.)
- Establish IV/IO access
 - While CPR is being performed, if you do not already have vascular access (IV/IO), another member of the resuscitation team should establish vascular access to get ready for medications.

Before the rhythm check, the Team Leader should ensure that the team is prepared to do the following:

- Rotate compressors
- Calculate appropriate shock dose to administer if VF/pVT persists
- Prepare drugs for administration if indicated

After 2 minutes of CPR, check the rhythm. *Limit interruptions in CPR to less than 10 seconds for a rhythm check.*

This rhythm check may indicate the following outcomes from the previous shock and CPR:

- VF/pVT becomes a "nonshockable" rhythm (asystole, PEA) or an organized rhythm with a pulse
- A shockable rhythm (VF/pVT) persists

Terminating VF/pVT (ie, rhythm is "nonshockable") (Step 12)

If the rhythm is nonshockable, check for an "organized" rhythm (ie, one with regular complexes [Figure 22]). If there are no signs of ROSC, go to Step 10.



Figure 22. VF converted to organized rhythm after defibrillation (successful shock).

If the rhythm is organized, palpate a central pulse. If there is return of spontaneous circulation (ROSC) go to the Post-Cardiac Arrest Care checklist (Part 13). If the rhythm is not organized, do not perform a pulse check; instead, immediately resume CPR (Step 10). Remember, rhythm and pulse checks should be brief (less than 10 seconds). In specialized environments (eg, an intensive care unit) with intra-arterial or other hemodynamic monitoring in place, providers may alter this sequence.

If PEA (an organized rhythm without a palpable pulse) or asystole presents at the rhythm check, resume CPR beginning with compressions and proceed to Step 10.

If you doubt the presence of a pulse, immediately resume CPR (Step 10). Chest compressions are unlikely to harm a child with a spontaneous rhythm and weak pulses. *If you don't detect a pulse or you are not sure, resume CPR.*

Persistent VF/pVT (Step 5)

If the rhythm check reveals a shockable rhythm (ie, persistent VF/pVT), prepare to deliver a second shock with a manual defibrillator (4 J/kg) or AED. Resume chest compressions while the defibrillator is charging. If IV/IO access is established, administer epinephrine while compressions continue. Consider inserting an advanced airway if one is not already in place. Once the defibrillator charges, “clear” the patient and deliver the shock (Step 5).

Resume CPR with chest compressions immediately after shock delivery (Step 6). A different compressor should perform the compressions (ie, compressors should rotate every 2 minutes). Give about 2 minutes of CPR. (For 2 rescuers, this will be about 10 cycles of 15 compressions followed by 2 breaths.)

Give epinephrine (Step 6)

- Perform CPR for 2 minutes
- Administer epinephrine every 3 to 5 minutes
- Consider an advanced airway

If VF/pVT persists, administer epinephrine every 3 to 5 minutes as soon as IV/IO access is available while compressions continue. For the IV/IO route, give epinephrine 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration) dose. For the ET route, give epinephrine 0.1 mg/kg (0.1 mL/kg of the 1 mg/mL concentration) dose. Repeat epinephrine every 3 to 5 minutes of cardiac arrest. This will generally result in epinephrine delivery after every second (ie, every other) rhythm check.

The Pediatric Cardiac Arrest Algorithm does not state a specific time for delivering the first dose of epinephrine. However, the *2020 AHA Guidelines for CPR and ECC* states that the earlier epinephrine is administered after CPR initiation, the more likely the patient is to survive. The 2020 Guidelines also state that for pediatric patients in any setting, it is reasonable to administer the initial dose of epinephrine within 5 minutes after the start of chest compressions.

A provider may choose to administer epinephrine before or after the second shock. When providers identify VF/pVT during the rhythm check, they can administer epinephrine during the CPR that precedes (during charging) or immediately follows the shock delivery. Conversely, if an organized rhythm is present during the rhythm check, it is reasonable to check for a pulse to avoid an unnecessary dose of epinephrine because it can produce adverse effects. For example, if the initial VF/pVT was related to a cardiomyopathy, myocarditis, or drug toxicity, epinephrine administration immediately after eliminating VF/pVT could induce recurrent VF/pVT.

Medication Administration During CPR

Ideally, administer IV/IO medications *during compressions* because blood flow generated by compressions helps to circulate the drugs. The *2020 AHA Guidelines for CPR and ECC* recommends administering medication during compressions, to avoid delaying shock delivery and so that the drugs have time to circulate before the next rhythm check.

Team members responsible for resuscitation drugs should prepare the *next* drug dose that might be needed after the next rhythm check. All team members should be familiar with the Pediatric Cardiac Arrest Algorithm and refer to it during the resuscitation attempt to anticipate the next interventions. Make drug tables, charts, or other references readily available to expedite the calculation of drug doses. Using a color-coded length-based resuscitation tape can help rapidly estimate appropriate drug doses.

Administering resuscitation drugs by the ET route results in lower blood concentrations than the same dose given intravascularly or by IO route. Studies also suggest that the lower epinephrine concentration achieved when the drug is delivered by the ET route may produce transient β -adrenergic effects (rather than the desired α -adrenergic vasoconstrictive effects). The β -adrenergic effects can be detrimental and cause hypotension, lower coronary artery perfusion pressure and flow, and reduce potential for ROSC. Another disadvantage of ET drug delivery is that providers must interrupt chest compressions for drug delivery by this route.

Inserting an Advanced Airway During CPR

The Team Leader will determine the best time for ET intubation or inserting another advanced airway. Because inserting an advanced airway will likely interrupt chest compressions, the Team Leader must weigh the relative benefits of the advanced airway vs benefits of minimizing interruptions in compressions. If you must insert an advanced airway, carefully plan and organize supplies and personnel to minimize the time that compressions are interrupted. Once you insert the advanced airway, confirm that it is in the correct position.

When an advanced airway is in place during CPR, provide continuous chest compressions. Refer to the [Critical Concepts box CPR With an Advanced Airway](#).



Critical Concepts: CPR With an Advanced Airway

Once an advanced airway (eg, ET tube) is in place, change the CPR sequence from “cycles” to continuous chest compressions and a regular ventilation rate. While one team member compresses the chest at a rate of 100 to 120/min, another team member ventilates with 1 breath every 2 to 3 seconds (a rate of about 20-30 breaths per minute).

Compressors should rotate every 2 minutes to reduce rescuer fatigue and provide high-quality chest compressions throughout the resuscitation. Limit interruptions of chest compressions to the minimum required for rhythm checks and shock delivery. However, providers may modify this sequence in special settings (eg, an intensive care unit) with continuous ECG and hemodynamic monitoring in place.

Check rhythm (Step 7)

After 2 minutes of CPR and epinephrine administration, recheck the rhythm. Try to limit interruptions in chest compressions for rhythm checks to less than 10 seconds. If the rhythm check reveals termination of VF/pVT, check for an organized rhythm.

- No organized rhythm (asystole/PEA): Go to Step 12.
- Organized rhythm: Check pulse.
 - If there are signs of ROSC, go to the [Post-Cardiac Arrest Care Checklist](#) (in [Part 13](#)).
 - If no signs of ROSC, go to Step 10.

However, if the rhythm check reveals persistence of VF/pVT, go to Step 8.

For persistent VF/pVT (Step 8)

Deliver shock

If VF/pVT persists, deliver 1 shock by manual defibrillator (4 J/kg or more, up to 10 J/kg or the maximum adult dose) or AED. Perform chest compressions, if possible, while the defibrillator is charging. Once the defibrillator charges, “clear” the victim and deliver the shock.


Immediately after the shock, resume CPR, beginning with chest compressions. Give about 2 minutes of CPR. (For 2 rescuers, this will be about 10 cycles of 15 compressions followed by 2 breaths.)

Antiarrhythmic medications (Step 8)

Immediately after resuming chest compressions, administer amiodarone or lidocaine. Give amiodarone 5 mg/kg IV/IO bolus; you may repeat 5 mg/kg IV/IO bolus up to 3 total doses for refractory VF/pulseless VT, or give lidocaine IV/IO at an initial 1 mg/kg loading dose. If torsades de pointes presents, give magnesium 25 to 50 mg/kg IV/IO bolus, maximum dose 2 g.

Summary of the VF/pVT Sequence

[Figure 23](#) summarizes the recommended sequence of CPR, rhythm checks, shocks, and administration of drugs for VF/pVT based on expert consensus.



sequence.

Think of managing pulseless arrest with VF/pVT as providing nearly continuous CPR. Ideally, interrupt CPR for only brief periods, such as for rhythm checks and shock delivery. Drug preparation and administration do not require interrupting CPR and should not delay shock delivery.

Prepare the next drug before rhythm check. Administer drug during chest compressions, as soon as possible after the rhythm check confirms VF/pVT. Do not delay shock. Continue CPR while the drugs are prepared and administered and the defibrillator is charging. Ideally, interrupt chest compressions only for ventilation (until advanced airway is placed), rhythm check, and actual shock delivery.

Nonshockable rhythm (Asystole/PEA, Step 9)

If the rhythm is nonshockable, asystole or PEA may be present. Managing this rhythm is outlined on the Asystole/PEA pathway of the [Pediatric Cardiac Arrest Algorithm](#) (Figure 21). If VF develops during the resuscitation attempt, return to the VF/pVT side of the Pediatric Cardiac Arrest Algorithm (Step 5 or 7).

Establish vascular access (Step 10)

For treating asystole or PEA, provide high-quality CPR, deliver epinephrine as soon as possible, and try to identify and treat potentially reversible causes of the arrest.

Continue high-quality CPR for about 2 minutes. During this time, establish vascular (IV/IO) access and consider advanced airway placement and capnography. As soon as you establish vascular access, give a bolus of epinephrine during chest compressions. For the IV/IO route, give 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration) dose. For the ET route, give 0.1 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration) dose. Administer epinephrine every 3 to 5 minutes if cardiac arrest persists. This generally results in epinephrine administration after every second (ie, every other) rhythm check.

Check rhythm


After about 2 minutes of CPR, check the rhythm. If the rhythm is shockable, go to Step 7, but if it is nonshockable, go to Step 11.

Nonshockable rhythm (Step 11)

If the rhythm check reveals a nonshockable rhythm, resume CPR immediately, beginning with chest compressions, and treat potentially reversible causes of cardiac arrest (recall the **H's** and **T's** mnemonic). After another 2 minutes of CPR, check the rhythm. If it is organized, check for a pulse. If ROSC, go to the [Post-Cardiac Arrest Care](#) Checklist (see [Part 13](#)). If there are no signs of ROSC, go to Step 10. If a shockable rhythm presents at any time, go to Step 7.

Summary of Asystole/PEA Treatment Sequence

Figure 24 summarizes the recommended CPR sequence, based on expert consensus, of rhythm checks, shocks, and drug delivery for asystole and PEA.



Managing any pulseless arrest includes providing nearly continuous CPR, interrupted by only brief rhythm checks. Do not interrupt CPR for drug preparation and administration. Administer IV/IO drugs during chest compressions. Refer to the [Critical Concepts](#) box [CPR With an Advanced Airway](#) earlier in this Part.

If you detect a shockable rhythm at any rhythm check, refer to [Figure 23](#).

Identifying and Treating Potentially Reversible Causes of Cardiac Arrest

The outcome of hypoxic-ischemic pediatric cardiac arrest is generally poor. Rapid recognition, immediate high-quality CPR, and correcting contributing factors and potentially reversible causes offer the best chance for a successful resuscitation. If you can quickly identify the condition and treat it, your resuscitative efforts may be successful.

When searching for potentially reversible causes or contributing factors, do the following:

- Ensure that high-quality CPR is being provided.
- Check that the advanced airway is open/patent and effective.
- Verify that the bag-mask device is connected to a source of high-flow O₂.
- Ensure that ventilation produces visible chest rise without excessive volume or rate.
- Consider potentially reversible causes by recalling the H's and T's listed in Part 5.

ROSC

If resuscitative efforts successfully restore an organized rhythm (or you find other evidence of ROSC, such as an abrupt and sustained increase in PETCO₂ or visible pulsations on an arterial waveform), check the child's pulse for a perfusing rhythm. If a pulse is present, continue with post-cardiac arrest care.

Pediatric Cardiac Arrest: Special Circumstances

The following special circumstances resulting in pediatric cardiac arrest require specific management:

- Trauma
- Drowning
- Anaphylaxis
- Poisoning
- Congenital heart disease: single ventricle
- Pulmonary hypertension

Cardiac Arrest Due to Trauma

Trauma-associated cardiac arrest in children represents a significant subgroup of pediatric OHCA. Improper resuscitation (including inadequate volume resuscitation) is a major cause of preventable pediatric trauma deaths. Despite rapid and effective out-of-hospital and trauma center response, the survival rate is low for pediatric OHCA due to trauma and very low for pediatric OHCA due to blunt trauma. Factors that may improve outcome from traumatic OHCA include treatable penetrating injuries and prompt transport (typically 10 minutes or less) to a trauma care facility.

Traumatic cardiac arrest in children has multiple possible causes, including

- Hypoxia secondary to respiratory arrest, airway obstruction, or tracheobronchial injury
- Injury to vital structures (eg, heart, aorta, pulmonary arteries)
- Severe brain injury with secondary cardiovascular collapse
- Upper cervical spinal cord injury with respiratory arrest, which may be accompanied by spinal shock, progressing to cardiac arrest
- Diminished cardiac output or PEA from tension pneumothorax, cardiac tamponade, or massive hemorrhage

Basic and advanced life support techniques for the pediatric trauma victim in cardiac arrest are fundamentally the same as for the child with nontraumatic cardiac arrest: support of circulation, airway, and breathing. The focus of a resuscitation attempt in an out-of-hospital setting is to

- Provide high-quality CPR, if needed
- Maintain adequate airway, ventilation, and circulation
- Anticipate airway obstruction by dental fragments, blood, or other debris (use a suction device, if necessary)
- Minimize cervical spine motion (if indicated), while opening the airway and providing ventilation
- Control external bleeding with pressure by using a tourniquet and/or hemostatic dressings as indicated
- Minimize interventions that delay transport to definitive care
- Transport infants and children with multisystem trauma to a trauma center with pediatric expertise
- Establish IV/IO access and initiate volume resuscitation as needed

The following is a summary of key management principles for traumatic cardiac arrest in children (Table 24).

Table 24. Managing Traumatic Cardiac Arrest

Technique	Intervention
CPR	<ul style="list-style-type: none"> Perform high-quality CPR. Attach a monitor/defibrillator. Provide defibrillation as needed. Control visible hemorrhage with direct pressure or tourniquet.
Airway	<ul style="list-style-type: none"> Open and maintain the airway by using a jaw-thrust maneuver. Restrict cervical spine motion by manually stabilizing the head and neck if cervical spine injury is suspected.
Breathing	<ul style="list-style-type: none"> Avoid excessive ventilation. Ventilate with a bag-mask device using 100% O₂; use a 2-person bag-mask ventilation technique to maintain manual stabilization of the head and neck (if spinal motion restriction is indicated). If an advanced airway is inserted, 1 rescuer should stabilize the head and neck in a neutral position (if spinal motion restriction is indicated). Perform unilateral or bilateral needle decompressions for suspected tension pneumothoraxes. Seal any significant open pneumothorax and insert a thoracotomy tube.
Circulation	<ul style="list-style-type: none"> Assume that the patient is hypovolemic; establish IV/IO access and replace fluids rapidly (<i>do not place an IO in any bone that might be fractured</i>). Consider non-crossmatched type O-negative blood for female patients (and either O-negative or O-positive for male patients). Consider pericardiocentesis for possible cardiac tamponade (if suspected). Consider spinal shock (ie, loss of sympathetic innervation) resulting in fluid-refractory hypotension and bradycardia. Vasopressor therapy is indicated if spinal shock is suspected.

Cardiac Arrest Due to Drowning

Immediate high-quality CPR is the single most important factor influencing survival in drowning (Table 25). You may have difficulty performing chest compressions while the victim is still in the water, but you can ventilate immediately if you are trained to do so. Start chest compressions as soon as you can do so safely, and when the child is lying faceup on a firm surface.

Table 25. Managing Cardiac Arrest Due to Drowning

Technique	Intervention
CPR	<ul style="list-style-type: none"> Perform high-quality CPR. Attach a monitor/defibrillator; provide rapid defibrillation if needed. If defibrillation is indicated and the chest is covered with water, quickly wipe the child's chest to minimize electrical arcing between the defibrillation pads or paddles.
Airway	<ul style="list-style-type: none"> Open the airway. If there is reason to suspect a cervical spine injury (eg, diving injury), restrict spinal motion. If an advanced airway is inserted, one rescuer should stabilize the head and neck in a neutral position.
Breathing	<ul style="list-style-type: none"> Ventilate with a bag-mask device by using 100% O₂. Be prepared to suction the airway, because drowning victims often vomit swallowed water; decompress the stomach with a nasogastric/orogastric tube after the advanced airway is inserted.
Circulation	<ul style="list-style-type: none"> Continue high-quality CPR if needed.
Exposure	<ul style="list-style-type: none"> Evaluate core body temperature and attempt rewarming if the child is severely hypothermic (core temperature <30°C).

Except for consideration of cervical spine injury and hypothermia as contributing factors, BLS and advanced life support techniques for cardiac arrest due to drowning are fundamentally the same as for other children in cardiac arrest: support of airway, breathing, and circulation. Rescuers should remove drowning victims from the water and begin the resuscitation attempt as quickly as possible.

In hypothermia-associated cardiac arrest, it is often difficult to know when to terminate resuscitative efforts. Victims of drowning in icy water can possibly survive after prolonged submersion (as long as 40 minutes) and CPR (longer than 2 hours). When drowning occurs in icy water, experts recommend rewarming to a core temperature of at least 30°C before abandoning CPR efforts. The heart may not respond to resuscitative efforts until it achieves this core temperature.

Extracorporeal circulation is the most rapid and effective technique for rewarming severely hypothermic cardiac arrest victims after submersion in icy water. Although you may rewarm the patient with passive techniques or body cavity irrigation in the out-of-hospital or community hospital setting, you should rapidly transfer to a facility that is capable of performing pediatric ECPR/extracorporeal membrane oxygenation.

Cardiac Arrest Due to Anaphylaxis

Near-fatal anaphylaxis produces airway edema and obstruction and profound vasodilation, which significantly increases intravascular capacity and produces relative hypovolemia. Anaphylaxis often accompanies bronchoconstriction, which compromises oxygenation and further impairs tissue O₂ delivery. If cardiac arrest develops, provide CPR as the primary therapy and establish and maintain an adequate airway, bolus fluid administration, and epinephrine. Children with anaphylaxis are often young with healthy hearts and cardiovascular systems. They may respond to an airway with adequate oxygenation and ventilation and rapid correction of vasodilation and low intravascular volume. Effective CPR may maintain sufficient O₂ delivery until the catastrophic effects of the anaphylactic reaction resolve.

Managing cardiac arrest that results from anaphylaxis may include the following critical therapies ([Table 26](#)):

Table 26. Managing Cardiac Arrest Due to Anaphylaxis

Technique	Intervention
CPR	<ul style="list-style-type: none"> Perform high-quality CPR and rapid defibrillation as needed.
Airway	<ul style="list-style-type: none"> Open and maintain the airway by using manual maneuvers. If you perform ET intubation, be prepared for possible airway edema and the need to use a smaller ET tube than predicted by the child's age or length.
Breathing	<ul style="list-style-type: none"> Perform bag-mask ventilation using 100% O₂ until an advanced airway is inserted.
Circulation	<ul style="list-style-type: none"> Administer boluses of isotonic crystalloid as needed to treat shock. Insert 2 large-bore IVs or 2 IO catheters. Administer epinephrine in standard doses (0.01 mg/kg IV/IO [0.1 mL/kg of 0.1 mg/mL concentration]) or via the ET tube if you cannot obtain vascular access (0.1 mg/kg [0.1 mL/kg of 1 mg/mL concentration]) every 3-5 minutes during cardiac arrest. Infuse with epinephrine as needed.

Antihistamines and corticosteroids are considered adjuncts in the treatment of anaphylaxis, with varying degrees of evidence to support their use. In the setting of cardiac arrest due to anaphylaxis, the most important medication to give, and to continue to give if indicated, is epinephrine. If ROSC is achieved, administering antihistamines or corticosteroids, or both, can then be considered.

Cardiac Arrest Associated With Poisoning

Drug overdose or poisoning may cause cardiac arrest as a result of direct cardiac toxicity or the secondary effects of respiratory depression, airway obstruction and possible respiratory arrest, peripheral vasodilation, arrhythmias, and hypotension. The poisoning victim often has a healthy myocardium, but temporary cardiac dysfunction may be present until the effects of the drug or toxin have been reversed or metabolized. This takes a variable amount of time, often several hours, depending on the nature of the toxin, drug, or poison. Because the toxicity may be temporary, prolonged resuscitation efforts and use of advanced support techniques such as extracorporeal life support may result in good long-term survival.

Initiate advanced life support measures according to the Pediatric Cardiac Arrest Algorithm. Check glucose as soon as possible. If the patient is in cardiac arrest and hypoglycemic (eg, from a β -blocker or alcohol overdose), normalize glucose as soon as possible to improve the chances of a successful cardiac and neurologic outcome. PALS treatment for victims of a suspected poisoning should include searching for and treating reversible causes. We recommend early consultation with a poison control center or toxicologist.

Congenital Heart Disease: Single Ventricle

While the prevalence of complex cyanotic congenital heart disease in children is low, those with single ventricle (tricuspid/pulmonary atresia, hypoplastic left heart syndrome, and their variants) represent a large proportion of children who have cardiac arrest, particularly in the in-hospital setting. More infants and children are surviving after palliative surgical procedures; these patients may require resuscitation postoperatively or during readmission for critical illness.

Single-ventricle physiology is complex and varies with the specific lesion and stage of surgical repair. Therefore, it is important to obtain a history from the caretakers to determine the child's baseline hemodynamic status and arterial O₂ saturation. During cardiac arrest, standard resuscitation care is indicated for all infants with single-ventricle anatomy after stage I (Norwood) palliation or those with a univentricular heart and aortopulmonary shunt to provide pulmonary blood flow. In addition to standard resuscitation, specific measures include the following:

- Consider administering heparin for children with aortopulmonary or right ventricular-pulmonary artery shunt if you are concerned about shunt openness/patency.
- After resuscitation, titrate administered O₂ to achieve an O₂ saturation appropriate to maintain the optimum pulmonary-to-systemic blood flow ratio and adequate systemic perfusion and oxygenation (must be individualized for each patient).
- Do not rely on PETCO₂ to indicate CPR quality in a single-ventricle patient because pulmonary blood flow in these patients does not always reflect cardiac output (ie, it is influenced by additional factors).
- Consider permissive hypoventilation strategies or even negative-pressure ventilation in the periarrest state to improve cardiac output.
- Consider extracorporeal life support or extracorporeal membrane oxygenation for patients in cardiac arrest who have undergone stage I palliation (Norwood) or Fontan-type procedures.

Pulmonary Hypertension

In pulmonary hypertension, increased resistance to blood flow through the lungs may impair cardiac output, so follow standard PALS recommendations during cardiac arrest. Other measures include the following:

- Correct hypercarbia and acidosis if present.
- Consider giving a bolus of isotonic crystalloid (eg, normal saline) to maintain ventricular preload.
- If the patient received pulmonary vasodilators such as nitric oxide or prostacyclin immediately before the arrest, be sure that drug administration continues.
- Consider administering inhaled nitric oxide or prostacyclin (or IV prostacyclin) to reduce pulmonary vascular resistance.
- Consider instituting ECPR early during resuscitation.



Critical Concepts: ECPR

You may consider ECPR for pediatric patients with cardiac diagnoses who have IHCA in settings with existing extracorporeal membrane oxygenation protocols, expertise, and equipment. While evidence shows no overall benefit to using ECPR, observational data from a registry of pediatric IHCA showed improved survival to hospital discharge with the use of ECPR in patients with surgical cardiac diagnoses. For children with underlying cardiac disease, when ECPR is initiated in a critical care setting, long-term survival has been reported after prolonged CPR. When you use ECPR during cardiac arrest, the outcome for children with underlying cardiac disease is better than for those with noncardiac disease.

Part 6: Effective High-Performance Team Dynamics

Although a single CPR-trained bystander can resuscitate a patient within the first moments after collapse, most attempts require the concerted efforts of multiple healthcare providers to perform various interventions simultaneously. Effective teamwork divides the tasks while multiplying the chances of a successful outcome.

Successful high-performance teams not only have medical expertise and mastery of resuscitation skills, but they also demonstrate effective communication and team dynamics. This Part discusses the importance of team roles, behaviors of effective Team Leaders and team members, and elements of effective high-performance team dynamics.

Learning Objective

After completing this Part, you should be able to apply high-performance team dynamics to increase your patients' chance of survival.

During the course you will participate as both the Team Leader and a team member during case simulations, modeling the behaviors discussed in this Part.



Critical Concepts: Understanding Team Roles

Whether you are a team member or Team Leader during a resuscitation attempt, you should understand your role and other team members' roles. This will help you anticipate

- The actions that will be performed next
 - How to communicate and work as a team member or Team Leader
-

Roles in a High-Performance Team

Team Leader Role

Every high-performance team needs a leader to organize group efforts and make sure everything is done at the right time in the right way by monitoring and integrating individual team members' performances.

The Team Leader also models excellent team behavior and leadership skills for the team and others involved or interested in the resuscitation attempt and serves as a teacher or guide to help train future Team Leaders and improve team effectiveness. After resuscitation, the Team Leader can facilitate analysis, critique, and practice to prepare for the next resuscitation attempt. The Team Leader also helps team members understand why they must perform certain tasks in a specific way.

The Team Leader should be able to explain the importance of

- Pushing hard and fast in the center of the chest
- Ensuring complete chest recoil
- Minimizing interruptions in chest compressions
- Avoiding excessive ventilation

Whereas members of a high-performance team should focus on their individual tasks, the Team Leader must focus on comprehensive patient care.

Team Member Roles

Members of a high-performance team must be proficient in performing the skills authorized by their scope of practice. For a resuscitation attempt to be successful, team members should be

- Clear about role assignments
- Prepared to fulfill their role responsibilities
- Well practiced in resuscitation skills

The diagram illustrates the timeline of events for a cardiac arrest scenario. A vertical blue line marks time in seconds. Icons represent the arrival of the HCP (green person), AED (red person), and Resuscitation team (blue group of people). Actions and medications are listed to the right of the timeline, color-coded to match the icons. Arrows at the bottom indicate the direction of flow for each group.

Time (s)	Event	Action/Medication
0s	HCP 1 arrives	Verifies scene safety Shouts for help
40s	HCP 2 arrives with AED	Compressor/Airway
60s		AED/Monitor/Defibrillator
70s	HCP 3 arrives	Compressor
		AED/Monitor/Defibrillator
		IV/IO Medications
120s		Compressor
		AED/Monitor/Defibrillator
		IV/IO Medications
240s	Resuscitation team arrives	Compressor
250s		AED/Monitor/Defibrillator
		IV/IO Medications

Not only should everyone on the team know his or her own limitations and capabilities, but the Team Leader should also be aware of them. This allows the Team Leader to evaluate team resources and call for backup when needed. High-performance team members should anticipate these situations in which they might require assistance and inform the Team Leader.

During the stress of an attempted resuscitation, do not practice or explore a new skill, especially without seeking advice from more experienced personnel. If you need extra help, request it early rather than wait until the patient deteriorates to the point that help is critical. Asking for help is not a sign of weakness or incompetence; it is better to have more help than needed than not enough help, which might negatively affect patient outcome.

The Team Leader and team members should also accept offers from others to carry out an assigned task they are unable to complete, especially if completing the task is essential.

Constructive Interventions

During a resuscitation attempt, the Team Leader or a team member may need to intervene against inappropriate actions.

Because tactful constructive intervention is necessary, the Team Leader should avoid confrontation with team members. Instead, he or she should conduct a debriefing afterward for constructive criticism. The Team Leader should also ask that a different intervention be started if it has a higher priority and reassign a team member who is trying to function beyond his or her skill level. Conversely, team members should confidently question a colleague who is about to make a mistake and suggest an alternative drug or dose, if necessary.

Communication

Knowledge Sharing

Sharing information is critical to effective team performance. Team Leaders may become trapped in a specific treatment or diagnostic approach, a common human error called a *fixation error*. Examples of 3 common types of fixation errors are

- “Everything is OK.”
- “This and only this is the correct path.”
- “Do anything but this.”

When resuscitative efforts are ineffective, go back to the basics and talk as a team, with conversations like, “Well, we’ve observed the following on the Primary Assessment... Have we missed something?” Members of a high-performance team should inform the Team Leader of any changes in the patient’s condition to ensure that decisions are made with all available information. The Team Leader and team members should all examine clinical signs that are relevant to the treatment.

The Team Leader should encourage information sharing and ask for suggestions if the next best interventions are unclear, ask for ideas for differential diagnoses, and ask if anything has been overlooked (eg, if IV access should have been obtained or drugs should have been administered). Team members should share information with other team members and the Team Leader.

Summarizing and Reevaluating

An essential role of the Team Leader is monitoring and reevaluating

- The patient’s status
- Interventions that have been performed
- Assessment findings

The Team Leader should summarize this information out loud in periodic updates to the team. He or she should review the status of the resuscitation attempt and announce the next few steps. However, the patient’s condition can change, so the Team Leader should remain flexible to changing treatment plans and revisiting the initial differential diagnosis. When a change is needed, the Team Leader should inform arriving personnel of the current status and plans for further action. He or she should ask for information and summaries from the Timer/Recorder as well.

The Team Leader should draw continuous attention to decisions about differential diagnoses and review or maintain an ongoing record of drugs and treatments administered and the patient’s response. The Team Leader and team members should clearly communicate significant changes in the patient’s clinical condition and increase monitoring (eg, frequency of respirations and blood pressure) when the patient’s condition deteriorates.

Closed-Loop Communication

When communicating with high-performance team members, the Team Leader should use closed-loop communication by taking these steps:

1. The Team Leader gives a message, order, or assignment to a team member.
2. By receiving a clear response and eye contact, the Team Leader confirms that the team member heard and understood the message.
3. The Team Leader listens for the team member to confirm task performance before assigning another task.

The Team Leader should only give more tasks to a team member if the Team Leader has asked for or received confirmation of a completed assignment. Correspondingly, team members should verbally confirm the order with the Team Leader before giving drugs and should inform the Team Leader after giving the drug or performing the procedure.

For example, the Team Leader assigns another task after receiving oral confirmation that a task has been completed, such as, “Now that the IV is in, give 1 mg of epinephrine.” Team members close the loop by informing the Team Leader when a task begins or ends, such as, “The IV is in.”

Clear Messages

Unclear communication can cause unnecessary treatment delays or medication errors. Therefore, all healthcare providers should deliver clear, concise messages and orders in a calm, articulate, and direct manner without yelling or shouting, which can impair effective high-performance team interaction. Only one person should talk at a time.

The Team Leader should encourage team members to speak clearly, and team members should repeat medication orders and question an order if they have any doubts.

Mutual Respect

The best high-performance teams comprise members who share a mutual respect for each other and work together in a collegial, supportive manner. To have a high-performance team, everyone must abandon ego and respect each other during the resuscitation attempt, regardless of any additional training or experience that the Team Leader or specific team members may have.

The Team Leader and team members should speak in a friendly, controlled tone of voice and avoid shouting or displaying aggression if they are misunderstood initially. When one person raises his voice, others will respond similarly, behave aggressively or confuse directive behavior with aggression, or be uninterested in others. Instead, the Team Leader should acknowledge correctly completed assignments by saying, “Thanks—good job!”

Part 7: Recognizing Respiratory Distress and Failure

Respiratory distress is defined as abnormal respiratory rate or effort, encompassing a spectrum of signs from tachypnea with retractions to agonal gasps. It includes

- Increased work of breathing
- Inadequate respiratory effort (eg, shallow respirations) or rate (bradypnea)
- Irregular breathing

PALS providers must identify respiratory conditions that are treatable with simple measures, such as by clearing airway secretions or administering O₂. Yet it may be even more important to identify those respiratory conditions that are subtly but rapidly progressing toward respiratory failure. These conditions require timely interventions with more advanced airway techniques (eg, assisted bag-mask ventilation).

In infants and children, respiratory distress can quickly progress to respiratory failure and finally to cardiac arrest. Good outcome (ie, neurologically intact survival to hospital discharge) is more likely after respiratory arrest than after cardiac arrest. Once the child with a respiratory problem develops cardiac arrest, the outcome is often poor, so you can greatly improve outcome by identifying and managing respiratory distress and failure before the child deteriorates to cardiac arrest.

Learning Objective

After completing this Part, you should be able to differentiate between respiratory distress and failure.

You need to understand the concepts in this Part to be able to quickly identify signs of respiratory distress and respiratory failure as well as be able to recognize respiratory problems by type so that you can choose appropriate interventions.

Fundamental Issues Associated With Respiratory Problems

Children with respiratory problems have impaired oxygenation, ventilation, or both. This section discusses

- Impaired oxygenation and ventilation
- Physiology of respiratory disease

Impaired Oxygenation and Ventilation in Respiratory Problems

Physiology of the Respiratory System

The main function of the respiratory system is gas exchange, in which air is taken into the lungs with inspiration. From there, O₂ diffuses from the alveoli into the blood, where some O₂ dissolves in plasma. Most O₂ that enters the blood is bound to hemoglobin. The percent of hemoglobin that is bound to O₂ is called *oxygen saturation*. When blood passes through the lungs, CO₂ diffuses from the blood into the alveoli and is exhaled. Acute respiratory problems can result from alterations in any part of this system, from the airway to the alveoli (lung parenchyma). Central nervous system disease, such as seizures or head trauma, can impair respiration control, leading to decreased respiratory rate. Muscle weakness, either primary (eg, muscular dystrophy) or secondary (eg, fatigue), may also impair oxygenation or ventilation.

Children have a high metabolic rate, so O₂ demand per kilogram of body weight is high. Therefore, hypoxemia and tissue hypoxia can develop more rapidly in a child than in an adult if apnea or inadequate alveolar ventilation occurs.

Respiratory problems can result in

- Hypoxemia
- Hypercarbia
- A combination of both hypoxemia and hypercarbia

Hypoxemia (Low Oxygen Saturation)

Hypoxemia is defined as a decreased arterial oxygen saturation detected by pulse oximetry or directly measuring oxygen saturation in an arterial blood gas sample. It is generally defined as arterial oxygen saturation below 94% in a normal child breathing room air. Various conditions may lower the threshold, such as altitude or cyanotic heart disease. Permissive hypoxemia is a pulse oximetry percentage of less than 94%, which may be appropriate in certain circumstances (eg, some cases of congenital heart disease).

It is important to distinguish between *hypoxemia* and *tissue hypoxia*. Hypoxemia is low arterial O₂ saturation (SaO₂ less than 94%), while hypoxia is a pathologic condition in which the body as a whole (*generalized hypoxia*) or a region of the body (*tissue hypoxia*) is deprived of an adequate oxygen supply. Note that hypoxemia does not necessarily lead to tissue hypoxia and that tissue hypoxia may occur when arterial oxygen saturation is normal. For example, when hypoxemia is chronic (eg, unrepaired cyanotic heart disease), compensatory increases in blood flow (ie, increased cardiac output) or hemoglobin concentration (polycythemia) increase the O₂-carrying capacity of the blood and help maintain arterial O₂ content at near-normal concentrations despite hemoglobin saturation being low. This can help maintain oxygen delivery and tissue oxygenation even when hypoxemia is present. Conversely, if tissue perfusion is poor or the patient has severe anemia, tissue hypoxia may occur with normal arterial oxygen saturation.

In response to tissue hypoxia, the child may initially compensate by increasing respiratory rate and depth, or hyperventilate. *Hyperventilation* refers to increased alveolar ventilation resulting in a decrease in PaCO₂ to less than 35 mm Hg. This may be caused by an increased respiratory rate, increased tidal volume, or combination of both. To ensure that PaCO₂ does not decrease below 30 mm Hg, you must use capnography to guide hyperventilation. Tachycardia may also develop in response to hypoxemia to increase cardiac output. As tissue hypoxia worsens, these signs of cardiopulmonary distress become more severe ([Table 27](#)).

Table 27. Signs of Tissue Hypoxia

Early signs of tissue hypoxia	Late signs of tissue hypoxia
Fast respiratory rate (tachypnea)	Slow respiratory rate (bradypnea), inadequate respiratory effort, apnea
Increased respiratory effort: nasal flaring, retractions	Increased respiratory effort: head bobbing, seesaw respirations, grunting
Tachycardia	Bradycardia
Pallor, mottling, cyanosis*	Pallor, mottling, cyanosis*
Agitation, anxiety, irritability	Decreased level of consciousness

*Pallor, mottling, and cyanosis can appear as both early and late signs.



Critical Concepts: Arterial O₂ Content

Arterial O₂ content is the total amount of O₂ carried in the blood (in milliliters O₂ per deciliter of blood). It is the sum of the quantity of O₂ bound to hemoglobin plus the O₂ dissolved in arterial blood. O₂ content is determined largely by the hemoglobin (Hgb) concentration (grams per deciliter) and its saturation with O₂ (SaO₂). Use the following equation to calculate arterial O₂ content: **Arterial O₂ Content = [1.36 × Hgb concentration × Sao₂] + (0.003 × Pao₂)**

Under normal conditions, dissolved O₂ (0.003 × Pao₂) is an inconsequential portion of the total arterial O₂ content. But an increase in dissolved O₂ can produce a relatively significant increase in arterial O₂ content for a child with severe anemia.

Hypoxemia can be caused by several different mechanisms leading to respiratory distress and failure ([Table 28](#)).

Table 28. Mechanisms of Hypoxemia

Factor	Causes	Mechanism	Treatment
Low atmospheric Po₂	High altitude (decreased barometric pressure)	Decreased PaO ₂	Give supplemental O ₂
Alveolar hypoventilation	<ul style="list-style-type: none"> Central nervous system infection Traumatic brain injury Drug overdose Neuromuscular weakness Apnea 	Increased alveolar CO ₂ tension (hypercarbia) displaces alveolar O ₂ , resulting in decreased alveolar O ₂ tension (PaO ₂)	Restore normal ventilation; supplemental O ₂

Diffusion defect	Interstitial pneumonia	Impaired movement of O ₂ and CO ₂ between the alveoli and blood results in decreased PaO ₂ (hypoxemia) and, if severe, an increased PaCO ₂ (hypercarbia).	Apply noninvasive ventilation (either continuous positive airway pressure [CPAP] or biphasic/bilevel CPAP)
Ventilation/perfusion (V/Q) mismatch	<ul style="list-style-type: none"> • Pneumonia • Pulmonary edema • Atelectasis • Acute respiratory distress syndrome • Asthma • Bronchiolitis • Foreign body 	Mismatch of ventilation and perfusion: blood flow through areas of the lung that are inadequately ventilated results in incomplete oxygenation of the blood returning to the left side of the heart. The result is a decreased arterial O ₂ saturation and PaO ₂ and, to a lesser extent, increased PaCO ₂ .	Use positive end-expiratory pressure to increase mean airway pressure [*] ; supplemental O ₂ ; ventilatory support
Right-to-left shunt	<ul style="list-style-type: none"> • Cyanotic congenital heart disease • Extracardiac (anatomical) vascular shunt • Same causes listed for V/Q imbalance[‡] 	Shunting of unoxygenated blood from the right side of the heart to the left (or from the pulmonary artery into the aorta) results in a low PaO ₂ . Effects are similar to right-to-left shunt in the lungs.	Correct defect (supplemental O ₂ alone is insufficient)

^{*}Positive end-expiratory pressure should be used carefully in children with asthma; expert consultation is advised.

[‡]With pneumonia, acute respiratory distress syndrome, and other lung tissue diseases, the pathophysiology is often characterized by a mix of hypoxemia mechanisms. The most extreme form of V/Q mismatch would be a segment of lung with blood flow (Q) but no ventilation (V), a situation in which the blood would not become oxygenated. When it returns to the left side of the heart, it mixes with the oxygenated blood, resulting in a lower O₂ saturation. The degree of desaturation depends on the size of the unventilated lung segment.

Hypercarbia (Inadequate Ventilation)

Hypercarbia is an increased CO₂ tension in the arterial blood (PaCO₂). When hypercarbia is present, ventilation is inadequate.

CO₂ is a by-product of tissue metabolism. Normally, the lungs eliminate it to maintain acid-base homeostasis because, when ventilation is inadequate, CO₂ elimination is inadequate. The resulting increase in PaCO₂ causes the blood to become acidic (respiratory acidosis). Causes of hypercarbia are

- Airway obstruction (upper or lower)
- Lung tissue disease
- Decreased or inadequate respiratory effort (central hypoventilation)

Most children with hypercarbia present with respiratory distress and tachypnea. Children may become tachypneic in an attempt to eliminate excess CO₂. However, the child with hypercarbia may also present with poor respiratory effort, including decreased respiratory rate. In this case, hypercarbia results from inadequate ventilation secondary to impaired respiratory drive. This inadequate ventilation may result from drugs, such as a narcotic overdose. It may also result from a central nervous system disorder with respiratory muscle weakness preventing the development of compensatory tachypnea. Detecting an inadequate respiratory drive requires careful observation and assessment. Consequences of inadequate ventilation become more severe as the PaCO₂ increases and respiratory acidosis worsens.

Detecting Hypercarbia

Hypercarbia is more difficult to detect than hypoxemia because it produces no obvious clinical sign, such as cyanosis. Precisely measuring PCO₂ requires a blood sample (arterial, capillary, or venous). Exhaled CO₂ detectors are now available for use in children with or without advanced airways. The end-tidal CO₂ (or exhaled CO₂ measured at the end of exhalation) measured by capnography may not be identical to the arterial CO₂. However, if the airway is open/patent, there is no increased dead space from air trapping (eg, asthma), and cardiac output is adequate, the end-tidal CO₂ will increase in the presence of hypercarbia.

When to Suspect Hypercarbia

Decreased level of consciousness is a critical symptom of both inadequate ventilation and hypoxia. If a child's clinical condition deteriorates from agitation and anxiety to decreased responsiveness despite supplemental O₂, this may indicate that the PaCO₂ is rising. Note that even if the pulse oximeter indicates adequate O₂ saturation, ventilation may be impaired and hypercarbia may be present. If a child with respiratory distress has a decreased level of consciousness despite adequate oxygenation, ventilation may be inadequate and hypercarbia and respiratory acidosis may be present.

Signs of inadequate ventilation are nonspecific and include one or more of the following:

- Tachypnea or inadequate respiratory rate for age and clinical condition
- Nasal flaring, retractions
- Change in level of consciousness: initial anxiety, agitation, and then decreased level of consciousness

Physiology of Respiratory Disease

One may accomplish normal spontaneous breathing with minimal work. Breathing is quiet with unlabored, smooth inspiration and passive expiration. In children with respiratory disease, "work of breathing" becomes more apparent. Important factors associated with increased work of breathing include

- Increased airway resistance (upper and lower)
- Decreased lung compliance
- Use of accessory muscles of respiration
- Disordered central nervous system control of breathing

Airway Resistance


Airway resistance, or the impedance to airflow within the airways, is primarily increased by reducing the size of the conducting airways (either by airway constriction or inflammation). Turbulent airflow also causes increased airway resistance. Airflow may become turbulent when the flow rate increases, even if the airway size remains unchanged. When airway resistance increases, work of breathing increases in an attempt to maintain airflow.

Larger airways provide lower resistance to airflow than smaller airways. Airway resistance decreases as lung volume increases (inflation) because airway dilation accompanies lung inflation. Conditions such as edema, bronchoconstriction, secretions, mucus, or a mediastinal mass impinging on large or small airways can decrease airway size, thereby increasing airway resistance. Resistance in the upper airway, particularly in the nasal or nasopharyngeal passages, can represent a significant portion of total airway resistance, especially in infants.

Airway Resistance in Laminar Airflow

During normal breathing, because airflow is laminar (ie, quiet, smooth, and orderly) and airway resistance is relatively low, only a small driving pressure (ie, difference in pressure between the atmosphere and the pleural space) is needed to produce adequate airflow. When airflow is laminar (quiet), resistance to airflow is inversely proportional to the fourth power of the airway radius, so even a small reduction in airway diameter results in a significant increase in airway resistance and work of breathing.

Figure 26 shows normal airways on the left and edematous airways (with 1 mm of circumferential edema) on the right. Resistance to flow is inversely proportional to the fourth power of the radius of the airway lumen for laminar flow and to the fifth power of the radius when air flow is turbulent. The net result is a 75% decrease in cross-sectional area and a 16-fold increase in airway resistance in the infant vs a 44% decrease in cross-sectional area and a 3-fold increase in airway resistance in the adult during quiet breathing. Turbulent flow in an infant (eg, crying) increases airway resistance and thus the work of breathing from 16- to 32-fold.



Airway Resistance in Turbulent Airflow

When airflow is turbulent (ie, irregular), resistance is inversely proportional to the fifth power of the radius of the airway lumen, a 10-fold increase over the resistance to airflow that occurs during reduction in the airway radius when airflow is normal and laminar. In this state, a much larger driving pressure is required to produce the same rate of airflow. Therefore, patient agitation (causing rapid, turbulent airflow) results in a much greater increase in airway resistance and work of breathing than with quiet, laminar flow. To prevent turbulent airflow from generating (eg, during crying), try to keep a child with airway obstruction as calm as possible.

Lung Compliance

Compliance refers to the distensibility of the lung, chest wall, or both. *Lung compliance*, specifically, is defined as the change in lung volume produced by a change in driving pressure across the lung. When lung compliance is high, the lungs easily inflate (a large change in volume produced by a slight change in driving pressure). The lungs are stiffer in a child with low lung compliance, so it takes more effort to inflate them. To create a significant pressure gradient to produce air flow into the stiff lung, the diaphragm contracts more forcefully, increasing intrathoracic volume and reducing intrathoracic pressure. Poorly compliant lungs will also lead to increased work of breathing. During mechanical ventilation, increased positive airway pressure is needed to achieve adequate ventilation when lung compliance decreases.

Causes of decreased compliance include extrapulmonary conditions like pneumothorax and pleural effusion as well as pneumonia and inflammatory lung tissue diseases, such as ARDS, fibrosis, and pulmonary edema. These conditions are associated with an increase in the water content in the interstitial space and alveoli. The impact of this on lung compliance is similar to the expansion that occurs when a sponge becomes saturated with water. A normal sponge re-expands quickly when compressed, while a wet sponge is harder to compress and re-expands more slowly because its normal elasticity is opposed by the extra weight of the fluid in the sponge.

The chest wall in infants and young children is compliant. Therefore, relatively small pressure changes can move the chest wall. During normal breathing, diaphragm contraction in infants pulls the lower ribs slightly inward but does not cause significant chest retraction. However, forcefully contracting the diaphragm results in a large drop in pressure within the chest, pulling the child's compliant chest inward (ie, retracting it) and pushing the abdomen out during inspiration, which can actually limit lung expansion during inspiration. These muscular contractions are useful in older children and adolescents when the chest wall is more rigid, but in infants and young children, the compliant chest wall (which is not fully calcified) actually can be counterproductive.

Children with neuromuscular disorders have a weak chest wall and weak respiratory muscles that can make breathing and coughing ineffective. Muscle weakness can result in characteristic seesaw breathing (the chest wall retracts and the abdomen expands simultaneously).

Inspiratory and Expiratory Flow

The inspiratory muscles of respiration include the diaphragm, intercostal muscles, and accessory muscles (primarily of the abdomen and neck). During spontaneous breathing, inspiratory muscles (chiefly the diaphragm) increase intrathoracic volume, resulting in decreased intrathoracic pressure. When intrathoracic pressure is less than atmospheric pressure, air flows into the lungs (*inspiration*); the intercostal muscles stabilize the chest wall to optimize intrathoracic inspiratory air flow. Accessory muscles of respiration are not typically needed during normal ventilation; however, in respiratory disorders that increase airway resistance or reduce lung compliance, activation of these accessory muscles helps to improve the child's ability to augment inspiratory air flow.

Expiratory flow results from the inspiratory muscles relaxing and the lung and chest wall elastically recoiling, decreasing intrathoracic volume and, thus, increasing intrathoracic pressure to a higher level than that of the atmospheric pressure. During spontaneous breathing, expiratory flow is primarily a passive process. However, expiration may become an active process in the presence of increased lower airway resistance and may require the abdominal wall muscles and the intercostal muscles.



Critical Concepts: The Diaphragm's Role in Breathing

The normal diaphragm is dome-shaped and contracts most forcefully in this shape. When the diaphragm flattens, as occurs with lung hyperinflation (eg, acute asthma), contraction is less forceful and ventilation is less efficient. Respiration is compromised if diaphragm movement is impeded by abdominal distention and high intra-abdominal pressure (eg, gastric inflation) or by air trapping caused by airway obstruction. During infancy and early childhood, the

intercostal muscles serve primarily to stabilize the chest wall. They cannot effectively lift the chest wall to increase intrathoracic volume and compensate for loss of diaphragm motion.

Central Nervous System Control of Breathing

Breathing is controlled by complex mechanisms involving

- Brainstem respiratory centers
- Central and peripheral chemoreceptors
- Voluntary control

A group of respiratory centers located in the brainstem control spontaneous breathing. Voluntary control from the cerebral cortex, such as breath holding, panting, and sighing, can also override breathing. Conditions like infection of the central nervous system, traumatic brain injury, and drug overdose can impair respiratory drive, resulting in hypoventilation or even apnea.

Note that central chemoreceptors respond to changes in the hydrogen ion concentration of cerebrospinal fluid, which is largely determined by the arterial CO_2 tension (PaCO_2). Peripheral chemoreceptors (eg, in the carotid body) respond primarily to a decrease in arterial O_2 (PaO_2); some receptors also respond to an increase in PaCO_2 .

Identifying Respiratory Problems by Severity

Identifying the severity of a respiratory problem will help you decide the most appropriate interventions. Be alert for signs of

- Respiratory distress
- Respiratory failure

Respiratory Distress

Respiratory distress is a clinical state characterized by increased respiratory rate and effort, spanning a spectrum from mild tachypnea with increased effort to severe distress with impending respiratory failure. A description of respiratory distress severity typically includes a description of the respiratory rate and effort, quality of breath sounds, and mental status. These indicators may vary in severity. Note that signs of severe respiratory distress can indicate respiratory failure.

- Mild respiratory distress
 - Mild tachypnea
 - Mild increase in respiratory effort (eg, nasal flaring, retractions)
 - Abnormal airway sounds (eg, stridor, wheezing)
- Severe respiratory distress
 - Marked tachypnea
 - Marked increase in respiratory effort (eg, nasal flaring, retractions)
 - Paradoxical thoracoabdominal breathing (eg, seesaw breathing)
 - Accessory muscle use (eg, head bobbing)
 - Abnormal airway sounds (eg, grunting)
 - Decreased level of consciousness (eg, less responsive)
- Impending respiratory arrest
 - Bradypnea, apnea, or respiratory pauses
 - Low oxygen saturation (hypoxemia) despite high-flow supplemental oxygen
 - Inadequate respiratory effort (eg, shallow respirations)
 - Decreased level of consciousness (eg, unresponsive)
 - Bradycardia

Respiratory distress is apparent when a child tries to maintain adequate gas exchange despite airway obstruction, reduced lung compliance, or lung tissue disease. As the child tires or as respiratory function or effort or both deteriorate, adequate gas exchange cannot be maintained. When this happens, clinical signs of respiratory failure develop.

Respiratory Failure

Respiratory failure is a clinical state of inadequate oxygenation, ventilation, or both. It is recognized typically by abnormal appearance (particularly an altered level of consciousness, which may be characterized by agitation or a depressed level of consciousness), poor color, and reduced responsiveness. Although respiratory failure often results from respiratory distress progression, it may occur with little or no respiratory effort. At times, recognizing respiratory failure requires laboratory data (eg, blood gas) to confirm the diagnosis. In other patients, the clinical examination suffices to identify respiratory failure.

Suspect *probable respiratory failure* if some of the signs in the following list are present.

- Signs of severe respiratory distress
 - Marked tachypnea
 - Increased or decreased respiratory effort
 - Poor distal air movement
 - Tachycardia
 - Low oxygen saturation (hypoxemia) despite high-flow oxygen administration
 - Cyanosis
- Signs of probable respiratory failure
 - Very rapid or inadequate respiratory rate or possible apnea
 - Significant, inadequate, or absent respiratory effort
 - Absent distal air movement
 - Extreme tachycardia; bradycardia often indicates life-threatening deterioration
 - Low oxygen saturation (hypoxemia) despite high-flow supplemental oxygen
 - Decreased level of consciousness
 - Cyanosis

Respiratory failure can result from upper or lower airway obstruction, lung tissue disease, and disordered control of breathing (eg, apnea or shallow, slow respirations). *When respiratory effort is inadequate, respiratory failure can occur without typical signs of respiratory distress.* Respiratory failure is a clinical state that *requires intervention* to prevent deterioration to cardiac arrest.

Strict criteria for respiratory failure is difficult to define because the baseline respiratory function of an infant or child may be abnormal. For example, an infant with cyanotic congenital heart disease and a baseline arterial O₂ saturation (SaO₂) of 75% is not in respiratory failure on the basis of low O₂ saturation. But the same degree of hypoxemia would be one sign of respiratory failure in a child with normal baseline cardiopulmonary physiology.

Auscultation With a Stethoscope

Use a stethoscope to auscultate the following points during a respiratory physical exam:

- Anterior (on either side of the sternum)
- Posterior
- Lateral (under the axillae)

For more information, refer to Chest Expansion and Air Movement in Part 4.

Identifying Respiratory Problems by Type

Respiratory distress or failure can be classified into one or more of the following types:

- Upper airway obstruction
- Lower airway obstruction
- Lung tissue disease
- Disordered control of breathing

Respiratory problems do not always occur one at a time. A child may have more than 1 cause of respiratory distress or failure, such as disordered control of breathing caused by a head injury and then pneumonia (lung tissue disease). A patient may also exhibit symptoms consistent with more than 1 class of respiratory abnormality.

Upper Airway Obstruction

Obstruction of the upper airways (ie, the airways outside the thorax) can occur in the nose, pharynx, or larynx and can range from mild to severe.

Causes of Upper Airway Obstruction

Common causes of upper airway obstruction are foreign-body aspiration (eg, food or a small object), and airway swelling (eg, anaphylaxis, tonsillar hypertrophy, croup, or epiglottitis). Other causes include a mass that compromises the airway lumen (eg, pharyngeal or peritonsillar abscess, retropharyngeal abscess, or tumor), thick secretions obstructing the nasal passages, or any congenital airway abnormality (eg, congenital subglottic stenosis) resulting in narrowing of the airway, or poor control of the upper airway due to a decreased level of consciousness. It also may be hospital acquired. For example, subglottic stenosis may develop secondary to trauma induced by endotracheal intubation.

Due to their small airway, infants and small children may easily develop upper airway obstruction. Because an infant's tongue is large in proportion to the oropharyngeal cavity, if the infant has a decreased level of consciousness, the muscles may relax and allow the tongue to fall back and obstruct the oropharynx. Infants also have a prominent occiput. If the infant with a decreased level of consciousness is supine, resting on the large occiput can cause flexion of the neck, resulting in upper airway obstruction.

Signs of Upper Airway Obstruction

The major clinical signs, such as stridor, hoarseness, or a change in voice or cry, typically occur during the inspiratory phase of the respiratory cycle. Inspiratory retractions, use of accessory muscles, and nasal flaring are often present. The respiratory rate is often only mildly elevated because upper airway obstruction is worse with faster breathing. Examples include foreign body obstruction, croup, and epiglottitis. Other signs of upper airway obstruction include

- Increased respiratory rate and effort
- Drooling, snoring, or gurgling sounds
- Poor chest rise
- Poor air entry on auscultation

Lower Airway Obstruction

Obstruction of the lower airways (ie, the airways within the thorax) can occur in the lower trachea, the bronchi, or the bronchioles.

Causes of Lower Airway Obstruction

Common causes of lower airway obstruction are asthma and bronchiolitis.

Signs of Lower Airway Obstruction

The major clinical signs typically occur during the expiratory phase of the respiratory cycle, wherein the child often has wheezing and a prolonged expiratory phase that requires increased expiratory effort. The respiratory rate is usually elevated, particularly in infants. Inspiratory retractions become prominent when the lower airway obstruction impairs inspiration and exhalation, requiring increased respiratory effort. Examples include asthma and bronchiolitis. Other signs of lower airway obstruction include possible decreased air movement on auscultation and cough.

Airway Obstruction and Respiratory Rate

Small airways are obstructed through a variety of mechanisms in acute lower airway obstruction. In asthma, small airways are obstructed via smooth muscle bronchial constriction, mucus plugging, inflammation, and external collapse due to high intrathoracic pressures. In bronchiolitis, small airway obstruction occurs predominantly because of mucus plugging and bronchial edema. If these obstructions are severe, air trapping and lung hyperinflation occur, further exacerbating the obstruction of the smaller airways. An infant with lower airway obstruction, by comparison, typically has a rapid respiratory rate. The infant has a compliant chest wall. If the infant attempts to breathe more deeply, the resulting decrease in intrapleural pressure may result in greater chest wall retractions. When lower airway obstruction is significant, it is more efficient for the infant to breathe at a fast rate with small tidal volumes to maintain minute ventilation, keeping a relatively larger volume of gas in the lungs.

Lung Tissue Disease

This condition describes disease involving the substance (ie, parenchyma or tissue) of the lung. In this state, the child's lungs become stiff because of fluid accumulation in the alveoli, interstitium, or both, requiring increased respiratory effort during inspiration and exhalation. Therefore, retractions and accessory muscle use are common.

Hypoxemia is often marked due to alveolar collapse or reduced oxygen diffusion caused by pulmonary edema fluid and inflammatory debris in alveoli. Tachypnea is common and often quite marked. Patients frequently attempt to counteract alveolar and small airway collapse by increasing efforts to maintain an elevated end-expiratory pressure, often manifested by grunting respirations.

Causes of Lung Tissue Disease

Causes of lung tissue disease vary and can include pneumonia from any cause (eg, bacterial, viral, chemical, aspiration) and cardiogenic (eg, congestive heart failure) and noncardiogenic pulmonary edema (eg, ARDS). Other potential causes include

- Pulmonary contusion (trauma)
- Allergic reaction
- Toxins
- Vasculitis
- Infiltrative disease
- Allergic, vascular, widespread inflammatory, environmental, and other factors

Signs of Lung Tissue Disease

Signs of lung tissue disease include tachypnea (often marked), increased respiratory effort, grunting, crackles (rales) and decreased air movement, diminished breath sounds, tachycardia, and hypoxemia (despite supplemental O₂ administration).

Children with lung tissue disease can often maintain ventilation (ie, CO₂ elimination) with a relatively small number of functional alveoli, but they cannot maintain oxygenation as effectively. As a result, hypoxemia is an earlier sign of lung tissue disease than hypercarbia. Compromised ventilation, indicated by hypercarbia, is typically a late manifestation of the disease process.

Grunting produces early glottic closure during expiration. Grunting is a compensatory mechanism to maintain positive airway pressure and prevent collapse of the alveoli and small airways.

Disordered Control of Breathing

In this state, there is inadequate respiratory effort, and the parent may say that the child is “breathing funny.” There may be periods of increased respiratory rate, effort, or both followed by decreased rate, effort, or both, or the child’s respiratory rate or effort may be continuously inadequate. Often the net effect is hypoventilation leading to hypoxemia and hypercarbia.

Causes of Disordered Control of Breathing

Disordered control of breathing may result from a host of conditions, including neurologic disorders (eg, seizures, central nervous system infections, head injury, brain tumor, hydrocephalus, neuromuscular disease), metabolic abnormalities, and drug overdose. Because disordered control of breathing is typically associated with conditions that impair neurologic function, these children often have a decreased level of consciousness.

Signs of Disordered Control of Breathing

Signs of disordered control of breathing include

- Variable or irregular respiratory rate and pattern (tachypnea alternating with bradypnea)
- Variable respiratory effort
- Shallow breathing with inadequate effort (frequently resulting in hypoxemia and hypercarbia)
- Central apnea (ie, apnea without any respiratory effort)
- Normal or decreased air movement

Summary: Recognizing Respiratory Problems Flowchart

Table 29 summarizes recognition and identification of respiratory problems. Note that this chart does not include all respiratory emergencies but instead provides key characteristics for a limited number of diseases.

Table 29. Recognizing Respiratory Problems Flowchart

PALS: Signs of respiratory problems

Clinical signs		Upper airway obstruction	Lower airway obstruction	Lung tissue disease	Disordered control of breathing
Airway	Patency	Airway open and maintainable/not maintainable			
Breathing	Respiratory rate/effort	Increased			Variable
	Breath sounds	Stridor (typically inspiratory)	Barking cough Hoarseness Wheezing (typically expiratory) Prolonged expiratory phase	Grunting Crackles Decreased breath sounds	Normal
	Air movement	Decreased			Variable
	Circulation	Heart rate	Tachycardia (early); bradycardia (late)		
	Skin	Pallor, cool skin (early); cyanosis (late)			
Disability	Level of consciousness	Anxiety, agitation (early); lethargy, unresponsiveness (late)			
Exposure	Temperature	Variable			
PALS: Identifying respiratory problems by severity					
Progression of respiratory distress to respiratory failure*					
Airway	Respiratory distress: open and maintainable Respiratory failure: not maintainable				
Breathing	Respiratory distress: tachypnea Respiratory failure: bradypnea to apnea				
	Respiratory distress: work of breathing (nasal flaring/retractions) Respiratory failure: increased effort progresses to decreased effort and then to apnea				
	Respiratory distress: good air movement Respiratory failure: poor to absent air movement				
Circulation	Respiratory distress: tachycardia Respiratory failure: bradycardia				
	Respiratory distress: pallor Respiratory failure: cyanosis				
Disability	Respiratory distress: anxiety, agitation Respiratory failure: lethargy to unresponsiveness				
Exposure	Variable temperature				

*Respiratory failure requires immediate intervention.

Part 8: Managing Respiratory Distress and Failure

Respiratory problems are a major cause of cardiac arrest in children. In fact, many infants and children who require CPR (both in and out of hospital) have respiratory problems that progress to cardiopulmonary failure. It may not be possible to differentiate between respiratory distress and respiratory failure on the basis of clinical examination alone because respiratory failure can develop even without significant signs of distress. In children, clinical deterioration in respiratory function may progress rapidly, so there is little time to waste. Promptly recognizing and effectively managing respiratory problems are fundamental to PALS.



Critical Concepts: Intervene Quickly to Restore Respiratory Function

PALS providers must intervene quickly to restore adequate respiratory function. You can greatly improve outcome by identifying early and promptly managing respiratory distress and failure. Once respiratory failure progresses to cardiac arrest, outcome is often poor.

Learning Objective

After completing this Part, you should be able to perform early interventions for respiratory distress and failure.

During the course, you will participate in the Airway Management Skills Station. You will practice and demonstrate your proficiency in performing basic airway management skills, such as inserting airway adjuncts, effective bag-mask ventilation, and suctioning. Refer to the [Appendix](#) for a checklist of required competencies. Refer to the [Resources for Managing Respiratory Emergencies](#) in Part 8 of this manual for details on bag-mask ventilation, respiratory function monitoring devices (eg, monitoring of O₂ saturation by pulse oximetry, monitoring of exhaled CO₂), and O₂ delivery systems.

Rescue Breathing

Respiratory Arrest

Respiratory arrest is the absence of respirations (ie, apnea) with detectable cardiac activity. The provider must provide rescue breathing to prevent cardiac arrest.

Rescue Breathing

Guidelines for rescue breathing for infants and children are as follows:

- Give 20 to 30 breaths per minute (about 1 breath every 2-3 seconds).
- Give each breath in 1 second.
- Each breath should result in visible chest rise.
- Check the pulse about every 2 minutes; if the child becomes pulseless, shout for help and provide compressions as well as ventilation (CPR).
- Use oxygen as soon as it is available.

Initial Management of Respiratory Distress and Failure

The first priority in managing a seriously ill or injured child who is not in cardiac arrest is evaluating the airway and breathing. If you identify signs of respiratory distress or failure, perform initial interventions that support or restore adequate oxygenation and ventilation. Because respiratory conditions are a major cause of cardiac arrest in infants and children, when you detect respiratory distress or failure, it is important to begin appropriate interventions quickly.

Initial interventions include rapidly evaluating respiratory function to identify the type and severity, rather than the precise etiology, of the respiratory problem. Once oxygenation and ventilation are stabilized, identify the cause of the problem to facilitate targeted interventions. Use the evaluate-identify-intervene sequence to monitor symptom progression or response to therapy and to prioritize further interventions.

Initially stabilizing and managing a child in respiratory distress or respiratory failure may include the interventions in [Table 30](#).

Table 30. Initial Management of Respiratory Distress or Failure

Evaluate	Interventions (as indicated)
Airway	<ul style="list-style-type: none"> • Support an open airway (allow child to assume position of comfort) or, if necessary, open the airway with <ul style="list-style-type: none"> – Head tilt–chin lift – Jaw thrust without head tilt if you suspect cervical spine injury. If this maneuver does not open the airway, use the head tilt–chin lift or jaw thrust with gentle head extension • Clear the airway if indicated (eg, suction nose and mouth, remove visualized foreign body). • Consider an oropharyngeal airway or nasopharyngeal airway to improve airway openness/patency.
Breathing	<ul style="list-style-type: none"> • Monitor O₂ saturation by pulse oximetry. • Provide O₂ (humidified if available). Use a high-concentration delivery device such as a nonrebreathing mask for treating severe respiratory distress or possible respiratory failure. • Administer inhaled medication (eg, albuterol, epinephrine) as needed. • Assist ventilation with bag-mask device and supplemental O₂ if needed. • Prepare for inserting an advanced airway if indicated.
Circulation	<ul style="list-style-type: none"> • Monitor heart rate, heart rhythm, and blood pressure. • Establish vascular access (for fluid therapy and medications) as indicated.

Principles of Targeted Management

Once you stabilize oxygenation and ventilation, identify the type of respiratory problem to help prioritize the next interventions. This Part reviews principles of targeted management for the following 4 types of respiratory problems:

- Upper airway obstruction
- Lower airway obstruction
- Lung tissue disease
- Disordered control of breathing

Managing Upper Airway Obstruction

Upper airway obstruction is a mild to severe obstruction of the large airways outside the thorax (ie, in the nose, pharynx, or larynx). Causes of upper airway obstruction are airway swelling, foreign body, or infection. Other causes include edema or swelling of the soft tissue of the upper airway (large tonsils or adenoids), a mass in the airway, thick secretions, congenital narrowing of the upper airway, or poor control of the upper airway due to a decreased level of consciousness.

Infants and small children are especially prone to upper airway obstruction. If the infant has a decreased level of consciousness, the muscles may relax and allow the tongue to fall back and obstruct the oropharynx. Also, if the infant is supine, resting on the large occiput can cause flexion of the neck, resulting in upper airway obstruction. In young infants, nasal obstruction can impair ventilation. Secretions, blood, and debris in the nose, pharynx, and larynx from infection, inflammation, or trauma also can obstruct the airway. Remember that the smaller the airway is, the more easily it can become obstructed.

General Management of Upper Airway Obstruction

General management of upper airway obstruction includes the initial interventions in [Table 30](#). Additional measures focus on relieving the obstruction and may include opening the airway by

- Allowing the child to assume a position of comfort
- Performing manual airway maneuvers, such as a jaw thrust or head tilt–chin lift
- Removing a foreign body
- Suctioning the nose or mouth

- Reducing airway swelling with medications
- Minimizing agitation (agitation often worsens upper airway obstruction)
- Deciding whether an airway adjunct or advanced airway is needed
- Deciding early if a surgical airway (tracheostomy or needle cricothyroidotomy) is needed

You may use suctioning to help remove secretions, blood, or debris; however, if the upper airway obstruction is caused by edema from infection (eg, croup) or allergic reaction, carefully weigh potential benefits vs risks of suctioning. Suctioning may further agitate the child and increase respiratory distress, so consider allowing the child to assume a position of comfort. Give nebulized epinephrine, particularly if the swelling is beyond the tongue. Corticosteroids (inhaled, IV, oral, or intramuscular [IM]) also may help in this situation.

With severe upper airway obstruction, *call early* for advanced help because the provider with the greatest skill and experience in airway management is the most likely person to safely establish an airway. Failure to aggressively treat an acute partial upper airway obstruction may lead to complete airway obstruction and, ultimately, to cardiac arrest.

In less severe cases of upper airway obstruction, infants and children may benefit from specific airway adjuncts. For example, in a child with a decreased level of consciousness, an oropharyngeal airway or nasopharyngeal airway may help to relieve obstruction caused by the tongue. Use an oropharyngeal airway only if the child is deeply unconscious with no gag reflex. A child with a gag reflex may tolerate a nasopharyngeal airway, which should be inserted carefully to avoid nasopharyngeal trauma and bleeding. Avoid using a nasopharyngeal airway in children with increased bleeding risk or severe head or face trauma.

An infant or child with upper airway obstruction from redundant tissues or tissue edema may benefit from noninvasive ventilation with positive airway pressure.

Specific Management of Upper Airway Obstruction by Etiology

Specific causes of upper airway obstruction require specific interventions. This section reviews management of upper airway obstruction due to

- Croup
- Anaphylaxis
- Foreign-body airway obstruction (FBAO)

Manage Croup Based on Severity

Manage croup according to your assessment of clinical severity. The following characteristics of croup are listed by degree of severity:

- **Mild croup:** Occasional barking cough, little or no stridor at rest, absent or mild retractions
- **Moderate croup:** Frequent barking cough, easily audible stridor at rest, retractions at rest, little or no agitation, good air entry by auscultating the peripheral lung fields
- **Severe croup:** Frequent barking cough, prominent inspiratory and occasional expiratory stridor, marked retractions, significant agitation, decreased air entry by auscultating the lungs
- **Severe respiratory distress/impending respiratory failure:** Barking cough (may not be prominent if the child's respiratory effort is growing weaker with the development of severe hypoxemia and hypercarbia), audible stridor at rest (can be difficult to hear with failing respiratory effort), retractions (may not be severe if respiratory effort is failing)
- **Respiratory failure/impending respiratory arrest:** Poor air movement on auscultation, lethargy or decreased level of consciousness, and, sometimes, pallor or cyanosis despite administering supplemental O₂

O₂ saturation may be slightly low in mild and moderate croup and is commonly well below normal in severe croup.

General management for upper airway obstruction includes the initial interventions in [Table 30](#) and may include the specific interventions for managing croup listed in [Table 31](#).

Table 31. Managing Croup

Severity of croup	Interventions
Mild	<ul style="list-style-type: none"> • Consider dexamethasone.
Moderate to severe	<ul style="list-style-type: none"> • Administer humidified O₂. • Give nothing by mouth. • Administer nebulized epinephrine.

	<ul style="list-style-type: none"> Observe for at least 2 hours after giving nebulized epinephrine to ensure continued improvement (no recurrence of stridor). Administer dexamethasone. Consider using heliox (helium-oxygen mixture) for severe disease if the child requires no higher than 40% inspired oxygen concentration.
Respiratory failure or impending respiratory failure	<ul style="list-style-type: none"> Administer a high concentration of O₂; use a nonrebreathing mask if available. Administer dexamethasone IV/IM. Provide assisted ventilation (ie, bag-mask ventilation timed to support the child's own inspiration) for persistent, severe hypoxemia (<90% O₂ saturation) despite O₂ administration, inadequate ventilation, or changes in level of consciousness. Perform endotracheal (ET) intubation if indicated; to avoid injury to the subglottic area, use a smaller ET tube size (a half size smaller than predicted for the child's age). Prepare for surgical airway if needed.



Critical Concepts: Endotracheal Intubation

Endotracheal intubation of the child with upper airway obstruction is a high-risk procedure and should be performed by a team with significant pediatric airway expertise. Use neuromuscular blockade only if you are confident the child's oxygenation and ventilation can be supported with bag-mask ventilation.

Managing Anaphylaxis

In addition to the initial interventions in [Table 30](#), specific interventions that may be used for managing anaphylaxis are listed in [Table 32](#).

Table 32. Managing Anaphylaxis

Severity of allergic reaction	Interventions
Mild	<ul style="list-style-type: none"> Remove the offending agent (eg, stop the IV antibiotic). Get help. Ask the child or caregiver about any history of allergy or anaphylaxis; look for a medical alert bracelet or necklace. Consider an oral dose of antihistamine.
Moderate to severe	<ul style="list-style-type: none"> Administer IM epinephrine by autoinjector or regular syringe every 10 to 15 minutes as needed; repeat doses as needed. Treat bronchospasm (wheezing) with albuterol administered by metered-dose inhaler or nebulizer solution. Give continuous nebulization if indicated (ie, severe bronchospasm). For severe respiratory distress, anticipate further airway swelling and prepare for ET intubation. To treat hypotension: <ul style="list-style-type: none"> Place the child in the supine position as tolerated. Use pharmacological adjuncts once stabilized: <ul style="list-style-type: none"> Methylprednisolone or equivalent corticosteroid IV Diphenhydramine IV Administer isotonic crystalloid (eg, normal saline or lactated Ringer's) 20 mL/kg bolus IV (repeat as needed). For hypotension unresponsive to fluids and IM epinephrine, administer an epinephrine infusion titrated to achieve adequate blood pressure for age.

Managing Foreign-Body Airway Obstruction

If you suspect an FBAO that is mild (the child is able to make sounds and cough forcefully), do not intervene. Call for help and allow the child to try to clear the obstruction by coughing. If you suspect that the FBAO is severe (child makes no sound, is unable to speak or cough, has poor or no air exchange, makes a high-pitched noise while inhaling or no noise at all, has increased respiratory difficulty), perform the maneuvers in [Table 33](#).

Table 33. Managing Foreign-Body Airway Obstruction

Interventions for a responsive infant or child with FBAO	
Infant (younger than 1 year)	Child (1 year to adolescent [puberty])

<ol style="list-style-type: none"> 1. Confirm severe airway obstruction. 2. Give up to 5 back slaps and up to 5 chest thrusts. 3. Repeat Step 2 until the object is expelled or victim becomes unresponsive. 	<ol style="list-style-type: none"> 1. Ask, "Are you choking?" If the child nods or otherwise indicates "yes," say you're going to help. 2. Stand or kneel behind the child. Give abdominal thrusts. 3. Repeat abdominal thrusts until the object is expelled or victim becomes unresponsive.
Child or infant becomes unresponsive	
<ol style="list-style-type: none"> 1. Shout for help. Send someone to activate the emergency response system. 2. Lower the child to the floor. If the child is unresponsive with no breathing or only gasping, begin CPR (no pulse check). 3. Each time you open the airway to deliver breaths, look into the mouth. If you see an object that can be easily removed, remove it. If you do not see an object, continue CPR. <i>Note:</i> Do not perform a <i>blind finger sweep</i> to dislodge a foreign body. This may push the foreign body farther into the airway. It also may cause trauma and bleeding. 4. Continue CPR for 5 cycles or about 2 minutes.* If you are alone, leave the child and activate the emergency response system. Continue CPR until more skilled providers arrive. 	

*Providing effective ventilation by using a bag-mask device on a child with an FBAO may be difficult. Consider using the 2-person bag-mask ventilation technique.

Managing Lower Airway Obstruction

Lower airway obstruction, commonly caused by bronchiolitis and asthma, involves the smaller bronchi and bronchioles inside the thorax.

General Management of Lower Airway Obstruction

General management of lower airway obstruction includes the initial interventions in [Table 30](#).

In a child with severe respiratory distress or respiratory failure, first restore adequate oxygenation; immediately correcting hypercarbia to the child's baseline level is not required because most children can tolerate hypercarbia without adverse effects.

If a child with lower airway obstruction needs bag-mask ventilation, provide effective ventilation at a relatively slow rate, allowing more time for expiration. This reduces the risk that air will remain inside the chest at the end of expiration. Providing too many breaths or breaths with too much volume may result in complications, such as

- Air entering the stomach (gastric distention), resulting in increased risk of vomiting and aspiration and possible prevention of adequate diaphragm movement, limiting effective ventilation
- Risk of pneumothorax (air leak into space surrounding the lungs), resulting in decreased blood return to the heart and risk of lung collapse and resultant complications (severe hypoxemia, obstructive shock)
- Severe air trapping, resulting in severe decrease in oxygenation and decreased venous return to the heart and cardiac output

Specific Management of Lower Airway Obstruction by Etiology

Specific causes of lower airway obstruction require specific interventions. This section reviews management of lower airway obstruction due to the following:

- Bronchiolitis
- Acute asthma

Distinguishing between bronchiolitis and asthma in a wheezing infant can be difficult. However, a history of wheezing episodes suggests that the infant has reversible bronchospasm (ie, asthma). Consider a trial of bronchodilators if the diagnosis is unclear.

Managing Bronchiolitis

In addition to the initial interventions in [Table 30](#), specific measures that may be used for managing bronchiolitis include performing oral or nasal suctioning as needed and considering laboratory and other tests, which may include viral studies, chest x-ray, and arterial blood gas.

Randomized controlled trials of bronchodilator or corticosteroid therapy for bronchiolitis have shown mixed results. Some infants improve when treated with nebulized epinephrine or albuterol, whereas nebulizer therapy aggravates

respiratory symptoms in other infants. Consider a trial of nebulized epinephrine or albuterol treatment, and discontinue it if you observe no improvement. Administer supplemental O₂ if O₂ saturation is less than 94%.

Managing Acute Asthma

Manage asthma according to your assessment of clinical severity ([Table 34](#)). In addition to the initial interventions in [Table 30](#), use specific interventions for managing acute asthma ([Table 35](#)).

Table 34. Classifying Mild, Moderate, and Severe Asthma

Parameter*	Mild	Moderate	Severe	Respiratory arrest imminent
Breathless	Walking Can lie down	Talking (Infant will have softer, shorter cry; difficulty feeding) Prefers sitting	At rest (Infant will stop feeding) Hunched forward	
Talks in	Sentences	Phrases	Words	
Alertness	May be agitated	Usually agitated	Usually agitated	Drowsy or confused
Respiratory rate[‡]	Increased	Increased	Often >30/min	
Accessory muscles and suprasternal retractions	Usually not	Usually	Usually	Paradoxical thoracoabdominal movement
Wheeze	Moderate, often only end-expiration	Loud	Usually loud	Absence of wheeze
Pulse/min[‡]	<100	100-120	>120	Bradycardia
Pulsus paradoxus	Absent <10 mm Hg	May be present 10-25 mm Hg	Often present >25 mm Hg (adult) 25-40 mm Hg (child)	Absence suggests respiratory muscle fatigue
PEF (if used in clinical practice) after initial bronchodilator % predicted or % personal best	>80%	Approximately 60%-80%	<60% predicted or personal best (<100 L/min adults) or response lasts <2 hours	
Paco₂	Normal, test usually not necessary <45 mm Hg	>60 mm Hg <45 mm Hg	<60 mm Hg; possible cyanosis >45 mm Hg; possible respiratory failure	
Spo₂ (room air)%	>95%	91%-95%	<90%	

*The presence of several parameters, but not necessarily all, indicates the general classification of the attack.

[‡]Guide to limits of normal respiratory rate in infants and children: age <2 months, respiratory rate <60/min; age 2-12 months, respiratory rate <50/min; age 1-5 years, respiratory rate <40/min; age 6-8 years, respiratory rate <30/min.

[‡]Guide to limits of normal pulse rate in infants and children: infant (2-12 months), pulse rate <160/min; toddler (1-2 years), pulse rate <120/min; preschool/school age (2-8 years), pulse rate <110/min.

Adapted from National Heart, Lung, and Blood Institute and World Health Organization. *Global Strategy for Asthma Management and Prevention NHLBI/WHO Workshop Report*. US Department of Health and Human Services; 1997. Publication 97-4051.

Table 35. Managing Acute Asthma

Asthma severity	Severity interventions
Mild to moderate	<ul style="list-style-type: none"> Administer humidified O₂ in high concentration via nasal cannula or O₂ mask; titrate according to pulse oximetry. Keep O₂ saturation ≥94%. Administer albuterol by metered-dose inhaler or nebulizer solution. Administer oral corticosteroids.
Moderate to severe	<ul style="list-style-type: none"> Administer humidified O₂ in high concentrations to keep O₂ saturation 95%; use a nonrebreathing mask if needed. If this is unsuccessful, further support such as noninvasive positive-pressure ventilation or high-flow nasal cannula may be indicated. Administer albuterol by metered-dose inhaler (with spacer) or nebulizer solution. If wheezing and aeration are not alleviated, continuous albuterol administration may be required.

	<ul style="list-style-type: none"> • Administer ipratropium bromide by nebulizer solution. Albuterol and ipratropium may be mixed for nebulization. Consider establishing vascular access for administering fluids and medications. • Administer corticosteroids PO/IV. • Consider administering magnesium sulfate by slow (15-30 minutes) IV bolus infusion while monitoring heart rate and blood pressure. • Perform diagnostic assessments (eg, arterial blood gas, chest x-ray) as indicated.
Severe to impending respiratory failure	<p>All of the previous therapies are indicated in addition to the following:</p> <ul style="list-style-type: none"> • Administer O₂ in high concentrations; use a nonrebreathing mask if available. • Administer albuterol by continuous nebulizer. • Administer corticosteroid IV if not already given. • Consider administration of IM epinephrine. • Terbutaline remains an alternative, subcutaneously or by continuous IV infusion. Practically, however, terbutaline can often be difficult to obtain and administer in a timely manner; IM epinephrine is preferred in an emergent situation. • Consider bilevel positive airway pressure (noninvasive positive-pressure ventilation), especially in alert, cooperative children. • Consider ET intubation for children with refractory hypoxemia (low O₂ saturation), worsening clinical condition (eg, decreasing level of consciousness, irregular breathing), or both despite the aggressive medical management described already. Intubation in an asthmatic child carries significant risk for respiratory and circulatory complications. Because of the high intrathoracic pressures in patients with asthma, we highly recommend inserting a cuffed ET tube.

Managing Lung Tissue Disease

Lung tissue disease (also called *parenchymal lung disease*) refers to various clinical conditions. Common causes of lung tissue disease are pneumonia (eg, infectious, chemical, aspiration) and cardiogenic pulmonary edema, but ARDS and traumatic pulmonary contusion are other causes. Lung tissue disease can also result from allergic, vascular, widespread inflammatory, environmental, and other factors.

General Management of Lung Tissue Disease

General management of lung tissue disease includes the initial interventions in [Table 30](#). In children with hypoxemia refractory to high inspired O₂ concentrations, other modalities that offer varying degrees of inspiratory and expiratory pressure support are usually helpful; these include humidified high-flow nasal cannula, CPAP or bilevel positive airway pressure (mechanical ventilation), and positive expiratory pressure (CPAP, noninvasive ventilation, or mechanical ventilation with positive end-expiratory pressure [PEEP]).

Specific Management of Lung Tissue Disease by Etiology

Specific causes of lung tissue disease require specific interventions. This section reviews the management of lung tissue disease from the following causes:

- Infectious pneumonia
- Chemical pneumonitis
- Aspiration pneumonitis
- Cardiogenic pulmonary edema
- Noncardiogenic pulmonary edema (ARDS)

Managing Infectious Pneumonia

Infectious pneumonia results from viral, bacterial, or fungal inflammation of the alveoli. Viruses, bacteria (eg, *Streptococcus pneumoniae*), and atypical bacteria (eg, *Mycoplasma pneumoniae* and *Chlamydia pneumoniae*) commonly cause acute community-acquired pneumonia in children. Methicillin-resistant *Staphylococcus aureus* is increasingly common and may cause empyema (ie, a collection of pus and fluid in the pleural cavity).

In addition to the initial interventions in [Table 30](#), specific interventions for managing acute infectious pneumonia include

- Performing diagnostic assessments (eg, arterial blood gas, chest x-ray, viral studies, complete blood count, blood culture, sputum gram stain and culture) as indicated

- Administering antibiotic therapy (the goal is to administer within the first hour after medical contact); it may not be necessary to draw blood cultures before giving antibiotics, so follow facility protocol
- Treating wheezing with albuterol by metered-dose inhaler or nebulizer solution
- Considering the use of noninvasive positive-pressure ventilation (eg, humidified high-flow nasal cannula, bilevel positive airway pressure, CPAP); in severe cases, endotracheal intubation and mechanical ventilation may be required
- Reducing metabolic demand by normalizing temperature (ie, treating fever)

Managing Chemical Pneumonitis

Chemical pneumonitis is an inflammation of the lung tissue caused by inhaling or aspirating toxic liquids, gases, or particulate matter such as dust or fumes. Aspirating hydrocarbons or inhaling irritant gases (eg, chlorine) can result in noncardiogenic pulmonary edema with increased capillary permeability.

In addition to the initial interventions in [Table 30](#), managing chemical pneumonitis may include specific interventions, such as

- Treating wheezing with nebulized bronchodilator
- Considering using CPAP or noninvasive ventilation. Intubation and mechanical ventilation may be required. Consider early intubation particularly if the child requires transport to a tertiary care facility, is not tolerating secretions, or demonstrates evidence of upper airway edema and obstruction.

Obtaining early consultation in a child with rapidly progressive symptoms. Consider referral to a specialized center for advanced technologies (eg, high-frequency oscillation or pediatric extracorporeal membrane oxygenation).

Managing Aspiration Pneumonitis

Aspiration pneumonitis is a form of chemical pneumonitis resulting from the toxic effects of aspirated oral secretions or stomach acid and enzymes and the subsequent inflammatory response.

In addition to the initial interventions for the general management of lung tissue disease in [Table 30](#), managing aspiration pneumonitis includes specific interventions, such as

- Considering using CPAP or noninvasive ventilation. Intubation and mechanical ventilation may be required in severe cases.
- Considering administering antibiotics if the child has a fever and an infiltrate is present on a chest x-ray. Prophylactic antimicrobial therapy is not indicated.

Managing Cardiogenic Pulmonary Edema

In cardiogenic pulmonary edema, high pressure in the pulmonary capillaries causes fluid to leak into the lung interstitium and alveoli. The most common cause of acute cardiogenic pulmonary edema in children is left ventricular myocardial dysfunction, which can result from congenital heart disease, myocarditis, cardiomyopathy, inflammatory processes, hypoxia, and cardiac-depressant drugs (eg, β -adrenergic blockers, tricyclic antidepressants, calcium channel blockers).

In addition to the initial interventions in [Table 30](#), specific interventions for managing cardiogenic pulmonary edema include

- Providing ventilator support (ie, noninvasive ventilation or mechanical ventilation with PEEP) as needed
- Considering diuretics to reduce left atrial pressure, inotropic infusions, and afterload-reducing agents to improve ventricular function, and obtaining expert consultation
- Reducing metabolic demand by normalizing temperature (treat fever)

Indications for ventilator support (noninvasive ventilation or ET intubation with mechanical ventilation) in children with cardiogenic pulmonary edema include

- Persistent hypoxemia despite oxygen administration and noninvasive ventilation
- Impending respiratory failure
- Hemodynamic compromise (eg, hypotension, severe tachycardia, signs of shock)

Add PEEP during mechanical ventilation to help reduce the need for high O_2 concentrations. You may usually start it at about 5 cm H_2O and adjust upward until O_2 saturation and oxygen delivery improve. Too much PEEP may create pulmonary hyperinflation that impedes both systemic and venous return to the heart, thus reducing cardiac output and O_2 delivery.

Managing Noncardiogenic Pulmonary Edema

ARDS usually follows a pulmonary (eg, pneumonia or aspiration) or systemic (eg, sepsis, pancreatitis, trauma) disease process that injures the interface between the alveoli and pulmonary capillaries and triggers release of inflammatory mediators. As a result, oxygen diffusion into the blood and, to a lesser degree, CO₂ diffusion from the blood to the alveoli is compromised. Recognizing and treating bacteremia, shock, and respiratory failure early may help prevent the progression to ARDS.

Characteristics of ARDS include

- Acute onset (within 7 days after insult)
- PaO₂/FIO₂ 300 or less (with full face-mask bilevel ventilation or CPAP 5 cm H₂O or greater)
- Oxygenation index (OI: [FIO₂ × mean airway pressure × 100]/PaO₂) 4 or greater
- New infiltrate on chest x-ray consistent with acute pulmonary parenchymal disease
- No evidence for a cardiogenic or fluid overload cause of pulmonary edema

In addition to the initial interventions in [Table 30](#), specific interventions for managing ARDS may include

- Monitoring heart rate and rhythm, blood pressure, respiratory rate, pulse oximetry, and end-tidal CO₂
- Obtaining laboratory studies, including arterial blood gas, central venous blood gas, and complete blood count
- Providing ventilatory support (ie, noninvasive ventilation or mechanical ventilation with PEEP) as needed

Indications for ventilatory support (noninvasive ventilation or ET intubation with mechanical ventilation) in children with ARDS are

- Worsening clinical and radiographic lung disease
- Hypoxemia refractory to high concentrations of inspired O₂

Correcting hypoxemia is the most important intervention, accomplished by increasing PEEP until O₂ saturation is adequate. “Permissive” hypercarbia is a treatment approach that recognizes that correcting increased PaCO₂ is less important than correcting hypoxemia. Maintaining low tidal volumes (5-8 mL/kg; lower for children with decreased lung compliance) and keeping peak inspiratory pressure less than 30 to 35 cm H₂O is more important than correcting the PaCO₂.

When you anticipate ET intubation in children with lung tissue disease, anticipate using PEEP and airway pressures as high as 29 to 32 cm H₂O. To ensure that both are provided effectively, use a cuffed ET tube to prevent glottic air leak. When using a cuffed tube, carefully monitor cuff inflation pressure and maintain it according to manufacturer’s recommendations (typically less than 20 to 25 cm H₂O).

Managing Disordered Control of Breathing

Disordered control of breathing results in an abnormal respiratory pattern that produces inadequate minute ventilation. Common causes are neurologic disorders, including increased intracranial pressure (ICP), neuromuscular disease (weakness), central nervous system infections, head injury, brain tumor, and hydrocephalus, as well as conditions that depress the level of consciousness, such as deep sedation, central nervous system infection, seizures, metabolic disorders such as hyperammonemia, poisoning, or drug overdose.

General Management of Disordered Control of Breathing

General management to treat disordered control of breathing includes the initial interventions in [Table 30](#).

Specific Management of Disordered Control of Breathing by Etiology

Specific causes of disordered control of breathing require specific interventions. This section reviews managing disordered control of breathing caused by the following:

- Increased ICP
- Neuromuscular disease
- Poisoning or drug overdose

Managing Respiratory Distress/Failure With Increased ICP

Increased ICP can be a complication of meningitis, encephalitis, intracranial abscess, subarachnoid hemorrhage, subdural or epidural hematoma, traumatic brain injury, hypoxic/ischemic insult, hydrocephalus, and central nervous

system tumor. An irregular respiratory pattern is one sign of increased ICP, and a combination of irregular breathing or apnea, increased mean arterial pressure, and bradycardia is called Cushing's triad. This triad suggests a marked increase in ICP and impending brain herniation. However, children with increased ICP also can present with irregular breathing, hypertension, and tachycardia rather than with bradycardia.

If you suspect increased ICP, obtain neurosurgical consult. In addition to the initial interventions in [Table 30](#), specific interventions for disordered control of breathing due to increased ICP may include

- Ensuring the head is midline, manually stabilizing the cervical spine, and using a jaw-thrust maneuver if you suspect trauma and need to open the airway
- Verifying open/patent airway, adequate oxygenation, and adequate ventilation. You may occasionally use a brief period of mild hyperventilation as temporizing rescue therapy in response to signs of impending brain herniation (eg, irregular respirations or apnea, bradycardia, hypertension, unequal or dilated pupil[s] not responsive to light, decerebrate or decorticate posturing) in the initial 48 hours after injury. If you use hyperventilation, you may consider advanced neuromonitoring for evaluating cerebral ischemia.
- Administering 20 mL/kg IV isotonic crystalloid (normal saline or lactated Ringer's) if the child has poor perfusion or other evidence of poor end-organ function
- Administering pharmacologic therapy for managing increased ICP (eg, osmotic agents, hypertonic saline)
- Treating agitation and pain aggressively once the airway is established and ventilation is adequate
- Avoiding hypotension because it can compromise cerebral blood flow in the setting of elevated ICP.
- Hypertension, as part of Cushing's triad, is often present when ICP is increased. Therapeutic measures in the acute phase should be aimed at reducing the ICP, not administering antihypertensive medications.
- Avoiding and aggressively treating fever

Avoid severe prophylactic hyperventilation (to PaCO₂ less than 30 mm Hg) because hyperventilation can depress cardiac output by impairing venous return. Hyperventilation must be used cautiously—if at all—in the child with traumatic brain injury because excessive hyperventilation may cause cerebral vasoconstriction, leading to brain ischemia and a worse outcome. In patients with severe traumatic brain injury, consider hyperventilation in the initial 48 hours after injury only if there are acute signs of cerebral herniation. It must be guided by neuromonitoring for evaluating cerebral ischemia.

Respiratory Management in Neuromuscular Disease

Chronic progressive neuromuscular diseases can affect the muscles of respiration. Affected children can develop an ineffective cough and difficulty managing secretions. Complications include atelectasis, restrictive lung disease, pneumonia (including aspiration pneumonitis and pneumonia), chronic respiratory insufficiency, and respiratory failure. Consider the initial interventions in [Table 30](#) for disordered control of breathing due to neuromuscular disease. For children with advanced restrictive lung disease, you may use long-term noninvasive ventilation.

Managing Respiratory Distress/Failure in Poisoning or Drug Overdose


One of the most common causes of respiratory distress or failure after a poisoning or drug overdose is depression of central respiratory drive; less common causes are respiratory muscle weakness or paralysis, loss of consciousness, and upper airway obstruction by the tongue.

Complications of disordered breathing in this setting include upper airway obstruction, poor respiratory effort and rate, hypoxemia, aspiration, and respiratory failure. Complications from a decreased level of consciousness (eg, aspiration pneumonitis and noncardiogenic pulmonary edema) may also result in respiratory failure. If you suspect poisoning, contact your local poison control center.

Supporting the airway and ventilation is the main therapeutic intervention for managing respiratory distress or failure caused by poisoning or drug overdose. In addition to the initial interventions in [Table 30](#), specific interventions for disordered control of breathing due to poisoning or drug overdose may include

- Contacting a poison control center (1-800-222-1222)
- Suctioning the airway in case of vomiting
- Administering an antidote as indicated. Administer naloxone for opioid overdose (IM and intranasal preparations available) in addition to standard BLS support for patients with known or suspected opioid overdose who have respiratory arrest but with a definite pulse.
- Performing diagnostic assessments as indicated (eg, arterial blood gas, ECG, chest x-ray, electrolytes, glucose, serum osmolality, drug screen)
- Potentially preparing for transfer when in a rural or nonchildren's hospital

Figure 27 is a step-by-step guide to providing care for a victim with a suspected opioid-associated emergency.



Suspected opioid poisoning (Step 1)

- Check for responsiveness.
- Shout for nearby help.
- Activate the emergency response system.
- If you are alone, get naloxone and an AED if available. If someone else is present, send that person to get them.

Is the person breathing normally? (Step 2)

- If Yes (the person is breathing normally), go to Steps 3 and 4.
- If No (the person is not breathing normally), go to Step 5.

Prevent deterioration (Step 3)

- Tap and shout. Check for responsiveness by tapping the child's shoulders. Shout, "Are you OK?"
- Open and reposition the airway if needed to maintain normal breathing. This may be necessary if the victim is not responsive or is responsive but unable to maintain an open airway due to a depressed level of consciousness.
- Consider administering naloxone, if available. If you suspect an opioid overdose, it is reasonable to give naloxone according to package directions and per local protocol. Monitor for response.
- Transport to the hospital. If the victim is not already in a healthcare setting, they should be transported by EMS to a hospital.

Ongoing assessment of responsiveness and breathing (Step 4)

Continue to assess responsiveness and breathing until the child is transferred to advanced care. Children with opioid-associated emergencies may not be able to maintain an open airway or breathe normally. Even those who receive naloxone may develop respiratory problems that can lead to cardiac arrest.

Does the child have a pulse? (Step 5)

Assess for a pulse for no more than 10 seconds.

- If yes (a pulse is present), go to Step 6.
- If no (a pulse is not present), go to Step 7.

Support ventilation (Step 6)

- Open and reposition the airway before giving rescue breaths.
- Provide rescue breathing or bag-mask ventilation. This can help prevent cardiac arrest. Continue until spontaneous, normal breathing occurs. Reassess the victim's breathing and pulse every 2 minutes. If there is no pulse, provide CPR (see Step 7).
- Give naloxone according to package directions and per local protocol.

Start CPR (Step 7)

- If the child is not breathing normally and no pulse is present, provide high-quality CPR including ventilation. Use the AED as soon as it is available.
- Consider naloxone. If naloxone is available and you suspect an opioid overdose, it is reasonable to give it according to package directions and per local protocol. High-quality CPR should take priority over giving naloxone.

- Refer to Pediatric Cardiac Arrest Algorithm.



Critical Concepts: Medications to Avoid in Children With Neuromuscular Disease

Recall that the use of succinylcholine for intubating children with neuromuscular diseases may trigger life-threatening conditions, such as hyperkalemia or malignant hyperthermia. Several commonly used pediatric acute care drugs, such as aminoglycosides, have intrinsic neuromuscular blocking activity that can worsen respiratory muscle weakness.

Summary: Managing Respiratory Emergencies Flowchart

The Managing Respiratory Emergencies Flowchart summarizes general management of respiratory emergencies and specific management by etiology (Table 36). This chart does not include all respiratory emergencies but provides key management strategies for a limited number of diseases.

Table 36. Managing Respiratory Emergencies Flowchart

Managing respiratory emergencies flowchart		
<ul style="list-style-type: none">• Airway positioning• Suction as needed	<ul style="list-style-type: none">• Oxygen• Pulse oximetry	<ul style="list-style-type: none">• ECG monitor as indicated• BLS as indicated
Upper airway obstruction		
Specific management for selected conditions		
Croup	Anaphylaxis	Aspiration foreign body
<ul style="list-style-type: none">• Nebulized epinephrine• Corticosteroids	<ul style="list-style-type: none">• IM epinephrine (or autoinjector)• Albuterol• Antihistamines• Corticosteroids	<ul style="list-style-type: none">• Allow position of comfort• Specialty consultation
Lower airway obstruction		
Specific management for selected conditions		
Bronchiolitis	Asthma	
<ul style="list-style-type: none">• Nasal suctioning• Consider bronchodilator trial	<ul style="list-style-type: none">• Albuterol ± ipratropium• Corticosteroids• Magnesium sulfate• IM epinephrine (if severe)• Terbutaline	
Lung tissue disease		
Specific management for selected conditions		
Pneumonia/pneumonitis	Pulmonary edema	
Infectious, chemical, aspiration	Cardiogenic or noncardiogenic (ARDS)	
<ul style="list-style-type: none">• Albuterol• Antibiotics (as indicated)• Consider noninvasive or invasive ventilatory support with PEEP	<ul style="list-style-type: none">• Consider noninvasive or invasive ventilatory support with PEEP• Consider vasoactive support• Consider diuretic	
Disordered control of breathing		
Specific management for selected conditions		
Increased ICP	Poisoning/overdose	Neuromuscular disease
<ul style="list-style-type: none">• Avoid hypoxemia• Avoid hypercarbia• Avoid hyperthermia• Avoid hypotension	<ul style="list-style-type: none">• Antidote (if available)• Contact poison control	<ul style="list-style-type: none">• Consider noninvasive or invasive ventilatory support

Resources for Managing Respiratory Emergencies

Bag-Mask Ventilation

Bag-mask ventilation can adequately oxygenate and ventilate for a child with no breathing or inadequate breathing despite an open/patent airway. Signs of inadequate breathing are apnea, abnormal respiratory rate, inadequate breath sounds, and hypoxemia despite supplemental O₂. When properly performed, bag-mask ventilation is as effective as ventilation through an endotracheal (ET) tube for short periods and may be safer. In the out-of-hospital setting, bag-mask ventilation is especially useful if the transport time is short or providers are inexperienced in inserting advanced airways or have insufficient opportunities to maintain competence in this skill.

Preparing for the Course

All healthcare providers who care for infants and children should be able to ventilate effectively with a bag and mask. In this course you will be required to demonstrate effective bag-mask ventilation during BLS testing, in the Airway Management Skills Station, and in some case simulations.

How to Select and Prepare the Equipment

For ventilation to be effective with a bag-mask device, you must know how to select the face mask, prepare the ventilation bag (such as the self-inflating or flow-inflating bag), and provide supplemental O₂ if needed.

Face Mask

Select a transparent face mask, if available, that extends from the bridge of the child's nose to the cleft of the chin, covering the nose and mouth but not compressing the eyes ([Figure 28](#)). A transparent mask allows you to see the color of the child's lips and condensation on the mask (which indicates exhalation) as well as regurgitation. The mask should have a soft rim (eg, flexible cuff) that molds easily to create a tight seal against the face. If the face-mask seal is not tight, O₂ intended for ventilation will escape under the mask and ventilation will not be effective.




Figure 28. Proper area of the face for face-mask application. Note that no pressure is applied to the eyes.

Self-Inflating Bag

A self-inflating bag, typically used for initial resuscitation, consists of a bag with an intake valve and a nonrebreathing outlet valve. The intake valve allows the bag to fill with either O₂ or room air. When you compress the bag, the intake valve closes and the nonrebreathing outlet valve opens, allowing either room air or an air-O₂ gas mixture to flow to the child. When the child exhales, the nonrebreathing outlet valve closes to prevent the child from rebreathing CO₂ and exhaled gases are vented.

[Figure 29](#) demonstrates how these bags work. [Figures 29A](#) and [29B](#) show bags with O₂ reservoirs. When you release these bags, O₂ flows into the bag from the O₂ source and from the reservoir, so the concentration of O₂ in the bag remains 100%, and O₂ continuously flows into the reservoir. [Figures 29C](#) and [29D](#) show bags without O₂ reservoirs. When you release these bags, O₂ flows into the bag from the O₂ source, but ambient air is also entrained into the bag, so the bag becomes filled with a mixture of O₂ and ambient air. With both setups, exhaled patient air flows into the atmosphere near the mask and bag connection (see the gray arrows coming from the mask in [Figures 29A](#) and [29C](#)).






Figure 29A. Self-inflating ventilation bag with face mask, with (A and B) and without (C and D) O₂ reservoir. A, Re-expansion of the bag with O₂ reservoir.




Figure 29B. Compressing the bag with O₂ reservoir delivers 100% O₂ to the patient (purple arrow).




Figure 29C. Re-expansion of the bag without an O₂ reservoir.




Figure 29D. Compressing the bag without O₂ reservoir delivers O₂ mixed with room air (aqua arrow).

The amount of delivered O₂ is affected by tidal volume and peak inspiratory flow rate. Even with supplemental O₂ attached, the delivered O₂ concentration varies from 30% to 80%. To deliver a high O₂ concentration (60% to 95%), attach an O₂ reservoir to the intake valve. Maintain an O₂ flow of 10 to 15 L/min into a reservoir attached to a pediatric bag and a flow of at least 15 L/min into an adult bag.



Critical Concepts: Use O₂ During Resuscitation

- Attach an O₂ reservoir to the self-inflating bag as soon as possible during a resuscitation attempt. Frequently verify that O₂ is attached and flowing to the bag. Remember to listen for O₂ flow and check O₂ tank pressure or verify connection to a wall O₂ source.
- Once circulation is adequate and when appropriate equipment is available, titrate O₂ administration to maintain an O₂ saturation of 94% to 99%.

Check whether the bag has a pop-off valve. Many self-inflating bags have a pressure-limited pop-off valve set at 35 to 45 cm H₂O to prevent excessive airway pressures from developing. However, if the child has poor lung compliance, has high airway resistance, or needs CPR, an automatic pop-off valve may prevent delivery of sufficient tidal volume, resulting in inadequate ventilation and chest expansion. Ventilation bags used during CPR should have *no pop-off valve* or the valve should be twisted into the closed position.



Critical Concepts: Continuous O₂ Flow Not Possible With Some Self-Inflating Bags

Self-inflating bag-mask devices with a fish-mouth or leaf-flap–operated nonrebreathing outlet valve do not provide a continuous flow of O₂ to the mask. Such valves open only if the bag is squeezed or the mask is sealed tightly to the face and the child generates significant inspiratory force to open the valve. Many infants cannot generate the inspiratory pressure required to open the outlet valve. Do not use this type of bag to provide supplemental O₂ to a spontaneously breathing infant or child.

Flow-Inflating Bag

You may use the second kind of basic ventilation bag, a flow-inflating bag ([Figure 30](#)) (also called an *anesthesia bag*), in the intensive care unit, delivery room, and operating room. A flow-inflating bag may require more operator experience to provide safe and effective ventilation than is needed with a self-inflating bag. To ventilate effectively with a flow-inflating bag, the provider must be able to adjust the flow of O₂, adjust the outlet control valve, ensure a proper seal with the face mask, and deliver the appropriate tidal volume at the correct rate. For these reasons, only trained and expert providers should use flow-inflating bags.



Figure 30A. Flow-inflating bag. A, With a pressure manometer.



Figure 30B. Without a pressure manometer.

Bag Size

Use a self-inflating bag with a volume of at least 450 to 500 mL or larger for infants and young children because smaller bags may not deliver an effective tidal volume over the longer inspiratory times required by full-term neonates and infants. In older children or adolescents, you may need to use an adult self-inflating bag (1000 mL or larger) to achieve chest rise and minute ventilation.

Providing PEEP During Bag-Mask Ventilation

Using PEEP may improve oxygenation in children with lung tissue disease or low lung volumes. Provide PEEP during ventilation with a bag by adding a compatible spring-loaded ball or disk or a magnetic-disc PEEP valve to the bag-mask or bag-tube system. Do not use self-inflating bag-mask devices equipped with PEEP valves to provide CPAP

during spontaneous breathing because the outlet valve in the bag will not open (and provide gas flow) unless the child generates significant negative inspiratory pressure.

How to Test the Bag-Mask Device

To ensure proper function of any bag and mask system, test all components before using by

- Checking the bag for leaks by occluding the patient outlet valve with your hand and squeezing the bag
- Checking the gas flow control valves (including any CPAP valve) to verify proper function
- Checking the pop-off valve (if present) to ensure that it can be closed
- Checking that the O₂ tubing is securely connected to the device and the O₂ source
- Listening for the sound of O₂ flowing into the bag
- Ensuring that the cuff of the mask (if present) is adequately inflated

How to Position the Child

Properly position the child to maintain an open airway. During bag-mask ventilation, you may need to move the child's head and neck gently through a range of positions to optimize ventilation. A "sniffing" position without hyperextending the neck is usually best for infants and toddlers.

To achieve a "sniffing" position, place the child supine and flex the neck forward at the shoulder level while extending the head. Position the opening of the external ear canal at the level of or in front of the anterior aspect of the shoulder while the head is extended. Avoid hyperextending the neck because this may obstruct the airway.

Children older than 2 years may require padding under the occiput, while younger children and infants may need padding under the shoulders or upper torso to prevent excessive flexion of the neck that can occur when the prominent occiput rests on a flat surface.

Figure 31 shows these positions. In Figure 31A, the child is on a flat surface (eg, bed or table), and the oral (O), pharyngeal (P), and tracheal (T) axes pass through 3 divergent planes. In Figure 31B, a folded sheet or towel is placed under the occiput aligns the pharyngeal and tracheal axes. In Figure 31C, extension of the atlanto-occipital joint results in the alignment of the oral, pharyngeal, and tracheal axes when the head is extended and the chin is lifted. Note that the external ear canal is anterior to the shoulder. Figure 31D shows the incorrect position with neck flexion, and Figure 31E shows the correct position for ventilation and ET intubation for an infant. Note that the external ear canal is anterior to the shoulder.




Figure 31A. Correct positioning of the child older than 2 years for ventilation and ET intubation. A, Child on a flat surface (eg, bed or table).




Figure 31B. A folded sheet or towel under the occiput.






Figure 31C. Extension of the atlanto-occipital joint.



Figure 31D. Incorrect position with neck flexion.



Figure 31E. Correct position for ventilation and ET intubation for an infant.

Modified from Coté CJ, Todres ID. The pediatric airway. In: Coté CJ, Ryan JF, Todres ID, Goudsouzian NG, eds. *A Practice of Anesthesia for Infants and Children*. 2nd ed. WB Saunders Co; 1993:55-83, copyright Elsevier.

How to Perform Bag-Mask Ventilation

One or two providers can operate a bag-mask device while another rescuer provides compressions, but because effective bag-mask ventilation requires complex steps, it is not recommended for a lone rescuer to perform during CPR. Instead, the lone rescuer should use the mouth-to-barrier device technique for ventilation during CPR. Bag-mask ventilation can be provided effectively during 2-rescuer CPR.

1-Person Bag-Mask Ventilation Technique

If 1 healthcare provider is performing bag-mask ventilation, the provider must open the airway and keep the mask sealed to the child's face with one hand ([Figure 32](#)) and squeeze the bag with the other hand. Form a tight seal between the mask and the child's face and open the airway by using the E-C clamp technique, described in the following steps.

1. If you do not suspect cervical spine injury, tilt the head back and lift the jaw against the mask, pressing and sealing the mask on the face. This technique moves the tongue away from the posterior pharynx, moves the jaw forward, and opens the mouth. If possible, the mouth should be open under the mask as a result of either lifting the jaw or inserting an oropharyngeal airway.
2. With the other hand, squeeze the ventilation bag until the chest rises. Deliver each breath over 1 second. Avoid excessive ventilation (refer to [How to Deliver Effective Ventilation](#) in this Part).



Figure 32. One-handed E-C clamp face-mask application technique. Three fingers of

one hand lift the jaw (forming the “E”) while the thumb and index finger hold the mask to the face (making a “C”).



Critical Concepts: E-C Clamp Technique

The technique of opening the airway and making a seal between the mask and the face is called the E-C clamp technique. The third, fourth, and fifth fingers of one hand (forming an “E”) are positioned along the jaw to lift it forward; then the thumb and index finger of the same hand (forming a “C”) make a seal to hold the mask to the face. Avoid pressing on the soft tissues underneath the chin (the submental area) because this can push the tongue into the posterior pharynx, resulting in airway compression and obstruction.

2-Person Bag-Mask Ventilation Technique

If 2 healthcare providers are available to perform bag-mask ventilation, one provider uses both hands to open the airway and keep the mask sealed to the child’s face and the other provider squeezes the bag ([Figure 33](#)). Both providers should ensure that chest rise is visible. Be careful to avoid delivering too large a tidal volume, which may result in excessive ventilation.



Figure 33. Two-person bag-mask ventilation technique.

The 2-person technique may provide more effective bag-mask ventilation than a 1-person technique. Also, 2-person bag-mask ventilation may be necessary when

- Making a seal between the face and the mask is difficult
- The provider’s hands are too small to reach from the front of the mask to behind the jaw or to open the airway and create a seal between the face and the mask
- Significant airway resistance (ie, asthma) or poor lung compliance (ie, pneumonia or pulmonary edema) occur
- Restricting spinal motion is necessary

How to Deliver Effective Ventilation

Avoid excessive ventilation; use only the force and tidal volume necessary to just make the chest rise.



Critical Concepts: Effective Ventilation With a Bag-Mask Device

Give each breath slowly over about 1 second. Watch for chest rise. If the chest does not rise, reopen the airway. Verify that there is a tight seal between the mask and the face. Reattempt ventilation.

Precaution

Healthcare providers often deliver excessive ventilation during CPR, which is harmful because it

- Increases intrathoracic pressure and impedes venous return, thus decreasing filling of the heart between compressions, reducing blood flow generated by the next compression, and reducing coronary perfusion and cerebral blood flow
- Causes air trapping and barotrauma in children with small airway obstruction
- Increases the risk of regurgitation and aspiration in children without an advanced airway

Clinical Parameters of Oxygenation and Ventilation

Frequently monitor the following parameters to assess the effectiveness of oxygenation and ventilation:

- Visible chest rise with each breath
- O₂ saturation
- Exhaled CO₂
- Heart rate
- Blood pressure
- Distal air entry
- Signs of improvement or deterioration (eg, appearance, color, agitation)

Troubleshooting Ineffective Ventilation

If you cannot achieve effective ventilation (ie, the chest does not rise), do the following:

- Reposition/reopen the airway: attempt to further lift the jaw and ensure that the child is placed in a sniffing position
- Verify mask size and ensure a tight face-mask seal
- Suction the airway if needed
- Check the O₂ source
- Check the ventilation bag and mask
- Treat gastric inflation
- Consider 2-person bag-mask ventilation and inserting an oropharyngeal airway (OPA)

Spontaneously Breathing Child

In a spontaneously breathing child who requires bag-mask ventilation, it may be difficult to coordinate the bagged breaths with the child's inspirations when using a self-inflating bag. Not only may breaths provided with a self-inflating bag be ineffective, but poorly timed breaths may stimulate coughing, vomiting, laryngospasm, and gastric inflation, which prevent effective ventilation. In these situations, a flow-inflating bag (eg, anesthesia bag) may be more beneficial; these bags provide CPAP and flows that can be manipulated to the needs of the child. If necessary, the healthcare provider can also augment the child's spontaneous inspirations by squeezing the self-inflating bag.

Detecting for Changes in Lung Compliance

When performing bag-mask ventilation, be aware of the child's lung compliance. A poorly compliant lung is "stiff" or difficult to inflate. A sudden increase in lung stiffness during ventilation with a bag may indicate airway obstruction, decreased lung compliance, or development of a pneumothorax. Lung distention from excessive inflating pressures, PEEP, or rapid assisted respiratory rates with short exhalation time may also cause the feel of "stiff lungs" during ventilation.

Causes and Prevention of Gastric Inflation

Stomach inflation or distention frequently develops during bag-mask ventilation. Gastric inflation is more likely to develop during assisted ventilation if

- A partial airway obstruction is present
- High airway pressures are needed, such as in a child with poor lung compliance
- The bag-mask ventilation rate is too fast
- The tidal volume delivered is excessive
- The peak inspiratory pressure created is excessive (eg, more than 30 cm H₂O)
- The child is unconscious or is in cardiac arrest (because the gastro-esophageal sphincter opens at a lower than normal pressure)

Gastric inflation can impair a child's ventilation by limiting lung volumes, interfering with effective ventilation, and causing regurgitation. To minimize gastric inflation

- Ventilate at a rate of 1 breath every 2 to 3 seconds (about 20-30 breaths per minute)
- Use a manometer (if available) and avoiding, if at all possible, generating excessive peak inspiratory pressures (eg, more than 30 cm H₂O) by delivering each breath over about 1 second
- Deliver enough volume and pressure to produce visible chest rise
- Consider administering cricoid pressure; although cricoid pressure is not routinely recommended during endotracheal intubation of pediatric patients, it can be considered to reduce the risk of gastric inflation

- If cricoid pressure is applied, it must be discontinued if
 - A tracheal obstruction compromises bag-mask ventilation or
 - The speed or ease of intubation is compromised

Advanced providers may perform gastric decompression by inserting a nasogastric or orogastric tube.

Suctioning

Suction Devices

Suction devices can be either portable or wall-mounted units.

- Although portable suction devices are easy to transport, they may not have adequate suction power. A suction force of –80 to –120 mm Hg is generally needed to remove airway secretions.
- Bulb or syringe suction devices are simple to use and require no outside vacuum source. These devices may be inadequate in larger patients or when secretions are thick or copious.
- Wall-mounted suction units can provide a high suction force of more than –300 mm Hg.

Suction devices used in children should have adjustable suction regulators so that you can use sufficient suction force and minimize tissue trauma. Large-bore, noncollapsible suction tubing should always be joined to the suction unit. Semirigid pharyngeal tips (tonsil suction tips) and appropriate sizes of catheters should be available.

Indications

You may need to suction secretions, blood, or vomit from the oropharynx, nasopharynx, or trachea to achieve or maintain an open/patent airway.

Complications

Complications of suctioning can include

- Hypoxia
- Vagal stimulation resulting in bradycardia
- Gagging and vomiting
- Soft tissue injury
- Agitation

Soft vs Rigid Catheters

Both soft, flexible and rigid suctioning catheters are available. Use a soft, flexible suction catheter for aspirating thin secretions from the oropharynx and nasopharynx and for suctioning an advanced airway (eg, ET tube). Use a rigid wide-bore suction cannula (“tonsil tip”) for suctioning the oropharynx, particularly if thick secretions, vomit, or blood are present.

Catheter Sizes

For a guide to selecting the appropriate-sized suction catheter, use a color-coded length-based resuscitation tape or other reference.

Oropharyngeal Suctioning Procedure

Follow these steps to suction the oropharynx:

1. Gently insert the distal end of the suction catheter or device into the oropharynx over the tongue. Guide it into the posterior pharynx (back of the throat).
2. Apply suction by covering the catheter side opening. At the same time, withdraw the catheter with a rotating or twisting motion.
3. Try to limit suction attempts to 10 seconds or less to help reduce the risk of hypoxemia (low oxygen saturation). You may give short periods of 100% oxygen immediately before and after each suctioning attempt.

Note: Suction attempts may need to be longer than 10 seconds if the airway is obstructed (eg, by blood). You cannot adequately oxygenate or ventilate unless the airway is open and clear.



Critical Concepts: Monitoring During Suctioning

Monitor the child's heart rate, oxygen saturation, and clinical appearance during suctioning. In general, if bradycardia develops or clinical appearance deteriorates, interrupt suctioning. Give high-flow oxygen and bag-mask ventilation, if needed, until the heart rate and clinical appearance return to normal.

Oropharyngeal Airway

Description

Usually made of plastic, the OPA consists of a flange, a short bite-block segment, and a curved body. It is shaped to provide an air channel and a passage for a suction catheter to the pharynx. The OPA fits over the tongue to prevent it and other soft structures of the throat from obstructing the airway. It is not intended to be used long-term as a bite-block device in agitated patients.

Indications

An OPA may relieve upper airway obstruction caused by the tongue. Using a correct-sized OPA will not damage laryngeal structures. It may be used in the *unconscious* child with no gag reflex if procedures to open the airway (eg, head tilt–chin lift or jaw thrust) fail to provide and maintain a clear, unobstructed airway; however, an OPA should not be used in a *conscious* or *semiconscious* child because it may stimulate gagging and vomiting. Before using an OPA, check if the child has a gag reflex. If so, do not use an OPA.

Complications

Choose the correct-sized OPA. A properly sized OPA relieves airway obstruction caused by the tongue without damaging the larynx. The tip of the OPA should end just at the angle of the jaw, so that once inserted, it will align with the glottis opening ([Figure 34A](#)). If the OPA is *too large*, it can block the airway or cause trauma to the laryngeal structures ([Figure 32B](#)). If the OPA is *too small* or is inserted improperly, it can push the tongue into the back of the throat and contribute to airway obstruction ([Figure 34C](#)).



Figure 34A. Selecting an OPA. A, Properly sized OPA.




Figure 34B. If the OPA is too large, it will obstruct the airway by pushing the epiglottis down.




Figure 34C. If the OPA is too small, it will worsen airway

obstruction by pushing the tongue into the back of the throat.

Modified from Coté CJ, Todres ID. The pediatric airway. In: Coté CJ, Ryan JF, Todres ID, Goudsouzian NG, eds. *A Practice of Anesthesia for Infants and Children*. 2nd ed. WB Saunders Co; 1993:55-83, copyright Elsevier.

Airway Selection and Insertion Procedure

OPA sizes range from 4 to 10 cm in length (Guedel sizes 000-4). Follow these steps to choose the correct-sized OPA and insert it into the airway:

1. Place the OPA against the side of the child's face. The tip of the OPA should extend from the corner of the mouth to the angle of the jaw ([Figure 34A](#)).
2. Gently insert the OPA directly into the oropharynx. The use of a tongue blade to depress the tongue may be helpful.
3. After inserting an OPA, monitor the child. Keep the head and jaw positioned properly to maintain an open/patent airway. Suction the airway as needed.

Oxygen Delivery Systems

Indications for Oxygen

For children with respiratory distress or shock, oxygen uptake by the lungs and oxygen delivery to the tissues typically reduce. At the same time, tissue demand for oxygen may increase. Give high-flow oxygen to all seriously ill or injured children with severe respiratory distress, shock, or changes in mental status. As soon as possible, add humidification to the oxygen delivery system to help prevent airway dryness.

Giving Oxygen to a Conscious Child

Because agitation can increase oxygen demand and respiratory distress, balance the need to improve oxygen delivery against possible agitation that may result from applying an oxygen delivery device to an alert child in respiratory distress. If a child is agitated by one method of oxygen delivery, try an alternative technique. For example, if the child is upset by an oxygen mask, try directing a "blow-by" stream of humidified oxygen toward the child's mouth and nose. It may be helpful to have a person familiar to the child, such as a parent, introduce the oxygen delivery equipment.

When giving oxygen to an alert child in respiratory distress, allow the child to remain in a position of comfort to minimize respiratory effort and help keep the airway as open as possible. For infants and young children, the best position might be in the arms of the parent or caregiver.

Giving Oxygen to a Child With a Decreased Level of Consciousness

If a child has a decreased level of consciousness, the airway may become obstructed by a combination of the following:

- Flexing the neck
- Relaxing the jaw
- Displacing the tongue against the back of the throat
- Accumulating secretions

If the child is unconscious with no cough or gag reflex, open the airway and insert an OPA. Use the head tilt–chin lift maneuver or a jaw thrust to open the airway.

If you don't suspect trauma and the child is breathing normally, roll the child onto her side in a neutral position. Place the child on her side only if you do not need to perform any other interventions.

Suction the oropharynx and nasopharynx to clear secretions, mucus, or blood if needed. Once the airway is open and clear, you can give oxygen by a variety of oxygen delivery systems.



Critical Concepts: Jaw Thrust

A properly performed jaw thrust is the most effective means to open the pediatric airway. Providers, however, are often inexperienced with this technique because it is difficult to practice the skill with some CPR manikins.

Types of Oxygen Delivery Systems

For a spontaneously breathing child who needs supplemental oxygen, you need to know which oxygen delivery system to use—either low flow or high flow. Consider the child's clinical status and the desired concentration of inspired oxygen when choosing the appropriate system. Low-flow oxygen delivery systems include a nasal cannula and simple oxygen mask, while high-flow oxygen delivery systems include a nonrebreathing mask with reservoir and a high-flow nasal cannula.

Several factors determine the concentration of inspired oxygen. These include oxygen flow into the device, the child's inspiratory flow, and how tightly the device fits against the child's face.

Low-Flow Oxygen Delivery Systems

A low-flow oxygen delivery system delivers air through a nasal cannula or a simple mask that does not fit tightly against the child's face. The oxygen flow into the delivery device is less than the child's inspiratory flow rate. When the child inhales, the child inspires some room air along with the oxygen provided by the device. As a result, the oxygen from the device mixes with room air, so the child receives a variable oxygen concentration. The higher the oxygen flow provided, the higher the inspired oxygen concentration.

Low-flow systems generally provide an inspired oxygen concentration of about 22% to 60% and are used for relatively stable children who require a relatively low inspired oxygen concentration, such as when a child is not in severe respiratory distress or shock.

The nasal cannula and simple oxygen mask are examples of low-flow oxygen delivery systems.

Nasal Cannula

The nasal cannula is typically a low-flow oxygen delivery device that delivers an inspired oxygen concentration of 22% to 60%. The appropriate oxygen flow rate for the nasal cannula is 0.25 to 4 L/min.

The nasal cannula is suitable for infants and children who require only low concentrations of supplemental oxygen. Note that in small infants, a nasal cannula may deliver a high inspired oxygen concentration (refer, also, to [High-Flow Nasal Cannula](#) later in this Part). The oxygen flow rate alone cannot reliably determine the inspired oxygen concentration delivered via nasal cannula. Other factors have influence, such as

- The child's size
- Inspiratory flow rate
- Volume of inspired air
- Nasopharyngeal and oropharyngeal volume
- Nasal resistance (eg, oxygen delivery is compromised if nares are obstructed)
- Oropharyngeal resistance

Simple Oxygen Mask

The simple oxygen mask is a low-flow device that delivers an inspired oxygen concentration of 35% to 60%. The appropriate flow rate for the simple oxygen mask is 6 to 10 L/min.

The simple oxygen mask cannot deliver an inspired oxygen concentration greater than 60% because room air enters between the mask and the face and through ports in the side of the mask during inspiration. The oxygen concentration delivered to the child is reduced if

- The child's inspiratory flow is high
- The mask does not fit tightly against the face
- The oxygen flow into the mask is low

A minimum oxygen flow rate of 6 L/min is needed to maintain an increased inspired oxygen concentration and prevent rebreathing of exhaled CO₂.

Several types of oxygen masks can deliver humidified oxygen in a wide range of concentrations. The soft vinyl pediatric mask may agitate and upset infants and toddlers, thus increasing oxygen demand and potentially resulting in increased respiratory distress. This mask may be used effectively in older children.

High-Flow Oxygen Delivery Systems

High-flow oxygen systems reliably deliver an oxygen concentration of greater than 60%. In a high-flow oxygen delivery system, the oxygen flow rate is high, at least 10 L/min. A nonrebreathing mask is the most common example of a high-flow system; high-flow nasal cannula is another example.

High-flow systems should be used in emergency settings whenever the child has respiratory distress or shock.

Nonrebreathing Mask

The nonrebreathing mask ([Figure 35](#)) is a high-flow delivery device. An inspired oxygen concentration of 95% can be achieved with an oxygen flow rate of 10 to 15 L/min and the use of a well-sealed face mask.




Figure 35. Nonrebreathing mask with reservoir.

A nonrebreathing mask consists of a face mask and reservoir bag with the addition of 2 one-way valves:

- A valve in 1 or both exhalation port(s) to prevent room air from entering the mask during inspiration
- A valve placed between the reservoir bag and the mask to prevent the flow of exhaled gas into the reservoir

Adjust the oxygen flow rate into the mask to prevent bag collapse (usually greater than 10 L/min). The bag is filled with oxygen to meet the child's total maximum inspired flow requirements. During inspiration, the child draws 100% oxygen from the reservoir bag and the oxygen inflow. Room air does not enter the mask if the mask is tight-fitting and the delivery system is closed.

High-Flow Nasal Cannula

The nasal cannula can also be used as a high-flow delivery device by adjusting the oxygen flow rate from 4 L in infants to up to 40 L or more in adolescents. The flow can be titrated to provide additional inspiratory and expiratory pressure to improve the patient's work of breathing.

High-flow nasal cannula systems deliver a combination of both room air and oxygen. With these systems, the oxygen concentration is titrated based on the patient's needs and oxyhemoglobin saturations.

Nebulizer

Components

A nebulizer has the following components:

- Nebulizer reservoir
- Nebulizer cap
- T-piece
- Spacer
- Handheld mouthpiece or face mask
- Plastic oxygen tubing
- Oxygen source or compressed air

Older children may use the handheld mouthpiece instead of the face mask.

Steps for Using a Nebulizer With a Handheld Mouthpiece

General steps for using a nebulizer with a handheld mouthpiece ([Figure 36](#)) include the following:

1. Unscrew the cap of the nebulizer reservoir, add medication (eg, albuterol) to the nebulizer reservoir, and reattach the cap.
2. Connect the bottom of the T-piece to the top of the nebulizer reservoir, and then connect the spacer to one end of the T-piece and the mouthpiece to the other end.
3. Connect plastic tubing between the bottom of the nebulizer bottle and the pressurized oxygen/gas source.

4. Set the gas flow at 5 to 6 L/min.
5. Hold the nebulizer reservoir upright during delivery of the medication through the mouthpiece. Place the mouthpiece in the child's mouth and show him how to hold it. Tell the child, "Take long, slow, deep breaths through your mouth." Continue the treatment (about 8-10 minutes) until the nebulizer reservoir is empty and no mist flows from the T-piece.



Figure 36. Child receiving a treatment by nebulizer and handheld mouthpiece.

Steps for Using a Nebulizer With a Face Mask

Follow these steps to use a nebulizer with a face mask:

1. Unscrew the cap of the nebulizer reservoir, add medication (eg, albuterol) to the nebulizer reservoir, and reattach the cap.
2. Attach the face mask to the top of the nebulizer reservoir.
3. Connect plastic tubing between the bottom of the nebulizer reservoir and the pressurized oxygen/gas source.
4. Set the gas flow at 5 to 6 L/min.
5. Hold the nebulizer reservoir upright during delivery of the medication through a face mask. Place the face mask over the child's face so that it covers both the nose and mouth; press the mask to the face to ensure a light seal. Tell the child, "Take long, slow, deep breaths through your mouth." Continue the treatment (about 8-10 minutes) until the nebulizer bottle is empty and no mist flows from the mask.

Metered-Dose Inhaler

Using a Metered-Dose Inhaler With a Spacer Device

Follow these steps to use a metered-dose inhaler (MDI) with a spacer device (with and without a face mask):

1. Remove the cap from the spacer, and insert the mouthpiece of the MDI into the rubber-sealed end of the spacer ([Figure 37](#)). Once assembled, shake the MDI and the spacer vigorously. Place the mouthpiece of the spacer device into the child's mouth.

or

If you are using a spacer device with a face mask, place the mask over the child's face so that it covers both the nose and the mouth ([Figure 38](#)). Press the mask to the face to ensure a tight seal.

2. While the child is exhaling, activate the MDI by pressing down on the inhaler to release the medication into the spacer device (or have the child do so). Tell the child to take 3 to 5 slow, deep breaths through the mouthpiece and hold the last breath for 10 seconds.

or

If you are using a spacer device with a face mask, depress the inhaler and allow the child to breathe normally through the mask for 3 to 5 breaths after release of medication.




Figure 37. Child self-administering a treatment by MDI with a spacer device.



Figure 38. Child receiving a treatment by MDI with a mask spacer device.

Pulse Oximetry

When to Use Pulse Oximetry

When caring for a seriously ill or injured child, use pulse oximetry to monitor oxygen saturation and trends in oxygen saturation. The pulse oximeter measures the percent of hemoglobin that is fully saturated or bound with oxygen. Hemoglobin is usually bound with oxygen, but it may be bound with any agent that binds to hemoglobin (eg, carbon monoxide). Oxygen saturation does not equate to oxygen delivery to the tissues, and the pulse oximeter does not provide information about the effectiveness of ventilation (carbon dioxide elimination).

Confirm the Validity of Oximeter Data

The pulse oximeter requires pulsatile blood flow to determine oxygen saturation. Different brands of pulse oximeters vary in how quickly they reflect hypoxemia development and in their accuracy when the child has decreased blood flow; all pulse oximetry devices are inaccurate unless the pulse rate displayed by the oximeter is consistent with the heart rate displayed by the cardiac monitor. Note that skin pigment does not affect the accuracy or function of the pulse oximeter.



Critical Concepts: Validity of Oximeter Data

Confirm validity of oximeter data by evaluating the child's appearance. Also compare the heart rate displayed by the pulse oximeter with the heart rate displayed by the bedside cardiorespiratory monitor or counted during physical examination.

Immediately evaluate the child if the oximeter fails to detect a signal, displays an inaccurate pulse rate, indicates a weak signal, or indicates a fall in oxygen saturation.

If the oximeter fails to detect a signal or indicates a fall in oxygen saturation, immediately evaluate the child. Do not assume that the pulse oximeter is malfunctioning.

The pulse oximeter may not be functioning correctly if

- The displayed heart rate does not correlate with the child's heart rate
- The child's appearance does not correspond with the reported concentration of oxygen saturation

Accuracy of Values in Clinical Settings

Pulse oximetry can accurately estimate oxygen saturation but does not provide evidence of oxygen delivery and does not directly evaluate the effectiveness of ventilation (carbon dioxide concentration).

Pulse oximetry may be inaccurate in some settings ([Table 37](#)).

Table 37. Settings Where a Pulse Oximeter May Be Inaccurate

Setting	Cause/solution
Cardiac arrest	Cause: Absence of blood flow Solution: None. Many devices will not be useful during cardiac arrest.
Shock or hypothermia	Cause: Decreased blood flow Solution: Improve blood flow (treat the shock). You may be able to find an alternative site (particularly one that is closer to the heart) where the device can detect pulsatile blood flow.

Motion, shivering, or bright overhead lighting	Cause: False signals and inaccurate oxygen saturation values Solution: Move the sensor unit closer to the heart. Lightly cover the device in position (ie, if it is placed on a finger, lightly cover the finger to reduce ambient light).
Problem with the skin probe interface	Cause: Low or absent pulse signals Solution: Try an alternate site or alternate skin probe
Misaligning sensor with light source	Cause: Low or absent pulse signals Solution: Reposition the device so that the light source is located directly across the tissue bed from the sensor.
Cardiac arrhythmias with low cardiac output	Cause: Arrhythmias that interfere with detection of a pulse and calculation of pulse rate Solution: Assess pulse rate manually. Obtain expert consultation for arrhythmia management.

Correct Use of Oximetry Equipment

Correct probe positioning, typically on a finger or toe, is critical for accurate oxygen saturation values. Falsely low values can occur when the probe is not positioned correctly. When this occurs, repositioning the probe can result in immediate improvement in detecting oxygen saturation by the device.

In addition to applying the probe to a finger or toe, the following locations can be used to solve placement problems: If infant probes are unavailable, use an adult probe around the hand or foot of an infant. If blood flow is significantly reduced, and no signal is detected in the extremities, assess and support systemic perfusion and apply an infant probe to the earlobe.

Endotracheal Intubation

Potential Indications

Consider ET intubation if the child is unable to maintain effective airway, oxygenation, or ventilation despite initial intervention.

Preparing for Endotracheal Intubation

In the Airway Management Skills Station, you will need to know the equipment needed for ET intubation ([Table 38](#)).

Table 38. Pre-event Equipment Checklist for Endotracheal Intubation

Check	Equipment
	Universal precautions (gloves, masks, eye protection)
	Cardiac monitor, pulse oximeter, and blood pressure monitoring device
	End-tidal CO ₂ detector or exhaled CO ₂ capnography (or esophageal detector device, if appropriate)
	Intravenous and intraosseous infusion equipment
	Oxygen supply, bag mask (appropriate size)
	Oral/tracheal suction equipment (appropriate size); confirm that it is working
	Oral and nasopharyngeal airways (appropriate size)
	Endotracheal tubes with stylets (all sizes) and sizes 0.5 mm (i.d.) above and below anticipated size for patient it is reasonable to choose cuffed ETTs over uncuffed ETTs
	Laryngoscope (curved and straight blades) and/or video laryngoscope; backup laryngoscope available
	3-, 5-, and 10-mL syringes to test inflate endotracheal tube balloon
	Adhesive/cloth tape or commercial endotracheal tube holder to secure tube
	Towel or pad to align airway by placing under head or torso
	Specialty equipment as needed for difficult airway management or anticipated complications (supraglottic, transtracheal, and/or cricothyrotomy)

Sudden Deterioration in an Intubated Patient (DOPE Mnemonic)

Sudden deterioration in an intubated patient may be caused by one of several complications. Use the mnemonic **DOPE** to help remember these:

- Displacement of the tube: The tube may be displaced out of the trachea or advanced into the right or left main bronchus.
- Obstruction of the tube: Obstruction may be caused by secretions, blood, pus, a foreign body, or kinking of the tube.
- Pneumothorax: Simple pneumothorax usually results in a sudden deterioration in oxygenation (reflected by a sudden decrease in SpO_2) and decreased chest expansion and breath sounds on the involved side. Tension pneumothorax may result in the above plus evidence of hypotension and decreased cardiac output. The trachea is usually shifted away from the involved side.
- Equipment failure: Equipment may fail for several reasons, such as disconnection of the O_2 supply from the ventilation system, leak in the ventilator circuit, failure of power supply to the ventilator, and malfunction of valves in the bag or circuit.

Evaluating the Patient's Status

If an intubated patient's condition deteriorates, first support oxygenation and ventilation. While attempting this, rapidly assess the child and attempt to determine and correct the cause of deterioration. If the child is being mechanically ventilated, hand ventilate with a bag while you assess the patient's airway, ventilation, and oxygenation as follows:

- Observe for chest rise and symmetry of chest movement.
- Auscultate over both sides of the anterior chest and at the midaxillary line and over the stomach. Listen carefully over the lateral lung fields for asymmetry in breath sounds or abnormal sounds such as wheezing.
- Check monitors (eg, pulse oximetry and, if available, capnography).
- Check heart rate.
- Suction the ET tube if you suspect obstruction with secretions.
- Use sedatives or analgesics, with or without neuromuscular blockers, if needed to reduce the child's agitation and control ventilation. Administer these agents only *after* you rule out a correctable cause of the acute distress and are sure that you can provide positive-pressure ventilation.

Your initial assessment will determine the urgency of the required response. If you cannot verify that the ET tube is in the airway, direct visualization of the tube passing through the glottis is advised. If the child's condition is deteriorating and you strongly suspect that the tube is no longer in the trachea, you may need to remove it and ventilate with a bag-mask device.

Patient Agitation

Once you confirm the ET tube position and openness/patency and rule out failure of ventilation equipment and pneumothorax, evaluate oxygenation and perfusion. If oxygenation and perfusion are adequate or unchanged, it is possible that agitation, pain, or excessive movement is interfering with adequate ventilation.

If so, try one or more of the following:

- Analgesia (eg, fentanyl or morphine) to control pain
- Sedation (eg, lorazepam, midazolam) for anxiety or agitation
- Neuromuscular blocking agents and analgesia or sedation to optimize ventilation and minimize the risk of barotrauma and unintentional tube displacement

Continuous capnography is the gold standard during mechanical ventilation as an adjunct to clinical assessment. A sudden decrease in exhaled CO_2 can indicate ET tube displacement or cardiac arrest, while a gradual decrease in exhaled CO_2 may indicate development of ET tube obstruction or decreasing cardiac output. In addition, capnography may help to detect hypoventilation or hyperventilation, and so is particularly useful during transport and diagnostic procedures. Use a colorimetric detector or capnography during intrahospital and interhospital transport.

Part 9: Recognizing Shock

When shock is present, rapidly identifying and promptly intervening are critical to improving outcomes. If left untreated, shock can quickly progress to cardiopulmonary failure followed by cardiac arrest. Once an infant or child develops cardiac arrest secondary to inadequately treated shock, the outcome is poor.

This Part discusses the following:

- Pathophysiology of shock
- Effect of different types of shock on blood pressure
- Systolic blood pressure as a method of categorizing the severity of shock (ie, compensated or hypotensive)
- Etiology and signs of the 4 most common types of shock
- Systematic approach to evaluating the cardiovascular system

Your evaluation will help you identify the child's shock based on type and severity. These clinical findings will direct your interventions, as discussed in [Part 10: Managing Shock](#). The earlier you recognize shock, establish priorities, and start therapy, the better the child's chance for a good outcome.

Learning Objective

After completing this Part, you should be able to differentiate between compensated and hypotensive shock.

During the course, you will need to identify different types and severities of shock. Evaluating patient information will help you determine effective interventions.

Defining Shock

Shock is defined as a physiologic state characterized by inadequate tissue perfusion to meet metabolic demand and tissue oxygenation. It is often, but not always, characterized by inadequate peripheral and end-organ perfusion. In children, most shock is characterized by low cardiac output; however, in some types of shock (eg, that caused by sepsis or anaphylaxis), cardiac output may be high. All types of shock can result in impaired function of vital organs such as the brain (decreased level of consciousness) and kidneys (low urine output, ineffective filtering).



Critical Concepts: Shock and Blood Pressure

The definition of shock does not require the presence of hypotension. Shock can be present with a normal, increased, or decreased systolic blood pressure.

Shock can result from

- Inadequate blood volume or oxygen-carrying capacity (hypovolemic shock, including hemorrhagic shock)
- Inappropriate distribution of blood volume and flow (distributive shock)
- Impaired cardiac contractility (cardiogenic shock)
- Obstructed blood flow (obstructive shock)

Conditions such as fever, infection, injury, respiratory distress, and pain may contribute to shock by increasing tissue demand for O_2 and nutrients. Whether due to inadequate supply, increased demand, or a combination of both, *O_2 and nutrient delivery to the tissues is inadequate relative to metabolic needs*. Inadequate tissue perfusion can lead to tissue hypoxia, anaerobic metabolism, accumulating lactic acid and CO_2 , irreversible cell damage, and, ultimately, organ damage. Death may then result rapidly from cardiovascular collapse or more slowly from multiorgan failure.



Critical Concepts: Goal for Treating Shock

When treating shock, the goal is to improve systemic perfusion and O_2 delivery to help prevent end-organ injury and stop the progression to cardiopulmonary failure and cardiac arrest.

Pathophysiology of Shock

The major function of the cardiopulmonary system is to deliver O_2 to body tissues and remove metabolic by-products of cellular metabolism (primarily CO_2). When O_2 delivery cannot adequately meet tissue demand, cells use anaerobic metabolism to produce energy, but this generates lactic acid as a by-product. Anaerobic metabolism can only maintain limited cell function. Unless O_2 delivery is restored, organ dysfunction or failure will result.



Critical Concepts: Central Venous O_2 Saturation ($ScvO_2$) and Cardiac Output

In healthy children with normal metabolic demand, the arterial blood contains more O_2 than the tissues need. If demand increases and/or O_2 delivery decreases, the tissues will extract a greater percent of the O_2 delivered. This results in reduced O_2 saturation in the venous blood returning to the heart. The $ScvO_2$, therefore, can be used to assess the balance between O_2 delivery and demand. If metabolic demand and O_2 content are unchanged, a decreased $ScvO_2$ indicates a fall in cardiac output and, therefore, a fall in O_2 delivered to the tissues. In addition, greater extraction of O_2 occurs as the result of reduced delivery.

Components of Tissue Oxygen Delivery

Adequate tissue O_2 delivery (Figure 39) depends on

- Sufficient O_2 content in the blood
- Adequate blood flow to the tissues (cardiac output)
- Appropriate distribution of blood flow to the tissues




Figure 39. Factors influencing O_2 delivery.

The hemoglobin concentration and percent of the hemoglobin that is saturated with O_2 (ie, the arterial oxygen saturation, or SaO_2) primarily determines blood O_2 content. Plasma carries a small amount of dissolved oxygen.



Critical Concepts: Compensatory Mechanisms for Hypoxemia

Tissue hypoxia is present when a region of the body or an organ is deprived of adequate O_2 supply. Low O_2 saturation (hypoxemia) alone does not necessarily result in tissue hypoxia. Oxygen delivery to the tissues is the product of the arterial O_2 content (determined by the oxygen bound to hemoglobin plus dissolved O_2) and the volume of blood pumped per minute (cardiac output). O_2 delivery may be normal despite hypoxemia if cardiac output increases commensurate with the decrease in O_2 content.

When hypoxemia is chronic (eg, unrepaired cyanotic heart disease), hemoglobin concentration increases (polycythemia). The increased hemoglobin concentration will increase the O_2 -carrying capacity of the blood and help maintain arterial O_2 content at near-normal concentrations, despite hemoglobin oxygen saturation being low.

If cardiac output decreases or hypoxemia worsens, these compensatory mechanisms may not be sufficient to maintain tissue O_2 delivery, and tissue hypoxia will likely develop.

Cardiac output and vascular resistance determine adequate blood flow to the tissues. Cardiac output (the volume of blood pumped by the heart per minute) is the product of stroke volume (the volume of blood pumped by the ventricles with each contraction) and heart rate (number of times the ventricles contract per minute):

Cardiac Output = Stroke Volume × Heart Rate

According to this formula, if the heart rate decreases, stroke volume must increase commensurately to maintain the cardiac output. Cardiac output can increase either by an increase in heart rate, in stroke volume, or both. However, the increased cardiac output produced by increasing heart rate does have its limit. If the heart rate is too fast, as can happen with extreme tachyarrhythmias (eg, supraventricular tachycardia), stroke volume can fall because there is inadequate time to fill the heart (ie, the diastolic phase is too short). Some arrhythmias, such as complete heart block or junctional tachycardia, can be associated with a decrease in ventricular filling because atrial contraction (which normally contributes about 25% of ventricular filling) does not precede ventricular contraction.



Critical Concepts: Infant Cardiac Output Dependent on Heart Rate

Infants have a very small stroke volume with limited ability to increase, therefore making them dependent on an adequate heart rate to maintain or increase cardiac output. The adolescent and adult are better able to increase stroke volume, and cardiac output is less dependent on the heart rate.

Appropriate blood flow distribution is determined by the size of the blood vessels supplying a specific organ. This property is known as *vascular resistance*. If the vessel is large, vascular resistance is low; if the vessel is small, vascular resistance is high. Vascular resistance is adjusted by tissues to locally regulate blood flow to meet metabolic demands. Abnormally increased resistance (vasoconstriction) or decreased resistance (vasodilation) can interfere with blood flow distribution, even if cardiac output is adequate.

Stroke Volume

Stroke volume, the amount of blood ejected by the ventricles with each contraction, is determined by the following 3 factors:

- Preload: volume of blood present in the ventricle before contraction
- Contractility: strength of contraction
- Afterload: resistance against which the ventricle is ejecting

Inadequate *preload* is the most common cause of low stroke volume and, therefore, low cardiac output. Several conditions (eg, hemorrhage, severe dehydration, vasodilation) can cause inadequate preload, which results in hypovolemic shock.

You may assess preload indirectly by measuring the central venous pressure, but the relationship between central venous pressure and preload is complex. Preload to the ventricles is the volume of blood in the ventricles that stretches the ventricular fibers before a contraction (ie, the end-diastolic volume). However, the measurements used clinically measure pressure. The relationship between changes in ventricular volume and changes in ventricular end-diastolic pressure is affected by ventricular compliance (or, conversely, ventricular stiffness).

In general, assess the right ventricle preload by measuring the central venous pressure, measured in the superior vena cava or right atrium. Generally, increased central venous pressure corresponds to increased right ventricular end-diastolic volume and preload. However, if there is increased pressure around the right atrium and ventricle from a tension pneumothorax or pericardial tamponade, or if the right ventricle is stiff (ie, not compliant) because of a congenital heart defect or pulmonary hypertension, ventricular end-diastolic pressure increases despite no increase (or even a decrease) in right ventricular end-diastolic volume and preload.

Preload is not the same as total blood volume. At steady state, most of the blood (about 70%) is in the veins. If the veins dilate while the total blood volume is maintained, an inadequate amount of blood may return to the heart (reduced preload) because of a drop in the driving pressure. This is part of the problem with sepsis: there is often severe venodilation, so preload to the heart may be inadequate. In addition, the maldistribution of blood flow results in some tissue hypoxia.

Poor contractility (also referred to as *myocardial dysfunction*) impairs stroke volume and cardiac output, potentially leading to cardiogenic shock. Poor contractility can be due to an intrinsic problem with pump function or an acquired abnormality, such as an inflamed heart muscle (ie, myocarditis). Poor contractility also can occur from metabolic problems, such as hypoglycemia, or from toxic ingestions (eg, calcium channel blockers).

Increased afterload is an uncommon primary cause of low stroke volume and impaired cardiac output in children. When cardiac output reduces, the body responds with vasoconstriction to maintain blood pressure and blood flow to vital organs. Paradoxically, vasoconstriction increases impedance to ventricular ejection and further decreases stroke

volume and cardiac output. Certain conditions, such as severe pulmonary or systemic hypertension or congenital abnormalities of the aorta, can increase afterload so significantly that cardiogenic shock results.

Compensatory Mechanisms

As shock develops, compensatory mechanisms attempt to maintain O₂ delivery to vital organs. These mechanisms include

- Tachycardia
- Increased systemic vascular resistance (SVR) (vasoconstriction)
- Increased strength of cardiac contraction (contractility)
- Increase in venous smooth muscle tone

The body's first action to maintain cardiac output is to increase heart rate (tachycardia), which can increase cardiac output to a limited degree.

When O₂ delivery to the tissues is compromised, blood flow is redirected or shunted from nonvital organs and tissues (eg, skin, skeletal muscles, gut, kidneys) to vital organs (eg, brain, heart). This redirection occurs by a selective *increase in SVR (vasoconstriction)*. Clinically, this results in reduced peripheral perfusion (ie, delayed capillary refill, cool extremities, weaker peripheral pulses) and reduced perfusion to the gut and kidneys (decreased urine volume).

Another compensatory mechanism to maintain stroke volume and cardiac output is *increased strength of cardiac contractions (contractility)* with more complete emptying of the ventricles. Stroke volume may also be supported by *increased venous smooth muscle tone*, improving venous return to the heart and preload.

Effect on Blood Pressure

Blood pressure is determined by cardiac output and SVR. As cardiac output decreases, blood pressure can be maintained by an increase in SVR. In children with shock, this compensatory mechanism can be so effective that systolic blood pressure may initially remain normal or even slightly elevated. Pulse pressure, the difference between the systolic and diastolic blood pressure, often narrows because increased SVR raises the diastolic pressure. In contrast, if SVR is low (as in sepsis), diastolic blood pressure decreases and pulse pressure widens.

If cardiac output is inadequate, tissue perfusion is compromised, even if blood pressure is normal. Signs of poor tissue perfusion, including lactic acidosis and end-organ dysfunction, will be present even if blood pressure is normal.

When SVR cannot increase further, blood pressure begins to decline. Then, O₂ delivery to vital organs is severely compromised. Clinical signs include metabolic acidosis and evidence of end-organ dysfunction (eg, impaired mental status and decreased urine output). Ultimately O₂ delivery to the myocardium becomes inadequate, causing myocardial dysfunction, decreased stroke volume, and hypotension. These may rapidly lead to cardiovascular collapse, cardiac arrest, and irreversible end-organ injury.

Identifying Shock by Severity (Effect on Blood Pressure)

Shock severity is frequently characterized by its effect on systolic blood pressure. Shock is described as *compensated* if compensatory mechanisms can maintain a systolic blood pressure within a normal range (ie, above the fifth percentile systolic blood pressure for age). When compensatory mechanisms fail and systolic blood pressure declines, shock is classified as *hypotensive* (previously referred to as *decompensated*). It's important to remember that shock may be present even if the child's blood pressure is normal.

You can easily identify hypotensive shock by measuring blood pressure, while you may find it more difficult to diagnose compensated shock. Shock can range from mild to severe, with its manifestations affected by the type of shock and the child's compensatory responses. Use blood pressure to determine shock severity; however, children with both compensated and hypotensive shock are at high risk for deterioration. The child with low cardiac output (ie, hypovolemic shock) but normal mean blood pressure associated with severe vasoconstriction may have more end-organ compromise than the child with normal or increased cardiac output (ie, septic shock) and low diastolic blood pressure.

Because blood pressure is one method of categorizing shock severity, it is important to recognize that automated blood pressure devices are accurate only when there is adequate distal perfusion. If you cannot feel distal pulses, and the extremities are cool and poorly perfused, automated blood pressure values may not be reliable. Treat the child based on your entire clinical evaluation. If measuring blood pressure is not feasible, clinically evaluate tissue perfusion to guide treatment.

Compensated Shock

Compensated shock refers to a clinical state in which clinical signs of inadequate tissue perfusion appear, but the patient's blood pressure is in the normal range. In this stage of shock, the body can maintain blood pressure despite impaired delivery of O₂ and nutrients to the vital organs. Clinical findings include tachycardia, delayed capillary refill, mental status changes, and decreased urine output.



Critical Concepts: Systolic Blood Pressure in Identifying Shock

Compensated shock refers to the child with signs of poor perfusion but a normal systolic blood pressure (ie, with blood pressure compensation). Systolic blood pressure is used by convention and consensus to determine the presence or absence of hypotension with shock. Infants and children with compensated shock may be critically ill despite an adequate systolic blood pressure.

When O₂ delivery is limited, compensatory mechanisms try to maintain normal blood flow to the brain and heart and act as clues to the presence of shock, varying according to the type of shock. [Table 39](#) lists common compensatory mechanisms in shock and the cardiovascular signs associated with these mechanisms.

Table 39. Common Signs of Shock Resulting From Cardiovascular Compensatory Mechanisms

Compensatory mechanism	Area	Sign
Increased heart rate	Heart	Tachycardia
Increased SVR	Skin	Cold, pale, mottled, diaphoretic
	Peripheral circulation	Delayed capillary refill
	Pulses	Weak peripheral pulses; narrow pulse pressure (increased diastolic blood pressure)
Increased renal and splanchnic vascular resistance (redistribution of blood flow away from these areas)	Kidney	Oliguria (decreased urine output)
	Intestine	Vomiting, ileus
Cerebral autoregulation	Brain	Altered mental status, anxiety/restlessness, disorientation or decreased level of consciousness or even coma

Signs specific to shock type are discussed in [Identifying Shock by Type](#) later in this Part.

Hypotensive Shock

Hypotensive (decompensated) shock can result from many causes and is characterized by evidence of impaired perfusion that will rapidly progress to cardiac arrest if not corrected. Signs include abnormal clinical appearance and evidence of severely impaired perfusion (ie, weak or absent distal pulses and weak central pulses, cool extremities, mottled skin, or altered level of consciousness). Shock represents a continuum of severity, and the presence of signs and symptoms of shock should prompt immediate action rather than waiting for direct measurement of blood pressure to document hypotension.

Hypotension is a late finding in most types of shock and may signal impending cardiac arrest. It can occur early in septic shock because mediators of sepsis produce vasodilation and reduce SVR. In this setting, the child may initially appear to have warm extremities, brisk capillary refill, and full peripheral pulses despite hypotension.



Critical Concepts: Hypotension in Septic Shock


Hypotension formula: In children 1 to 10 years of age, hypotension is present if the systolic blood pressure is less than $70 \text{ mm Hg} + [\text{child's age in years} \times 2] \text{ mm Hg}$.

For more information, see [Table 17: Definition of Hypotension by Systolic Blood Pressure and Age](#) in [Part 4](#).

Systolic blood pressure is a complex combination of the patient's cardiac output, intravascular volume, and SVR. Septic shock is a dynamic process and does not always follow a specific clinical presentation; serial assessments are mandatory because the clinical presentation is likely to evolve over time. Hypotension can be an early or late sign in septic shock and must be immediately recognized and treated aggressively.

Accelerating Process

Shock progression is unpredictable. It may take hours for compensated shock to progress to hypotensive shock but only minutes for hypotensive shock to progress to cardiopulmonary failure and cardiac arrest. This progression is typically called an *accelerating process* (Figure 40).



Critical Concepts: Halting the Progression

Early recognition and rapid intervention are critical to halting the progression from compensated shock to hypotensive shock to cardiopulmonary failure and cardiac arrest.

These and other clinical manifestations are discussed in greater detail later in this Part.

Identifying Shock by Type

Shock can be categorized into 4 basic types (refer to [Table 48](#) later in this Part), including

- Hypovolemic: resulting from gastroenteritis, burns, hemorrhage, inadequate fluid intake, increased body fluid losses, sepsis, or osmotic diuresis
- Cardiogenic: resulting from congenital heart disease, myocarditis, cardiomyopathy, or arrhythmia
- Distributive: resulting from sepsis, anaphylaxis, or spinal cord injury
- Obstructive: resulting from tension pneumothorax, cardiac tamponade, pulmonary embolism, or constriction of the ductus arteriosus in infants with ductal-dependent congenital heart lesions (eg, coarctation, hypoplastic left ventricle)

Hypovolemic Shock

Hypovolemic shock refers to a clinical state of reduced intravascular volume. The most common type of shock in pediatric patients, it can be caused by extravascular fluid loss (eg, diarrhea, dehydration) or intravascular volume loss (eg, hemorrhage) and results in decreased preload and cardiac output. Volume loss leading to hypovolemic shock can result from

- Diarrhea
- Vomiting
- Hemorrhage (internal and external)
- Inadequate fluid intake
- Osmotic diuresis (eg, diabetic ketoacidosis [DKA])
- Third-space losses (fluid leak into tissues, eg, in sepsis)
- Large burns

Hypovolemic shock is the result of an absolute deficiency of intravascular blood volume, but, in fact, it typically represents depletion of both intravascular and extravascular fluid volume. As a result, adequate fluid resuscitation often requires administering fluid boluses that exceed the volume of the estimated intravascular deficit.

Tachypnea, a respiratory compensation to maintain acid-base balance, is often present in hypovolemic shock. The respiratory alkalosis that results from hyperventilation partially compensates for the metabolic acidosis (lactic

acidosis) that accompanies shock.

Physiology of Hypovolemic Shock

Hypovolemic shock is characterized by decreased preload leading to reduced stroke volume and low cardiac output. Tachycardia, increased SVR, and increased cardiac contractility are the main compensatory mechanisms.

- Preload: decreased
- Contractility: initially normal or increased
- Afterload: increased

Signs of Hypovolemic Shock



Table 40 outlines typical signs of hypovolemic shock found during the initial assessment and primary assessment.

Table 40. Findings Consistent With Hypovolemic Shock

Primary assessment	Finding
Airway	Typically open/patent unless level of consciousness is significantly impaired
Breathing	Tachypnea without increased effort (quiet tachypnea)
Circulation	<ul style="list-style-type: none">• Tachycardia• Adequate systolic blood pressure, narrow pulse pressure, or systolic hypotension with a narrow pulse pressure*• Weak or absent peripheral pulses• Normal or weak central pulses• Delayed capillary refill• Cool to cold, pale, mottled, diaphoretic skin• Dusky/pale distal extremities• Changes in level of consciousness• Oliguria
Disability	Decreasing level of consciousness as shock progresses
Exposure	Extremities often cooler than trunk

*Sign that distinguishes hypovolemic shock from distributive shock.

Although septic, anaphylactic, neurogenic, and other distributive forms of shock are not classified as hypovolemic, they are characterized by *relative* hypovolemia. The relative hypovolemia results from arterial and venous vasodilation, increased capillary permeability, and plasma loss into the interstitium (“third spacing” or capillary leak).

Distributive Shock

Distributive shock refers to a clinical state characterized by reduced SVR leading to maldistribution of blood volume and blood flow. This group includes

- Septic shock
- Anaphylactic shock
- Neurogenic shock (eg, spinal injury)

In septic and anaphylactic shock, there may also be increased capillary permeability, leading to volume loss from the intravascular space (ie, decreased preload). In neurogenic shock, there is loss of sympathetic tone leading to vasodilation and lack of compensatory mechanisms (ie, tachycardia and peripheral vasoconstriction).

Distributive shock caused by sepsis is typically characterized by reduced SVR resulting in maldistribution of blood flow. The vasodilation and venodilation cause blood pooling in the venous system and a relative hypovolemia. Septic

shock also causes increased capillary permeability, so the vascular space loses plasma. This increases hypovolemia severity. Myocardial contractility may also be depressed in septic shock.

In anaphylaxis, shock is caused by arterial and venous dilation and increased capillary permeability leading to reduced blood pressure. A compensatory increase in cardiac index may be seen, but ultimately, relative hypovolemia will impair blood pressure.

Neurogenic shock is characterized by generalized loss of vascular tone, most often after a high cervical spine injury, leading to severe vasodilation and hypotension. Normally, the sympathetic nervous system increases heart rate in response to hypotension. Children with neurogenic shock may be unable to generate a faster heart rate in response to hypotension. As a result, cardiac output and blood flow to tissues decrease dramatically.

Pathophysiology of Distributive Shock

In distributive shock, cardiac output may be increased, normal, or decreased. Although myocardial dysfunction may be present, stroke volume can be adequate, particularly if there is aggressive volume resuscitation and decreased SVR. Tachycardia and increased ventricular end-diastolic volume (from volume resuscitation) help maintain cardiac output. Tissue perfusion is compromised by maldistribution of blood flow. Some tissue beds (eg, splanchnic circulation, renal circulation) may be inadequately perfused; in other tissues (eg, some skeletal muscle and skin) perfusion may exceed metabolic needs. Hypoxic tissues generate lactic acid, leading to metabolic acidosis. Early in the clinical course, a child with distributive shock may present with decreased SVR and increased blood flow to the skin, producing warm extremities and bounding peripheral pulses (“warm shock”).

The typically high cardiac output and low SVR often observed in distributive shock differ from the low cardiac output and high SVR seen in cardiogenic and obstructive shock. As distributive shock progresses, concomitant hypovolemia and/or myocardial dysfunction produce a decrease in cardiac output. SVR can then increase, resulting in inadequate blood flow to the skin, cold extremities, and weak pulses (“cold shock”). The late phase of distributive shock, therefore, can be similar to the clinical picture of cardiogenic shock.

Distributive shock is most often characterized by many changes in cardiovascular function, including

- Low SVR, which typically leads to a lower diastolic blood pressure, is the etiology of the wide pulse pressure that is commonly, but not always, present in the early phases of distributive shock
- Increased blood flow to some peripheral tissue beds
- Inadequate perfusion of the splanchnic and kidney vascular beds
- Release of inflammatory and other mediators and vasoactive substances
- Volume depletion caused by capillary leak
- Accumulating lactic acid in poorly perfused tissue beds



Critical Concepts: Relative Hypovolemia

Although most types of distributive shock are not typically classified as hypovolemic shock, all are characterized by relative hypovolemia unless adequate fluid resuscitation is provided.

Signs of Distributive Shock



Table 41 outlines typical signs of distributive shock seen during the initial assessment and primary assessment.

Table 41. Findings Consistent With Distributive Shock

Primary assessment	Finding
Airway	Usually open/patent unless level of consciousness is significantly impaired
Breathing	Tachypnea, usually without increased work of breathing (“quiet tachypnea”) unless the child has additional compromise as seen in pneumonia, ARDS, pulmonary edema, or

	bronchospasm
Circulation	<ul style="list-style-type: none"> • Tachycardia (predominantly); bradycardia (very rarely) (eg, neurogenic shock typically due to catastrophic high cervical spinal cord injury); bounding (early)* or decreased peripheral pulses (late) • Flash (early)* or delayed capillary refill (late) • Warm, flushed skin peripherally (warm extremities, early)* or Pale, mottled skin with vasoconstriction (cold extremities, late) • Hypotension with a wide pulse pressure (when warm extremities are present, early)* or Hypotension with a narrow pulse pressure (when cold extremities are present, late) or Normotension may be present • Changes in level of consciousness • Oliguria
Disability	Changes in level of consciousness
Exposure	<ul style="list-style-type: none"> • Fever or hypothermia • Extremities may be warm (early) or cool (late) • Petechial or purpuric rash (may be seen in septic shock) or urticarial rash (anaphylaxis)*

*Sign that distinguishes distributive shock from other forms of shock.

Septic Shock

Sepsis represents an important cause of shock in infants and children. *Sepsis* and *septic shock* are terms used to characterize shock caused by an infectious agent or inflammatory stimulus.

Septic shock is the most common form of distributive shock. It is caused by an abnormal host immune response to infectious organisms or their by-products (eg, endotoxin) that lead the small blood vessels to dilate and to leak fluid into the tissues.

Infection is a pathologic process caused by pathogenic or potentially pathogenic microorganisms invading normally sterile tissue, fluid, or a body cavity. You may suspect or prove infection by positive result of a culture, tissue stain, or polymerase chain reaction test. Without these tests, evidence of infection includes positive findings on clinical examination, imaging, or laboratory tests consistent with tissue invasion by a pathogenic organism leading to a host response (eg, white blood cells in normally sterile body fluid, perforated viscus, chest radiograph consistent with pneumonia, petechial or purpuric rash, purpura fulminans).

Pathophysiology of Septic Shock

Septic shock in children typically evolves along a continuum from a systemic inflammatory response in the early stages to septic shock in the late stages. Because of the wide variability in clinical presentation and progression, this continuum may evolve over days or just a few hours.

- Preload: decreased
- Cardiac output: normal or increased (early); decreased (late)
- Afterload (eg, SVR): variable

Inflammatory Cascade Response to Sepsis

The pathophysiology of the septic cascade includes the following, often referred to as the *systemic inflammatory response*:

- The infectious organism or its by-products (eg, endotoxin) activates the immune system, including neutrophils, monocytes, and macrophages.
- These cells, or their interaction with the infecting organism, stimulate release or activation of inflammatory mediators (cytokines) that perpetuate the inflammatory response.
- Cytokines produce vasodilation and damage to the lining of the blood vessels (endothelium), causing increased capillary permeability.
- Cytokines activate the coagulation cascade and may result in microvascular thrombosis and disseminated intravascular coagulation (DIC).
- Specific inflammatory mediators can impair cardiac contractility and cause myocardial dysfunction.

The Challenge of Treating Septic Shock

In septic shock, inadequate perfusion combined with possible microvascular thrombosis leads to ischemia, which is diffuse and patchy so that individual organs have varying levels of hypoxia and ischemia. The variability of perfusion throughout the body is what makes treating sepsis so difficult.

Adrenal Insufficiency in Septic Shock

The adrenal glands are especially prone to microvascular thrombosis and hemorrhage in septic shock. Because adrenal glands produce cortisol, an important hormone in the body's stress response, children with sepsis may develop absolute or relative adrenal insufficiency. Adrenal insufficiency contributes to low SVR and myocardial dysfunction in septic shock.

Signs of Septic Shock

In the early stages, signs of septic shock are often subtle and may be difficult to recognize because peripheral perfusion may initially appear adequate. Because septic shock is triggered by an infection or its by-products, the child may have fever or hypothermia, and the white blood cell (WBC) count may be decreased, normal, or increased.

In addition to the findings listed in the Identifying Shock by Type section, the child with septic shock may have other abnormalities identified by diagnostic assessments. Examples include metabolic acidosis, respiratory alkalosis, leukocytosis (high WBC count), leukopenia (low WBC count), or left shift (increased percent of bands or immature white blood cells). With some types of infections, the child may develop a petechial or purpuric rash.

Central ScvO₂ in Septic Shock

In contrast to hypovolemic and cardiogenic shock, ScvO₂ may be normal, increased, or decreased in septic shock. There are 2 mechanisms to explain a normal, or even increased, ScvO₂, despite the presence of inadequate cardiac output/index. Because of this potential situation in septic shock, lactate is a more useful marker of adequate oxygen delivery than ScvO₂.

- Children with low SVR and increased cardiac output will extract less O₂ from the blood because some tissues are receiving more blood flow than they need. Other tissues don't receive enough blood flow, so they don't have the opportunity to extract the O₂.
- Children with sepsis may be unable to use O₂ at the cellular level.

Toxins and inflammatory mediators circulating in sepsis can prevent aerobic metabolism even in the setting of adequate O₂ delivery. As a result, lactic acidosis and end-organ dysfunction can occur even in the setting of normal or increased ScvO₂.

Identify and Intervene: Septic Shock

Early recognition and treatment of septic shock are critically important determinants of outcome. Evaluate temperature, heart rate, systemic perfusion, blood pressure, and clinical signs of end-organ function to identify sepsis and septic shock before severe organ dysfunction develops. If you suspect sepsis, and especially if shock develops, provide appropriate volume resuscitation and hemodynamic support (refer to [Part 10: Managing Shock](#) for details). Search for and treat the underlying cause.

Physiology of Anaphylactic Shock

Anaphylactic shock, an acute multisystem response caused by a severe reaction to a drug, vaccine, food, toxin, plant, venom, or other antigen, is characterized by venodilation, arterial vasodilation, and increased capillary permeability. It can occur within seconds to minutes after exposure.

Signs of Anaphylactic Shock

Signs and symptoms may include those listed in [Table 42](#).

Table 42. Findings Consistent With Anaphylactic Shock

Primary assessment	Finding	Resulting from
Airway	Angioedema (swelling of the face, lips, and tongue)*	Swelling of the tongue and tissues related to fluid leak from blood vessels
Breathing	Respiratory distress with stridor, wheezing, or both*	Constriction of airways by inflammatory response
Circulation	Hypotension	Vasodilation, hypovolemia, and diminished cardiac output

Circulation	Tachycardia	Inadequate blood flow to the tissues
Disability	Anxiety and agitation	Low oxygen concentration and reduced brain perfusion
Exposure	Urticaria (hives)*	Histamine release
Other	Nausea and vomiting	Histamine and other mediator release

*Sign that distinguishes anaphylactic shock from other forms of shock.

Angioedema may partially or completely obstruct the upper airway. Hypotension results from vasodilation, hypovolemia, and diminished cardiac output. Relative hypovolemia is caused by the vasodilation, and absolute volume loss is caused by capillary leak.

Neurogenic Shock

Neurogenic shock, also known as *spinal shock*, results from a cervical (neck) or upper thoracic (above T6) injury that disrupts the sympathetic nervous system innervation of blood vessels and of the heart.

Physiology of Neurogenic Shock

The sudden loss of sympathetic nervous system signals to the smooth muscle in the vessel walls results in uncontrolled vasodilation. The same disruption prevents tachycardia developing as a compensatory mechanism.

- Preload: decreased
- Contractility: normal
- Afterload: decreased

Signs of Neurogenic Shock

Primary signs of neurogenic shock are

- Hypotension with a wide pulse pressure
- Normal heart rate or bradycardia
- Hypothermia

Other signs may include increased respiratory rate, diaphragmatic breathing (the use of muscles in the diaphragm rather than the chest wall), and other evidence of a high thoracic or cervical spine injury (ie, motor or sensory deficits).

You must differentiate neurogenic shock from hypovolemic shock. Hypovolemic shock is typically associated with hypotension, a narrow pulse pressure from compensatory vasoconstriction, and compensatory tachycardia. In neurogenic shock, these compensatory mechanisms are not apparent because sympathetic innervation of the heart and blood vessels is interrupted.

Cardiogenic Shock

Cardiogenic shock refers to reduced cardiac output secondary to abnormal cardiac function or pump failure, resulting in decreased systolic function and cardiac output. Common causes of cardiogenic shock include

- Congenital heart disease
- Myocarditis (inflamed heart muscle)
- Cardiomyopathy (an inherited or acquired abnormality of pumping function)
- Arrhythmias
- Sepsis
- Poisoning or drug toxicity
- Myocardial injury (eg, trauma)

Physiology of Cardiogenic Shock

Marked tachycardia, high SVR, and decreased cardiac output characterize cardiogenic shock. End-diastolic volume within the left and right ventricles increases, resulting in congestion within the pulmonary and systemic venous systems. This pulmonary venous congestion leads to pulmonary edema and increased work of breathing. Typically, intravascular volume is normal or increased unless a concurrent illness causes hypovolemia (eg, in a child who has viral myocarditis with recent vomiting, fever, poor oral intake).

- Preload: variable

- Contractility: decreased
- Afterload: increased

Cardiogenic shock is often characterized by sequential compensatory and pathologic mechanisms, including

- Increased heart rate and left ventricular afterload, which increases left ventricular work and myocardial O₂ consumption
- Compensatory increase in SVR to redirect blood from peripheral and splanchnic tissues to the heart and brain
- Decreased stroke volume due to decreased myocardial contractility and increased afterload
- Increased venous tone, which increases central venous (right atrial) and pulmonary capillary (left atrial) pressures
- Diminished renal blood flow resulting in fluid retention
- Pulmonary edema resulting from myocardial failure and high left ventricular end-diastolic, left atrial, and pulmonary venous pressures, and from increased venous tone and fluid retention

The same compensatory mechanisms that maintain perfusion to the brain and heart in hypovolemic shock are often detrimental during cardiogenic shock. For example, compensatory peripheral vasoconstriction can maintain blood pressure in hypovolemic shock, but it increases left ventricular afterload (commonly thought of as increased resistance to left ventricular ejection) in cardiogenic shock.

Because the heart muscle also needs O₂, almost all children with severe or sustained shock may eventually have inadequate myocardial O₂ delivery relative to myocardial O₂ demand. Therefore, severe or sustained shock of any type eventually causes impaired myocardial function (ie, these children develop cardiogenic shock in addition to the primary cause of shock). Once myocardial function declines, the child's clinical status usually deteriorates rapidly.

Signs of Cardiogenic Shock



Table 43 outlines signs of cardiogenic shock typically found during the initial assessment and primary assessment of the child.

Table 43. Findings Consistent With Cardiogenic Shock

Primary assessment	Finding
Airway	Usually open/patent unless level of consciousness is significantly impaired
Breathing	<ul style="list-style-type: none"> • Tachypnea • Increased respiratory effort (retractions, nasal flaring, grunting) resulting from pulmonary edema*
Circulation	<ul style="list-style-type: none"> • Tachycardia • Normal or low blood pressure with a narrow pulse pressure • Weak or absent peripheral pulses • Normal and then weak central pulses • Delayed capillary refill with cool extremities • Signs of congestive heart failure (eg, pulmonary edema, hepatomegaly, jugular venous distention, gallop, murmur)* • Cyanosis (caused by cyanotic congenital heart disease or pulmonary edema)* • Cold, pale, mottled, diaphoretic skin • Changes in level of consciousness • Oliguria
Disability	Changes in level of consciousness
Exposure	Extremities often cooler than trunk

*Sign that distinguishes cardiogenic shock from other forms of shock.



Critical Concepts: Distinguishing Signs of Cardiogenic Shock

Increased respiratory effort often distinguishes cardiogenic shock from hypovolemic shock. Hypovolemic shock is characterized by “quiet tachypnea,” while children with cardiogenic shock may demonstrate retractions, grunting, and use of accessory muscles.

In cardiogenic shock, decreased arterial O_2 saturation secondary to pulmonary edema may present.

Rapidly providing volume resuscitation of *cardiogenic* shock in the setting of poor myocardial function can aggravate pulmonary edema and further impair myocardial function, further compromising oxygenation, ventilation, and cardiac output. Gradually provide volume resuscitation for cardiogenic shock; give smaller (5-10 mL/kg) boluses of isotonic crystalloid and deliver over a longer period of time (ie, over 10-20 minutes). Carefully monitor hemodynamic parameters during fluid infusion, and repeat infusion as needed.

Infants and children with cardiogenic shock often require medications to increase and redistribute cardiac output, improve myocardial function, and reduce SVR. Additional treatment includes reducing metabolic demand, such as decreasing work of breathing and controlling fever, which allows a limited cardiac output to better meet tissue metabolic demands. For more details, refer to [Part 10: Managing Shock](#).

Obstructive Shock

Obstructive shock refers to conditions that physically impair blood flow by limiting venous return to the heart or limit the pumping of blood from the heart, resulting in decreased cardiac output. Causes of obstructive shock include

- Pericardial tamponade
- Tension pneumothorax
- Ductal-dependent congenital heart defects (eg, coarctation of the aorta, hypoplastic left ventricle)
- Massive pulmonary embolism

The physical obstruction to blood flow results in low cardiac output, inadequate tissue perfusion, and a compensatory increase in SVR. The early clinical presentation of obstructive shock can be indistinguishable from hypovolemic shock. However, careful clinical examination may reveal signs of systemic or pulmonary venous congestion that are not consistent with hypovolemia. In the case of massive PE, tension pneumothorax, and tamponade, the onset is usually acute, as opposed to hypovolemia, which has a more gradual progression to shock. As the condition progresses, increased respiratory effort, cyanosis, and signs of vascular congestion become more apparent.

- Preload: variable
- Contractility: normal
- Afterload: increased

Pathophysiology and Clinical Signs of Obstructive Shock

Pathophysiology and clinical signs vary according to the cause of the obstructive shock.

Cardiac Tamponade



Cardiac tamponade is caused by accumulating fluid, blood, or air in the pericardial space. Increased intrapericardial pressure and compression of the heart impede systemic venous and pulmonary venous return. This reduces ventricular filling and causes decreased stroke volume and cardiac output. If untreated, cardiac tamponade results in cardiac arrest with pulseless electrical activity.

In children, cardiac tamponade most often occurs after penetrating trauma or cardiac surgery. It also may develop as a result of pericardial effusion complicating an inflammatory disorder, an infection of the pericardium, a tumor, or an extremely high white blood cell count. [Table 44](#) outlines signs of cardiac tamponade typically found during the initial assessment and primary assessment.

Table 44. Findings Consistent With Cardiac Tamponade

Primary assessment	Finding
Airway	Usually open/patent unless level of consciousness is significantly impaired
Breathing	Respiratory distress with increased respiratory rate and effort
Circulation	<ul style="list-style-type: none"> • Tachycardia • Poor peripheral perfusion (weak distal pulses, cool extremities, delayed capillary refill) • Muffled or diminished heart sounds* • Narrowed pulse pressure • Pulsus paradoxus (decrease in systolic blood pressure by >10 mm Hg during spontaneous inspiration)* • Distended neck veins (may be difficult to see in infants, especially with severe hypotension)
Disability	Changes in level of consciousness
Exposure	Extremities often cooler than trunk

*Sign unique to pericardial tamponade.

After cardiovascular surgery in children, signs of tamponade may be indistinguishable from those of cardiogenic shock. Favorable outcome depends on urgent diagnosis and immediate treatment. In children with a large pericardial effusion, the electrocardiogram typically shows small QRS complexes (low voltage), but echocardiography provides a definitive diagnosis. Pulsus paradoxus is an exaggerated manifestation of a normal variation in stroke volume that occurs during the phases of spontaneous respiration. Stroke volume decreases slightly during inspiration and increases slightly during expiration. In pulsus paradoxus, the systolic blood pressure declines by greater than 10 mm Hg on inspiration, compared with expiration. True assessment for pulsus paradoxus requires measuring blood pressure with a manual pressure cuff. Inflate the cuff until no sounds are heard (as usual). Slowly decrease the cuff pressure and note when you initially hear the Korotkoff sounds, which will be when the child is exhaling. Continue to slowly deflate the cuff and note when you consistently hear the Korotkoff sounds throughout the respiratory cycle. If the difference between these 2 points is greater than 10 mm Hg, the child has a clinically significant pulsus paradoxus.

You may be able to detect a significant pulsus paradoxus by palpating the pulse, noting a distinct variation in pulse amplitude as the child inhales and exhales. Pulsus paradoxus may also be apparent on arterial and pulse oximetry waveforms but is not as easily quantified unless the waveform can be saved on the monitor screen or printed for review.

Tension Pneumothorax



Tension pneumothorax is caused by air entering the pleural space and accumulating under pressure. This air can enter from lung tissue injured by an internal tear or from a penetrating chest injury. An air leak that enters the pleural space but then stops spontaneously is called a *simple pneumothorax*. An ongoing leak can result from positive-pressure ventilation or chest trauma that forces air out of the injured lung and into the pleural space. If air continues to leak into the pleural space, it accumulates under pressure, creating a tension pneumothorax. As this pressure increases, it compresses the underlying lung and pushes the mediastinum to the opposite side of the chest. Compression of the lung rapidly causes respiratory failure. The high intrathoracic pressure and direct pressure on mediastinal structures (heart and great vessels) impede venous return, resulting in a rapid decline in cardiac output and hypotension. Untreated tension pneumothorax leads to cardiac arrest characterized by pulseless electrical activity.

Suspect tension pneumothorax in a victim of chest trauma or in any intubated child who deteriorates suddenly while receiving positive-pressure ventilation (including bag-mask or noninvasive ventilation). [Table 45](#) outlines signs of tension pneumothorax typically found during the initial assessment and primary assessment.

Table 45. Findings Consistent With Tension Pneumothorax

--	--

Primary assessment	Finding
Airway	<ul style="list-style-type: none"> Variable depending on situation and primary cause of respiratory distress Advanced airway may already be in place Tracheal deviation toward contralateral side (the side opposite the side of the pneumothorax). This deviation may be difficult to appreciate in infants.*
Breathing	<ul style="list-style-type: none"> Respiratory distress with increased respiratory rate and effort Hyperresonance of affected side; hyperexpansion of affected side* Diminished or absent breath sounds on affected side*
Circulation	<ul style="list-style-type: none"> Distended neck veins (may be difficult to appreciate in infants or in children with severe hypotension) Pulsus paradoxus (decrease in systolic blood pressure by >10 mm Hg during spontaneous inspiration) Rapid deterioration in perfusion; commonly, rapid evolution from tachycardia to bradycardia and hypotension as cardiac output decreases
Disability	Changes in level of consciousness
Exposure	Extremities often cooler than trunk

*Sign(s) unique to tension pneumothorax.

Favorable outcome depends on immediate diagnosis and treatment.

Ductal-Dependent Lesions



Ductal-dependent congenital cardiac abnormalities usually present in the first days to weeks of life. Ductal-dependent lesions include

- Cyanotic congenital heart lesions (ductal dependent for pulmonary blood flow)
- Left ventricular outflow obstructive lesions (ductal dependent for systemic blood flow)

The congenital heart lesions that depend on the ductus for pulmonary blood flow present with cyanosis rather than signs of shock. The left ventricular outflow obstructive lesions often present with signs of obstructive shock in the first few days or weeks of life when the ductus arteriosus closes. These left heart and aortic lesions include coarctation of the aorta, interrupted aortic arch, critical aortic valve stenosis, and hypoplastic left heart syndrome. Restoring and maintaining openness/patency of the ductus arteriosus is critical for survival until surgical intervention is possible, because the ductus serves as a conduit for systemic blood flow that bypasses the obstruction. If the infant is to survive, you must quickly recognize the presence of a ductal-dependent lesion and promptly provide treatment to open and maintain an open/patent ductus arteriosus.

Table 46 outlines findings consistent with left ventricular outflow obstructive lesions that may be found while evaluating the child.

Table 46. Findings Consistent With Left Ventricular Outflow Obstructive Lesions

Primary assessment	Finding
Airway	Usually open/patent unless level of consciousness is significantly impaired
Breathing	Respiratory failure with signs of pulmonary edema or inadequate respiratory effort
Circulation	<ul style="list-style-type: none"> Rapid progressive deterioration in systemic perfusion Congestive heart failure (cardiomegaly, hepatomegaly) Higher preductal vs postductal blood pressure (coarctation or interrupted aortic arch)* Higher (greater than 3% to 4%) preductal vs postductal arterial O₂ saturation (coarctation or interrupted aortic arch)* Absent or weaker femoral pulses (coarctation or interrupted aortic arch)* Metabolic acidosis (elevated lactate)
Disability	Rapid deterioration in level of consciousness

Exposure	Cool skin
-----------------	-----------

*Sign unique to left ventricular outflow obstructive lesions.

Massive Pulmonary Embolism



Pulmonary embolism is a total or partial obstruction of the pulmonary artery or its branches by a blood clot, fat, air, amniotic fluid, catheter fragment, or injected matter. Most commonly, a pulmonary embolus is a thrombus that migrates to the pulmonary circulation. Pulmonary embolism is rare in children but may develop when an underlying condition predisposes the child to intravascular thrombosis. Examples include immobility, central venous catheters, sickle cell disease, malignancy, connective tissue disorders, and inherited disorders of coagulation (eg, antithrombin III, protein S, and protein C deficiencies).

Pulmonary embolism results in ventilation/perfusion mismatch, hypoxemia, increased pulmonary vascular resistance leading to right heart failure, decreased left ventricular filling, and decreased cardiac output. Pulmonary embolism may be difficult to diagnose because signs may be subtle and nonspecific (cyanosis, tachycardia, and hypotension). Providers may not suspect pulmonary embolus, especially in children. However, signs of systemic venous congestion and right heart failure as well as acute onset do help distinguish it from hypovolemic shock. Some children with pulmonary embolism will say they have chest pain, reflecting lack of oxygenated blood flow to the lung tissue itself.

Table 47 outlines findings consistent with pulmonary embolism that may be found while evaluating the child.

Table 47. Findings Consistent With Pulmonary Embolism

Primary assessment	Finding
Airway	Usually open/patent unless level of consciousness is significantly impaired
Breathing	Respiratory distress with increased respiratory rate and effort
Circulation	<ul style="list-style-type: none"> • Tachycardia • Cyanosis • Hypotension • Systemic venous congestion and right heart failure • Chest pain
Disability	Changes in level of consciousness
Exposure	Extremities may be cool and mottled

Summary

Treatment of obstructive shock is cause specific; immediately recognizing and correcting the underlying cause of the obstruction can save a life. The most critical tasks for PALS providers are prompt recognition, diagnosis, and treatment of obstructive shock. Without them, children with obstructive shock often progress rapidly to cardiopulmonary failure and cardiac arrest.

Recognizing Shock Flowchart

Table 48 summarizes recognition of the 4 basic types of shock.

Table 48. Recognizing Shock Flowchart

Clinical signs		Hypovolemic shock	Distributive shock	Cardiogenic shock	Obstructive shock
Airway	Patency	Airway open and maintainable/not maintainable			
Breathing	Respiratory rate	Increased			
	Respiratory effort	Normal to increased		Labored	

	Breath sounds	Normal	Normal (± crackles)	Crackles, grunting
Circulation	Systolic blood pressure	Compensated shock can progress to hypotensive shock if left untreated		
	Pulse pressure	Narrow	Variable	Narrow
	Heart rate	Increased		
	Peripheral pulse quality	Weak	Bounding or weak	Weak
	Skin	Pale, cool	Warm or cool	Pale, cool
	Capillary refill	Delayed	Variable	Delayed
	Urine output	Decreased		
Disability	Level of consciousness	Irritable early, lethargic late		
Exposure	Temperature	Variable		

Part 10: Managing Shock

Once you identify shock in a critically ill or injured child, intervene early to reduce morbidity and mortality. This Part discusses the goals and priorities of shock management, treatment fundamentals, general and advanced management, and specific management according to etiology.

Learning Objective

After completing this Part, you should be able to perform early interventions for treating shock.

During the course, you will be asked to manage children in shock. To do this, you must know general and specific treatment based on the different types of shock.

Goals of Shock Management

The goals in treating shock are to

- Improve O₂ delivery
- Balance tissue perfusion and metabolic demand
- Support organ function
- Prevent progression to cardiac arrest

Immediate intervention for a child in shock may be lifesaving. The more time that passes between the onset of signs of shock and the restoration of adequate O₂ delivery and organ perfusion, the poorer the outcome is. Once a child in shock progresses to cardiac arrest, prognosis is poor.

Warning Signs

Watch for signs that compensatory mechanisms are failing in a seriously ill or injured child. Once you recognize that the child's condition is deteriorating, act decisively with the resuscitation team to provide effective resuscitation therapy. Identifying compensated shock early is critical to effective treatment and good outcome. Warning signs that indicate progression from compensated to hypotensive shock include

- Increasing tachycardia
- Diminishing or absent peripheral pulses
- Weakening central pulses
- Narrowing pulse pressure
- Cold distal extremities with prolonged capillary refill
- Decreasing level of consciousness
- Hypotension (late finding)

Once the child develops hypotensive shock, organ perfusion is typically severely compromised, and organ dysfunction may develop even if the child does not progress to cardiac arrest.

Fundamentals of Shock Management

The acute treatment of shock focuses on restoring O₂ delivery to the tissues and improving the balance between tissue perfusion and metabolic demand. This treatment consists of

- Optimizing O₂ content of the blood
- Improving volume and distribution of cardiac output
- Reducing O₂ demand
- Correcting metabolic derangements

Try to identify and reverse the underlying cause of shock while promptly intervening.

Optimizing Oxygen Content of the Blood

O₂ content of the blood is determined by the hemoglobin concentration and its saturation with oxygen. To optimize O₂ content

- Administer a high concentration of O₂ (use nonrebreathing mask to deliver 100% O₂)
- Use invasive or noninvasive mechanical ventilation to improve oxygenation by correcting a ventilation/blood flow (V/Q) mismatch or other respiratory disorders
- If hemoglobin concentration is low, consider packed red blood cell (PRBC) transfusion

Improving Volume and Distribution of Cardiac Output

For most forms of shock, use bolus fluid administration to improve volume and distribution of cardiac output. You may consider noninvasive or invasive positive-pressure ventilation to reduce the work of breathing and improve oxygenation. Children in shock may also benefit from vasoactive agents such as vasopressors, vasodilators, inodilators, and/or inotropes.

Reducing Oxygen Demand

For all forms of shock, try to improve the balance between O₂ delivery and supply by reducing O₂ demand. The most common factors that contribute to increased O₂ demand are

- Increased work of breathing
- Pain and anxiety
- Fever

Support breathing with noninvasive or invasive ventilation and assisted ventilation. To facilitate intubation and mechanical ventilation, you may administer sedatives or analgesics and neuromuscular blockade. You may also need to control pain and anxiety with analgesics and sedatives. Use sedative and analgesic agents with extreme caution; they may suppress the child's endogenous stress response, impair compensatory mechanisms such as tachycardia, and reduce blood pressure. The sedative effects of these agents can also make it more difficult to evaluate the child's level of consciousness and response to treatment. Control fever by administering antipyretics and other cooling measures.

Correcting Metabolic Derangements

Many conditions that lead to shock may result in or be complicated by metabolic derangements, such as

- Hypoglycemia
- Hypocalcemia
- Hyperkalemia
- Metabolic (lactic) acidosis

All of these conditions can adversely affect cardiac contractility. Metabolic acidosis is characteristic of all forms of shock.

Hypoglycemia is low serum glucose concentration that, if left untreated, can cause seizures and brain injury. Glucose is vital for proper cardiac and brain function. Glucose stores may be low in infants and chronically ill children.

Hypocalcemia is a low serum ionized calcium concentration. Calcium is essential for effective cardiac function and vasomotor tone. Hypocalcemia can result from administering blood products, colloid, and buffering medications such as sodium bicarbonate.

Hyperkalemia is a high serum potassium concentration, which may result from renal dysfunction, cell death, excess potassium administration, or acidosis. Acidosis causes a shift of potassium from the intracellular to the extracellular—including the intravascular—space. As a result, acidosis, or a fall in serum pH, typically is associated with a rise in serum potassium. The serum potassium will fall when acidosis is corrected or alkalosis develops.

Metabolic acidosis develops from production of acids, such as lactic acid, when tissue perfusion is inadequate. Renal or gastrointestinal dysfunction can also cause metabolic acidosis. Renal dysfunction can cause organic acid retention or bicarbonate ion loss while gastrointestinal dysfunction, such as diarrhea, can result in bicarbonate ion loss. Severe metabolic acidosis may depress myocardial contractility and reduce the effect of vasopressors. Unless metabolic acidosis is due solely to bicarbonate losses, it does not respond well to buffer therapy. Treat the acidosis by attempting to restore tissue perfusion with fluid resuscitation and vasoactive agents. If treatment is effective, the metabolic acidosis will resolve.

On occasion, you may need buffers (eg, sodium bicarbonate) to acutely correct profound metabolic acidosis that is impairing vital organ function. Sodium bicarbonate works by combining with hydrogen ions (acids) to produce carbon dioxide and water; increased alveolar ventilation then eliminates the carbon dioxide. Ventilation support is always important in the critically ill child, but it is especially important for treating metabolic acidosis with sodium bicarbonate. One scenario in which sodium bicarbonate should be avoided is DKA, where the HCO_3^- can be profoundly low but giving HCO_3^- can result in worse outcomes.

Correcting metabolic derangements may be essential to optimizing organ function. Measure ionized calcium concentration (the active form of calcium in the body) and glucose concentration, and replenish them as indicated. It is important to measure glucose a few times during resuscitation, particularly if it is borderline or the patient has already received dextrose for a previously low glucose. Consider administering sodium bicarbonate to treat metabolic acidosis refractory to attempts to increase cardiac output or redistribute blood flow to vital organs.

General Management of Shock

Components of General Management

General management of shock consists of the following:

- Positioning
- Supporting airway, oxygenation, and ventilation
- Establishing vascular access
- Providing fluid resuscitation
- Monitoring
- Performing frequent reassessment
- Obtaining laboratory studies
- Providing medication therapy
- Consulting appropriate subspecialists

Note that several of these interventions may be implemented by the team simultaneously.

Positioning

Initial management of shock includes positioning the critically ill or injured child. If the child is responsive and hemodynamically stable, allow her to remain in the most comfortable position (eg, sitting in the arms of a caregiver) to decrease anxiety and activity as you form your initial assessment and conduct the primary assessment. If the child is hypotensive and breathing is not compromised, place the child in the supine position.

Supporting Airway, Oxygenation, and Ventilation

Maintain an open/patent airway and support oxygenation and ventilation. Give a high concentration of supplemental O_2 to all children with shock, usually by a high-flow O_2 delivery system. Sometimes you must combine O_2 delivery with ventilatory support if respirations are ineffective, mental status is impaired, or work of breathing is significantly increased. Appropriate interventions may include noninvasive positive airway pressure or mechanical ventilation after endotracheal intubation.

Vascular Access

Once the airway is open/patent and oxygenation and ventilation are supported, establish vascular access for fluid resuscitation and administering medications. For compensated shock, initial attempts at peripheral venous cannulation are appropriate. For hypotensive shock, you can best accomplish critical immediate vascular access by the intraosseous (IO) route if peripheral IV access is not readily achieved. Depending on the provider's experience and expertise and clinical circumstances, central venous access may be useful. However, gaining central venous access takes longer than placing IO access.



Critical Concepts: IO Access

If peripheral vascular access cannot be readily obtained in a child with compensated or hypotensive shock, be prepared to establish IO access.

For more information on establishing IO access, refer to [Intraosseous Access](#) in [Resources for Managing Circulatory Emergencies](#) in Part 10 of this manual.

Fluid Resuscitation

Once you establish vascular access, start fluid resuscitation immediately.



Critical Concepts: Fluid Resuscitation

In general, isotonic crystalloid should be given as a bolus over 5 to 20 minutes: 10 to 20 mL/kg for children with suspected septic shock and 20 mL/kg for all other shock etiologies (unless otherwise specified). In children with severe, hypotensive, hypovolemic shock, fluid should be given over 5 to 10 minutes. If you suspect cardiogenic shock, use smaller fluid boluses of 5 to 10 mL/kg given over 10 to 20 minutes. Carefully monitor for signs of pulmonary edema or worsening tissue perfusion. Stop the infusion if such signs occur. Be prepared to support oxygenation and ventilation as necessary.

Reassess* and repeat boluses to restore blood pressure and tissue perfusion.

*Repeat fluid boluses based on clinical signs of end-organ perfusion, including heart rate, capillary refill, level of consciousness, and urine output.

Monitoring

No single resuscitation end point has been identified as a consistent marker of adequate tissue perfusion and cellular homeostasis. However, you can assess the effectiveness of fluid resuscitation and medication therapy by frequent or continuous monitoring ([Table 49](#)).

Table 49. Monitoring in Circulatory Emergencies

Frequently or continuously monitor	Indication of positive response to shock therapy
Oxygen saturation with pulse oximetry	94% or higher when breathing room air
Heart rate	Appropriate for age and clinical condition. Typically will fall from a rapid rate toward a normal range (refer to Table 14: Normal Heart Rates in Part 4)
Peripheral pulses	Weak pulses will become stronger, and bounding pulses will be less bounding but still strong
Capillary refill	Will shorten toward 2 seconds or less
Skin color and temperature	Normal skin color and mucous membranes; warm extremities
Blood pressure	Increase to within normal range for age (Table 16: Normal Blood Pressures in Part 4) with normal pulse pressure
Level of consciousness	Child will respond more appropriately (ie, mental status improves)
Ongoing fluid losses	Bleeding and diarrhea controlled
Urine output	<i>Infants and young children</i> approximately 1.5 to 2 mL/kg per hour <i>Older children and adolescents</i> approximately 1 mL/kg per hour

As soon as possible, start noninvasive monitoring (SpO₂, heart rate, blood pressure), assess mental status, and measure temperature and measure urine output with an indwelling bladder catheter. Also consider invasive monitoring (eg, arterial and central venous catheterization), depending on the providers' experience and available resources. Signs that indicate clinical improvement toward a normal hemodynamic state are decreased serum lactate, smaller base deficit, and central venous oxygen saturation (ScvO₂) greater than 70%.



Critical Concepts: Accurately Assess Tissue Perfusion

Although measuring blood pressure is easy, it is important to assess other clinical parameters to evaluate tissue perfusion. Remember that blood pressure may be normal in children with severe shock, and noninvasive blood pressure measurement may be inaccurate if perfusion is poor.

Frequent Reassessment

Frequently reassess the child's respiratory, cardiovascular, and neurologic status to

- Evaluate trends in the child's condition
- Determine response to therapy
- Plan the next interventions

A child in shock is in a dynamic clinical condition that can deteriorate at any moment and require lifesaving interventions, such as endotracheal intubation. Frequently reassess until the child's condition becomes stable or the child is transferred to advanced care.



Critical Concepts: Monitor to Evaluate Trends

The condition of a child in shock is dynamic. Continuously monitor and frequently reassess to evaluate trends in the child's condition and determine response to therapy.

Laboratory Studies

Laboratory studies provide important information to help you

- Identify the etiology and severity of shock
- Evaluate organ dysfunction secondary to shock
- Identify metabolic derangements
- Evaluate response to therapy

Refer to [Part 13: Post–Cardiac Arrest Care](#) for additional information on evaluating end-organ function. Also consider expert consultation in diagnosis and management of end-organ failure.

[Table 50](#) outlines some laboratory studies that can help identify the etiology and severity of shock and guide therapy.

Table 50. Laboratory Studies to Evaluate Shock and Guide Therapy

Laboratory study	Finding	Possible etiology	Possible interventions
Complete blood cell count	Hemoglobin/hematocrit decreased	<ul style="list-style-type: none">• Hemorrhage• Fluid resuscitation (dilution)• Hemolysis	<ul style="list-style-type: none">• Administer 100% O₂• Control bleeding• Transfuse blood• Titrate fluid administration
	White blood cell count increased or decreased	<ul style="list-style-type: none">• Sepsis	<ul style="list-style-type: none">• Obtain appropriate cultures• Give antibiotics
	Platelets decreased	<ul style="list-style-type: none">• Disseminated intravascular coagulation• Decreased platelet production	<ul style="list-style-type: none">• Transfuse platelets if child has serious bleeding• Obtain prothrombin time/partial thromboplastin time, fibrinogen, and D-dimers
Glucose	Increased or decreased	<ul style="list-style-type: none">• Stress (usually increased but may be decreased in infants)• Sepsis• Decreased production (eg, liver failure)• Adrenal insufficiency	<ul style="list-style-type: none">• If hypoglycemia is present, give dextrose bolus and start infusion of dextrose-containing solution if needed• Severe hyperglycemia may require treatment (per institutional protocols or obtain expert consultation)
Potassium	Increased or decreased	<ul style="list-style-type: none">• Renal dysfunction• Acidosis (increases serum potassium concentration)• Diuresis (decreased)• Adrenal insufficiency (increased)	<ul style="list-style-type: none">• Treat significant or symptomatic hyperkalemia or hypokalemia• Correct acidosis
Calcium	Decreased (ionized)	<ul style="list-style-type: none">• Sepsis	<ul style="list-style-type: none">• Give calcium

	calcium concentration)	<ul style="list-style-type: none"> Transfusion of blood preserved with citrate-phosphate-dextran Colloid administration Buffering agents (eg, sodium bicarbonate) 	
Lactate	Increased as product of anaerobic metabolism from tissue hypoperfusion	<ul style="list-style-type: none"> Tissue hypoxia Increased glucose production (gluconeogenesis) Decreased metabolism (eg, liver failure, inborn error of metabolism, toxic ingestion) 	<ul style="list-style-type: none"> Improve tissue perfusion Treat acidosis if end-organ function is impaired Consider other unusual etiologies for lactic acidosis if lactate not improving with improved systemic oxygen delivery Consider buffering acidosis in specific situations
Arterial blood gas	pH decreased in acidosis; increased with alkalosis	<ul style="list-style-type: none"> Lactic acid accumulation caused by tissue hypoperfusion Renal failure Inborn error of metabolism Diabetic ketoacidosis Poisoning/overdose Diarrhea or ileostomy losses Hyper/hypoventilation (sepsis, poisoning) Vomiting 	<ul style="list-style-type: none"> Give fluid Support ventilation Correct shock Consider buffer Evaluate anion gap* to determine if acidosis is from increased unmeasured ions (increased anion gap) or is more likely from loss of bicarbonate (normal anion gap)
ScvO₂	Variable	<ul style="list-style-type: none"> Low central venous O₂ saturation—inadequate O₂ delivery or increased consumption High central venous O₂ saturation—maldistribution of blood flow or decreased O₂ utilization 	<ul style="list-style-type: none"> Attempt to maximize O₂ delivery and minimize O₂ demand

*Anion gap = ([serum Na⁺] – ([serum Cl[–]] + [serum HCO₃[–]])); normal = approximately 8 to 12 mEq/L.

Medication Therapy

Use medication therapy when managing shock to affect myocardial contractility, heart rate, and vascular resistance. The choice of agent(s) is determined by the child's physiologic state.

Vasoactive agents are indicated when shock persists despite adequate volume resuscitation to optimize preload. For example, a child with septic shock who remains hypotensive with signs of vasodilation despite fluid bolus administration may benefit from a vasoconstrictor. Administering vasoactive medications is potentially harmful if the child has not been adequately fluid resuscitated first. However, in children with cardiogenic shock, use vasoactive agents early because fluid resuscitation is not key to improving myocardial function and may contribute to pulmonary edema and respiratory failure. Most children with cardiogenic shock benefit from a vasodilator (provided that blood pressure is adequate) to decrease systemic vascular resistance (SVR) and increase cardiac output and tissue perfusion.

Inotropes, phosphodiesterase inhibitors (eg, the inodilator milrinone), vasodilators, and vasopressors are classes of pharmacologic agents commonly used in shock. [Table 51](#) lists vasoactive medications by class and pharmacologic effects.

Table 51. Vasoactive Therapy Used in Treating Shock

Class	Medication	Effect
Inotropes	<ul style="list-style-type: none"> Epinephrine Dobutamine Dopamine 	<ul style="list-style-type: none"> Increase cardiac contractility Increase heart rate Produce variable effects on SVR <p><i>Note:</i> Includes agents with both α-adrenergic and β-adrenergic effects</p>
Phosphodiesterase inhibitors	<ul style="list-style-type: none"> Milrinone 	<ul style="list-style-type: none"> Decrease SVR

(inodilators)		<ul style="list-style-type: none"> • Improve coronary artery blood flow • Improve contractility • Improve diastolic relaxation (filling)
Vasodilators	<ul style="list-style-type: none"> • Nitroglycerin • Nitroprusside 	<ul style="list-style-type: none"> • Decrease SVR and venous tone
Vasopressors (vasoconstrictors)	<ul style="list-style-type: none"> • Epinephrine (doses >0.1 mcg/kg per minute) • Norepinephrine • Dopamine (doses >10 mcg/kg per minute) • Vasopressin 	<ul style="list-style-type: none"> • Increase SVR • Increase myocardial contractility (except vasopressin)



Critical Concepts: Color-Coded Length-Based Tape

Use a color-coded length-based tape to determine the child's weight (if not known) for calculating drug doses and for selecting the correct sizes of resuscitation equipment. Refer to [Resources for Managing Circulatory Emergencies](#) in Part 10 of this manual for an example.

Subspecialty Consultation

For specific shock categories, some required lifesaving diagnostic assessments and therapeutic interventions may be beyond many PALS providers' scope of practice. For example, a provider may not be trained to interpret an echocardiogram or perform a thoracostomy or pericardiocentesis. Recognize limitations to your own scope of practice and call for help when needed. Early subspecialty consultation (eg, pediatric critical care, pediatric cardiology, pediatric surgery) is essential to shock management and may influence outcome.



Critical Concepts: Expert Consultation

When treating a child in shock, providers must obtain consultation from appropriate experts as soon as possible.

Summary: Initial Management Principles

This summary reviews initial shock management principles discussed in this section:

- Positioning the child
 - Stable—Allow to remain with caregiver in a position of comfort
 - Unstable—If hypotensive, place in supine position unless breathing is compromised
- Optimizing arterial O₂ content
 - Administer a high concentration of O₂ via a nonrebreathing mask; consider blood transfusion in cases of significant blood loss or other causes of severe anemia; and consider using continuous positive airway pressure, noninvasive positive airway pressure, or mechanical ventilation with positive end-expiratory pressure
- Supporting ventilation as indicated (invasive or noninvasive)
- Establishing vascular access
 - Consider IO access early
- Beginning fluid resuscitation
 - Give an isotonic crystalloid bolus of 20 mL/kg over 5 to 20 minutes (5-10 minutes with severe, hypotensive, hypovolemic shock); repeat 20 mL/kg boluses as needed to restore blood pressure and tissue/organ perfusion. For children with shock due to suspected sepsis, administration of 10 to 20 mL/kg is recommended, with repeat boluses of 10 to 20 mL/kg as well. Reassess the child after each bolus. For trauma and hemorrhage, administer PRBCs if the child does not respond to isotonic crystalloid. Modify bolus fluid therapy to deliver 5 to 10 mL/kg over 10 to 20 minutes if you suspect cardiogenic shock or severe myocardial dysfunction
- Monitoring
 - SpO₂, heart rate, respiratory rate and effort, blood pressure, level of consciousness, temperature, and urine output
- Performing frequent reassessment

- Evaluate trends and determine response to therapy
- Conducting laboratory studies
 - To identify shock etiology and severity, evaluate organ dysfunction secondary to shock, identify metabolic derangements, and evaluate the response to therapy
- Administering pharmacologic support—refer to [Table 51: Vasoactive Therapy Used in Treating of Shock](#)
 - To increase heart rate, improve myocardial function or redistribute cardiac output (increase contractility, reduce or increase SVR, improve organ perfusion), correct metabolic derangements, and manage pain and anxiety
 - Obtain subspecialty consultation

Fluid Therapy

The primary objective of fluid therapy in shock is to restore intravascular volume and tissue perfusion. Rapid fluid resuscitation is required for hypovolemic and distributive shock, including septic shock. Cardiogenic and obstructive shock, as well as special conditions such as severe poisonings or fluid loss with DKA, may dictate alternative approaches to fluid resuscitation.

In general, boluses of isotonic crystalloid will expand intravascular volume. Blood and blood products are generally not used for volume expansion in children with shock unless shock is due to hemorrhage. Blood products may also be indicated for correction of some coagulopathies.

Isotonic Crystalloid Solutions

For most children with shock, isotonic crystalloid solutions, such as normal saline or lactated Ringer's, are the preferred initial fluids for volume replacement in managing shock. They are inexpensive, readily available, and do not cause sensitivity reactions.



Critical Concepts: Quantity of Crystalloid Solution in Shock Resuscitation

Because isotonic crystalloids are distributed throughout the extracellular space, a large quantity of crystalloid solution may be needed to restore intravascular volume for children in shock. Rapid infusion of a large volume of fluid may be well tolerated by a healthy child but may cause pulmonary and peripheral edema in a critically ill child. Reassess after every fluid bolus.

Colloid Solutions

Colloid solutions (eg, 5% albumin and fresh frozen plasma) may be alternatives to crystalloid solutions. However, they have disadvantages for the acute resuscitation of a child in shock. They are less widely available than crystalloid solutions and may take time to prepare. Blood-derived colloid solutions may cause sensitivity reactions. Synthetic colloids may cause coagulopathies; their use is usually limited to 20 to 40 mL/kg. As with crystalloids, excessive administration of colloids can lead to pulmonary edema, particularly in children with cardiac or renal disease. Despite these limitations, fresh frozen plasma may be used in specific circumstances, such as massive hemorrhage and blood administration (refer to [Indications for Blood Product Administration](#) later in this part).

Rate and Volume of Fluid Administration

Start fluid resuscitation for shock with 20 mL/kg of isotonic crystalloid bolus over 5 to 20 minutes; in children with shock due to suspected sepsis, administer 10 to 20 mL/kg per bolus. Repeat boluses as needed to restore blood pressure and perfusion. You may often find it difficult to predict the volume of fluid deficit from the child's history, so use clinical examination and supporting laboratory studies to identify the volume needed. You may need to administer more than the estimated volume deficit. Reassess frequently and after administering each bolus.

Give fluid boluses rapidly for hypotensive and septic shock. Children with septic shock may require 60 mL/kg or more of isotonic crystalloid solution during the first hour of therapy; as much as 200 mL/kg or more may be required in the first 8 hours of therapy.



Critical Concepts: Fluid Administration

If you find or suspect myocardial dysfunction or obstructive shock, *give smaller volumes of fluid more slowly*. Administer boluses of 5 to 10 mL/kg over 10 to 20 minutes and reassess after each bolus; stop bolus administration if the child develops signs of worsening respiratory status, rales, or other evidence of pulmonary edema or hepatomegaly. Obtain further diagnostic assessment and expert consultation (eg, echocardiogram) to confirm suspicions and guide the next interventions. Be prepared to support the airway, oxygenation, and ventilation with positive end-expiratory pressure as needed if pulmonary edema develops.

Modifying fluid resuscitation is appropriate for children in shock associated with DKA. Children with DKA may be significantly dehydrated but often have high serum osmolality (caused by hyperglycemia). Rapidly administering crystalloid solution and reducing serum osmolality may contribute to risk of cerebral edema. Therefore, fluid management in DKA is complex. Consider giving an initial bolus of isotonic crystalloid 10 to 20 mL/kg over 1 to 2 hours. This constitutes a fluid bolus, but it is atypical in that it is given over a longer period of time than the usual fluid bolus. However, if the patient with DKA is in hypotensive shock, the treatment approach should default to more aggressive bolus fluids for shock per any other etiology of shock. Many institutions have local protocols about specifically managing DKA and its associated metabolic, electrolyte, and fluid derangements. Consult experts when possible. After giving a fluid bolus, reassess the patient.

Similarly, children who have ingested calcium channel blockers or β -adrenergic blockers may have myocardial dysfunction and are less tolerant of rapid volume expansion. When caring for children with severe febrile illness in settings with limited access to critical care resources (ie, mechanical ventilation and inotropic support), administer bolus IV fluids with extreme caution because it may be harmful. Once again, providers should reassess the patient after every fluid bolus.

Table 52 gives a summary of fluid boluses and rates of delivery based on the underlying cause of shock.

Table 52. Guide to Fluid Boluses and Rates of Delivery Based on Underlying Cause of Shock

Type of shock	Volume of fluid	Approximate rate of delivery
Hypovolemic shock Distributive shock (eg, anaphylaxis, neurogenic)	20 mL/kg bolus (repeat as needed)	Over 5-20 minutes; given over 5-10 minutes if severe and/or hypotensive
Cardiogenic shock (nonpoisoning)	5-10 mL/kg bolus (repeat as needed)	Over 10-20 minutes
Poisonings (eg, calcium channel blocker or β-adrenergic blocker)	5-10 mL/kg (repeat as needed)	Over 10-20 minutes
Septic shock	10-20 mL/kg (repeat as needed)	Over 5-20 minutes
DKA		
DKA with compensated shock	10-20 mL/kg	Per local protocol; if used, give over at least 1-2 hours*

*Refer to paragraph on modifying fluid resuscitation for details.

Rapid Fluid Delivery

IV fluid administration systems generally used for pediatric fluid therapy do not deliver fluid boluses as rapidly as required for managing some forms of shock. To help deliver fluid rapidly

- Place as large an IV catheter as possible, especially if blood or colloid administration is needed; ideally, insert 2 catheters.
- Place an in-line 3-way stopcock in the IV tubing system.
- Deliver fluid by using a 30- to 60-mL syringe to push fluids through the stopcock, or use a pressure bag (beware of risk of air embolism) or a rapid infusion device.
- If IV access cannot be established, establish IO access.

Note: Standard infusion pumps—even if set at the maximum infusion rate—do not provide a sufficiently rapid rate of fluid delivery, especially in larger children. For example, a 50-kg patient with septic shock should ideally receive 1 L of crystalloid in 5 to 10 minutes, but standard infusion pumps may have a maximum rate of 999 mL for the hourly rate.

Frequently Reassess During Fluid Resuscitation

Frequently reassess during fluid resuscitation to manage shock effectively. Monitor to

- Assess the physiologic response to therapy after each fluid bolus
- Determine the need for further fluid boluses
- Assess for signs of detrimental effects (eg, pulmonary edema) during and after fluid resuscitation

Signs of physiologic improvement include improved perfusion, increased blood pressure, decreasing heart rate (toward normal), decreased respiratory rate (toward normal), increased urine output, and improved mental status. If the child's condition does not improve or worsens after fluid boluses, try to identify the cause of the shock to help determine the next interventions. For example, persistently delayed capillary refill despite initial fluid administration may indicate ongoing hemorrhage or other fluid loss. Deterioration of the child's condition after fluid therapy may signal cardiogenic or obstructive shock. Increased work of breathing may indicate pulmonary edema.

Indications for Administering Blood Products

Administering PRBCs is recommended for replacing traumatic blood loss if the child's perfusion is inadequate despite receiving 2 to 3 boluses of 20 mL/kg of isotonic crystalloid. Under these circumstances, administer 10 mL/kg PRBCs as soon as available.

Fully crossmatched blood is generally not available in emergencies because most blood banks require about 1 hour for the crossmatching process. It may become available for children who are stabilized with crystalloid but have ongoing blood losses. Priorities for the type of blood or blood products used in order of preference are

- Crossmatched
- Type specific
- Type O-negative (O[–] preferred for female patients and either O[+] or O[–] for male patients)

Unmatched, type-specific blood may be used if ongoing blood loss results in hypotension despite administration of crystalloid. Most blood banks can supply type-specific blood within 10 minutes. Type-specific blood is ABO and Rh compatible but, unlike fully crossmatched blood, may have other antibodies that are incompatible with the patient's blood.

Use type O blood if blood should be administered immediately to prevent circulatory collapse or cardiopulmonary arrest, because it can be given to children of any blood type. O-negative blood is preferred for female patients of childbearing age to avoid Rh sensitization. Either O-negative or O-positive blood may be administered to male patients.

Complications of Rapid Administration of Blood Products

Rapid infusion of cold blood or blood products, particularly in large volume, may produce several complications, including

- Hypothermia
- Myocardial dysfunction
- Ionized hypocalcemia

Hypothermia may adversely affect cardiovascular function and coagulation as well as compromise several metabolic functions, including metabolism of citrate, which is present in stored blood. Inadequate citrate clearance, in turn, causes ionized hypocalcemia. The combined effects of hypothermia and ionized hypocalcemia can result in significant myocardial dysfunction and hypotension.

To minimize these problems, warm blood and blood products, if possible, with an approved commercial blood-warming device before or during rapid IV administration. Prepare calcium if the child becomes hypotensive during rapid transfusion; in some cases, it may be beneficial to administer calcium empirically to prevent hypocalcemia.

Glucose

Monitor blood glucose concentration as a component of shock management. Hypoglycemia, a common finding in critically ill children, can result in brain injury if not rapidly identified and effectively treated. In one pediatric study, hypoglycemia was present in 18% of children who received resuscitative care in an emergency department for decreased level of consciousness, status epilepticus, respiratory failure, cardiopulmonary failure, or cardiac arrest.

Glucose Monitoring

Measure serum glucose concentration as soon as possible in all critically ill infants and children (eg, those with altered mental status, respiratory compromise, shock) from capillary, venous, or arterial blood samples with a point-of-care device or by laboratory analysis. Small infants and chronically ill children have higher glucose utilization rates and limited stores of glycogen. This limited supply may rapidly deplete during episodes of physiologic stress, resulting in hypoglycemia. Infants receiving non-glucose-containing IV fluids are at increased risk for developing hypoglycemia.



Critical Concepts: Identify Hypoglycemia

In all critically ill or injured children, perform a rapid glucose test to rule out hypoglycemia as a cause of or a contributing factor to shock or decreased level of consciousness.

Hyperglycemia, also frequently present in seriously ill or injured children, may result from a relative insulin-resistant state induced by high concentrations of endogenous catecholamines and cortisol. Although controlling serum glucose concentration within a narrow range by using insulin infusion improved survival in critically ill adult and pediatric patients, tight glucose control was also associated with more frequent episodes of hypoglycemia. Insufficient data support routine use of this tight control of glucose concentration in critically ill children. Consider treating hyperglycemia in high-risk groups, such as brain-injured children, while monitoring closely to prevent hypoglycemia.

Diagnosing Hypoglycemia

Hypoglycemia may be difficult to recognize clinically by the child's appearance. Some children have no outward signs or symptoms (ie, asymptomatic hypoglycemia), while others may show nonspecific clinical signs, such as

- Poor perfusion
- Diaphoresis
- Tachycardia
- Hypothermia
- Irritability or lethargy
- Hypotension

These clinical signs are also common to many other conditions, including hypoxemia, ischemia, or shock.

Although single threshold values are not applicable to every patient, the following lowest acceptable glucose concentrations can be used to define hypoglycemia:

- Preterm and term neonates: less than 40 mg/dL
- Infants, children, and adolescents: less than 60 mg/dL

The reported low range of normal glucose is typically related to sample measurements obtained in nonstressed, fasting infants and children. It is difficult to extrapolate these thresholds to the glucose concentration required by a stressed, critically ill, or injured child.

Managing Hypoglycemia

If the glucose concentration is low and the child has minimal symptoms and normal mental status, you may administer glucose orally (eg, with orange juice or other glucose-containing fluid). If the concentration is very low or the child is symptomatic, give IV glucose at a dose of 0.5 to 1 g/kg. IV dextrose is commonly administered as D₂₅W (2-4 mL/kg) or D₁₀W (5-10 mL/kg). Dextrose is the same substance as glucose. Reassess the serum glucose concentration after dextrose administration. Provide a continuous infusion of glucose-containing IV fluid to prevent recurrent hypoglycemia.

Do not routinely infuse dextrose-containing fluids for volume resuscitation of shock. This can cause hyperglycemia, increase the serum osmolality, and produce an osmotic diuresis that will further exacerbate hypovolemia and shock. Electrolyte imbalances (eg, hyponatremia) can also develop.

Management According to Type of Shock

Effective management of shock targets treatment to the etiology of the shock. For the purposes of the PALS Provider Course, shock is categorized into 4 types, based on the underlying cause. However, this classification method oversimplifies the physiologic state seen in individual patients. Some children with shock have elements of

hypovolemic, distributive, and cardiogenic shock, with 1 type being dominant. Any child with severe shock may develop characteristics of myocardial dysfunction and maldistribution of blood flow. For a more comprehensive discussion of shock by etiology, refer to [Part 9: Recognizing Shock](#).

Managing the following types of shock is discussed in this section:

- Hypovolemic
- Distributive
- Cardiogenic
- Obstructive

Managing Hypovolemic Shock

Rapid administration of isotonic crystalloids is the primary therapy for hypovolemic shock. Children with hypovolemic shock who receive an appropriate volume of fluid within the first hour after resuscitation have the best chance for survival and recovery. Timely administration of fluid is key to preventing deterioration from compensated hypovolemic shock to hypotensive and refractory shock.



Critical Concepts: Timely Fluid Resuscitation in Hypovolemic Shock

It is important to provide rapid, adequate fluid resuscitation for hypovolemic shock. Avoid the common errors of inadequate or delayed administration of fluid resuscitation.

Other components in effectively managing hypovolemic shock are

- Identifying the type of volume loss (nonhemorrhagic vs hemorrhagic)
- Replacing volume deficit
- Preventing and replacing ongoing losses (eg, bleeding, GI losses)
- Restoring acid-base balance
- Correcting metabolic derangements

Determining Adequate Fluid Resuscitation

Dehydration is defined as a loss of water with varying loss of electrolytes leading to a hypertonic (hypernatremic), isotonic, or hypotonic (hyponatremic) state. The losses can be from some combination of the interstitial, intracellular, and intravascular compartments; the relative loss from each component helps determine clinical symptoms. Severity of dehydration is generally related to the percentage of total body water loss (ie, percent dehydration), but the percentage is not consistent across all age groups because the relative proportion of fluid loss based on total body weight is size dependent.

Adequate fluid resuscitation in hypovolemic shock is determined by the

- Extent of volume depletion
- Type of volume loss (eg, blood, electrolyte-containing fluid, or electrolyte-and-protein-containing fluid)

The extent of volume depletion may be underestimated and undertreated. In many cases, volume loss is compounded by inadequate fluid intake. Use vital signs and physical examination to assess the child's response to each fluid bolus. The clinical parameters used to help determine the percentage of dehydration include

- General appearance
- Presence or absence of tears and appearance of eyes (normal vs sunken)
- Moisture of mucous membranes
- Skin elasticity (ie, skin turgor)
- Respiratory rate and depth
- Heart rate
- Blood pressure
- Capillary refill time
- Urine output
- Mental status

Clinically significant dehydration in children is generally associated with at least 5% volume depletion (ie, 5% or greater loss in body weight) corresponding to a fluid deficit of 50 mL/kg or greater. Therefore, treating a child with clinically evident dehydration with administration of a single 20 mL/kg bolus of isotonic crystalloid may be insufficient. Conversely, it is usually unnecessary to completely correct the estimated deficit within the first hour. After perfusion is restored and the child is no longer in shock, the total fluid deficit may be corrected over the next 24 to 48 hours.

Although all forms of hypovolemic shock are initially treated with rapid infusion of isotonic crystalloid, early identification of the type of volume loss can optimize further treatment. Fluid losses may be classified as nonhemorrhagic and hemorrhagic. Nonhemorrhagic losses include electrolyte-containing fluids (eg, diarrhea, vomiting, osmotic diuresis associated with DKA) and protein-and-electrolyte-containing fluids (eg, losses associated with burns and peritonitis).

Nonhemorrhagic Hypovolemic Shock

Common sources of nonhemorrhagic fluid loss are gastrointestinal (ie, vomiting and diarrhea), urinary (eg, diabetes insipidus [DI]), and capillary leak (eg, burns). Hypovolemia caused by nonhemorrhagic fluid loss is generally classified in terms of percent loss of body weight ([Table 53](#)). Correlation of blood pressure and fluid deficits is imprecise. As a general rule, however, shock may be observed in children with fluid deficits of 50 to 100 mL/kg (particularly in hyponatremic dehydration), but it is more consistently observed with deficits of 100 mL/kg or greater.

Table 53. Stages and Signs of Dehydration

Severity of dehydration	Infant estimated weight loss (mL/kg)*	Adolescent estimated weight loss (mL/kg)*	Clinical signs	Pitfalls in assessment
Mild	5% (50)	3% (30)	<ul style="list-style-type: none"> • Dry mucous membranes • Oliguria 	<ul style="list-style-type: none"> • Oral mucosa may be dry in chronic mouth breathers • Frequency and amount of urine are difficult to assess during diarrhea, especially with infants wearing diapers
Moderate	10% (100)	5% to 6% (50-60)	<ul style="list-style-type: none"> • Poor skin turgor • Sunken fontanel • Marked oliguria • Tachycardia • Quiet tachypnea 	<ul style="list-style-type: none"> • Affected by sodium concentration; increased sodium concentration better maintains intravascular volume • Fontanel open only in infants • Oliguria is affected by fever, sodium concentration, and underlying disease
Severe	15% (150)	7% to 9% (70-90)	<ul style="list-style-type: none"> • Marked tachycardia • Weak to absent distal pulses • Narrow pulse pressure • Increased respiratory rate • Anuria • Hypotension and altered mental status (late findings) 	<ul style="list-style-type: none"> • Clinical signs are affected by fever, sodium concentration, and underlying disease; increased sodium concentration better maintains intravascular volume

*mL/kg refers to the estimated corresponding fluid deficit normalized to body weight.

Modified with permission from Roberts KB. Fluid and electrolytes: parenteral fluid therapy. *Pediatr Rev.* 2001;22(11):380-387.

Rapidly infuse 20 mL/kg boluses of isotonic crystalloid to effectively treat children with hypovolemic shock secondary to dehydration. Failure to improve after at least 3 boluses (ie, 60 mL/kg) of isotonic crystalloid indicates that

- The extent of fluid losses may be underestimated
- The type of fluid replacement may need to be altered (eg, need for colloid or blood)
- There are ongoing fluid losses (eg, occult bleeding)
- Your initial assumption about the etiology of the shock may be incorrect (ie, consider alternative or combined types of shock)

Ongoing fluid losses (eg, diarrhea, burns) must be replaced in addition to correcting existing fluid deficits. Colloid is not routinely indicated as the initial treatment of hypovolemic shock. Albumin and other colloids, however, have been used successfully for volume replacement in children with large “third-space” losses or albumin deficits.

Hemorrhagic Hypovolemic Shock

Hemorrhagic hypovolemic shock is classified according to an estimated percent of total blood volume loss ([Table 54](#)).

Table 54. Systemic Responses to Blood Loss in Pediatric Patients

System	Mild blood volume loss (<30%)	Moderate blood volume loss (30%-45%)	Severe blood volume loss (>45%)
Cardiovascular	Increased heart rate; weak, thready peripheral pulses; normal systolic blood pressure ($80-90 + 2 \times \text{age in years}$); normal pulse pressure	Markedly increased heart rate; weak, thready central pulses; low normal systolic blood pressure ($70-80 + 2 \times \text{age in years}$); narrowed pulse pressure	Tachycardia followed by bradycardia; very weak or absent central pulses; absent peripheral pulses; hypotension ($<70 + 2 \times \text{age in years}$); narrowed pulse pressure (or undetectable diastolic blood pressure)
Central nervous system	Anxious; irritable; confused	Lethargic; dulled response to pain _‡	Comatose
Skin	Cool, mottled; prolonged capillary refill	Cyanotic; markedly prolonged capillary refill	Pale and cold
Urine output[‡]	Low to very low	Minimal	None

_‡The child's dulled response to pain with this degree of blood loss (30% to 45%) may be indicated by a decreased response to IV catheter insertion.

[‡]After initial decompression by urinary catheter. Low normal is 2 mL/kg per hour (infant), 1.5 mL/kg per hour (younger child), 1 mL/kg per hour (older child), and 0.5 mL/kg per hour (adolescent).

Reproduced from American College of Surgeons Committee on Trauma. Pediatric trauma. In: *Advanced Trauma Life Support for Doctors: ATLS Student Course Manual*. 9th ed. American College of Surgeons; 2012:257.

In children, the dividing line between mild and compensated vs moderate or severe hypotensive hemorrhagic shock is thought to correlate with an acute loss of about 30% of blood volume. The estimated total blood volume of a child is 75 to 80 mL/kg; a 30% blood volume loss, therefore, represents a blood loss of about 25 mL/kg.

Fluid resuscitation in hemorrhagic shock begins with rapid infusion of isotonic crystalloid in boluses of 20 mL/kg. Because isotonic crystalloids are distributed throughout the extracellular space, it may be necessary to give up to 3 boluses of 20 mL/kg (60 mL/kg) of fluid to replace a 25% loss of blood volume; approximately 3 mL of crystalloid is needed for every 1 mL of blood lost. If the child remains hemodynamically unstable despite 2 to 3 boluses of 20 mL/kg isotonic crystalloid, PRBCs should be administered. If ongoing administration of PRBCs is needed, platelets and fresh frozen plasma should be strongly considered to avoid coagulopathy associated with multiple PRBC infusions.



Critical Concepts: 3 mL to 1 mL Rule

For fluid resuscitation in hemorrhagic shock, give about 3 mL of isotonic crystalloid for every 1 mL of blood lost.

For blood replacement, use PRBCs in 10 mL/kg boluses. Whole blood (20 mL/kg) can be given in place of PRBCs, but it is harder and more time-consuming to obtain. Also, the risk of transfusion reaction is significantly increased if the blood is not crossmatched. To minimize adverse effects, warm the blood if a blood-warming device is available, especially when transfusing rapidly.

Indications for transfusion in hemorrhagic shock include

- Crystalloid-refractory hypotension or poor perfusion
- Known significant blood loss

Crystalloid-refractory hemorrhagic shock is defined as persistent hypotension despite administration of 40 to 60 mL/kg crystalloid. Children with rapid hemorrhage may demonstrate a normal or low initial hemoglobin concentration. Transfuse blood for a low hemoglobin concentration because anemia increases the risk of tissue hypoxia from inadequate arterial O₂ content and O₂ delivery.

Medication Therapy

Vasoactive agents are not routinely indicated for managing hypovolemic shock. Moribund children with profound hypovolemic shock and hypotension may require short-term administration of vasoactive agents, such as epinephrine, to restore cardiac contractility and vascular tone once adequate fluid resuscitation is provided.

Acid-Base Balance

Early in the progression of hypovolemic shock, the child may develop tachypnea and respiratory alkalosis. However, the alkalosis does not completely correct the metabolic (lactic) acidosis produced by hypovolemic shock. A child with long-standing or severe shock may have severe acidosis because the child eventually develops fatigue or cardiorespiratory failure. Children with head or chest injuries may not demonstrate compensatory tachypnea.

Persistent acidosis and poor perfusion are indications of inadequate resuscitation or, in hemorrhagic shock, of ongoing blood loss. Sodium bicarbonate is not recommended for treating metabolic acidosis secondary to hypovolemic shock. As long as fluid resuscitation improves perfusion and end-organ function, metabolic acidosis is well tolerated and will correct gradually. Sodium bicarbonate administration is indicated if the metabolic acidosis is caused by significant bicarbonate losses from renal or gastrointestinal losses (ie, a non-anion gap metabolic acidosis) because it is difficult to compensate for an ongoing bicarbonate loss.

Specific Treatment Considerations

Follow the initial management principles listed in the [Summary: Initial Management Principles](#) section in addition to these considerations specific to hypovolemic shock:

- Initiate fluid resuscitation as quickly as possible.
 - *In all patients*, rapidly infuse isotonic crystalloid (normal saline or lactated Ringer's) in 20 mL/kg boluses; repeat as needed.
 - In patients with crystalloid-refractory hemorrhagic shock, give a transfusion of PRBCs, 10 mL/kg.
 - If loss of protein-containing fluids is documented or suspected (low albumin concentration), consider administration of colloid-containing fluids if the child fails to respond to crystalloid resuscitation.
- Correct metabolic derangements.
- Identify the type of volume loss (hemorrhagic or nonhemorrhagic) to determine best treatment.
- Control any external hemorrhage with direct pressure; measure and replace ongoing losses (eg, continued nonhemorrhagic losses with diarrhea).
- Consider other laboratory studies:
 - Complete blood cell count
 - Type and crossmatch
 - Arterial blood gas with particular attention to the base deficit
 - Electrolyte panel to calculate anion gap, glucose, and ionized calcium
 - Serum or plasma lactate concentration
 - Diagnostic imaging to identify the source of bleeding or volume loss

Managing Distributive Shock

Initial management of distributive shock focuses on expanding intravascular volume to correct hypovolemia and fill the expanded dilated vascular space. Use vasoactive agents if the child remains hypotensive or poorly perfused despite rapid bolus fluid administration or if the diastolic pressure remains low with a wide pulse pressure.

This section discusses managing the following types of distributive shock:

- Septic shock
- Anaphylactic shock

- Neurogenic shock

Managing Septic Shock

The clinical, hemodynamic, and metabolic changes observed in septic shock result from the host's response to an infection, including the release or activation of inflammatory mediators. The primary goals in the initial management of septic shock are

- Restoration of hemodynamic stability
- Support of organ function
- Identification and control of infection


Fundamental principles of management include increasing tissue O₂ delivery by optimizing cardiac output and arterial O₂ content and minimizing O₂ consumption.

Identifying septic shock early is key to initiating resuscitation and preventing multisystem organ failure and cardiac arrest from developing. Hemodynamic support to maintain O₂ delivery can reduce pediatric morbidity and mortality from septic shock.

Overview of Pediatric Septic Shock Algorithm

The recommended treatment approach to restore hemodynamic stability for septic shock in children is presented in the [Pediatric Septic Shock Algorithm](#) (Figure 41). It outlines 3 phases requiring rapid recognition, stabilization/resuscitation, and critical care management, as follows:

- **Early detection of signs of septic shock**, which may include alteration in mental status, heart rate, temperature, and perfusion. Note that systolic or diastolic hypotension may or may not be present.
- **Initial stabilization/resuscitation:**
 - Within 10 to 15 minutes after recognizing signs of shock, monitor and support airway, oxygenation, and circulation, monitor heart rate and pulse oximetry, and establish vascular access (IV/O). Initiate fluid resuscitation and give 10-20 mL/kg of isotonic crystalloid bolus over 5 to 20 minutes.
 - Reassess child during and after each fluid bolus. Stop rapid fluid bolus administration if rales, respiratory distress, or hepatomegaly develops. If additional boluses are needed to treat shock, it may be necessary to administer smaller boluses over longer intervals.
 - Within the first hour after onset of shock, administer antibiotics, give fluid boluses (repeating, as needed, to correct shock), and initiate vasoactive drug therapy if shock persists despite administration of fluid boluses.
 - Critical care: If signs of shock persist, critical care expertise is needed to establish hemodynamic monitoring, continue initial vasoactive drug therapy, and titrate additional drug therapy, as needed, to treat shock.



Once shock has been effectively treated, further management includes continued monitoring and support of organ function, treatment of the source of the infection, and evaluation of the child's clinical course and therapy.

Identify signs of septic shock (Step 1).

- Altered mental status
- Altered heart rate
- Altered temperature
- Altered perfusion
- Hypotension

Initial stabilization/resuscitation for septic shock (Step 2)

Adequate treatment during the first hour after the onset of symptoms is critical to maximize survival for a child in septic shock. Immediate (10-15 minutes) components of management of septic shock are

- Support adequate airway, breathing, and circulation.
- Monitor heart rate, blood pressure, and pulse oximetry.
 - Establish IV/IO access.
 - Give boluses of 10 to 20 mL/kg (10 mL/kg for neonates and those with preexisting heart disease) of isotonic crystalloid.

Within the first hour (Step 3)

- Draw labs (if possible, draw a blood sample for culture before antibiotic administration, but do not delay antibiotics).
- Administer antibiotics within 1 hour after first medical contact.
- Assess carefully after each fluid bolus; stop fluid boluses if rales, respiratory distress, or hepatomegaly develops.
- Give antipyretics if needed.

Do signs of shock persist? (Step 4)

If signs of shock do not persist, consider critical care consultation (Step 5).

If signs of shock persist, go to Step 6:

- Obtain critical care consultation.
- Initiate and titrate epinephrine or norepinephrine.

Critical care therapies (Step 7)

- Establish central venous and intra-arterial pressure monitoring.
- Continue epinephrine/norepinephrine, and fluid bolus therapy as needed to treat shock.
- Verify adequate airway, oxygenation, and ventilation.
- Consider stress-dose hydrocortisone if hemodynamics remain inadequate despite adequate fluid resuscitation and vasoactive drug therapy.

Rapid, appropriate fluid bolus administration is a priority. Inadequate intravascular volume rapidly leads to low stroke volume and hypotension. A child in septic shock often requires a large volume of fluid to restore perfusion. It may be necessary to rapidly infuse 10 to 20 mL/kg of isotonic crystalloid solution (up to 40-60 mL/kg total). Titrate the volume and rate of fluid administration during and after each bolus by assessing the child's mental status, heart rate, temperature, blood pressure, and organ perfusion (including presence and quality of peripheral pulses, capillary refill, skin temperature and color, and urine output).

The spirit of the recommendations about fluid therapy for treating septic shock is a continued emphasis on fluid resuscitation for both compensated (detected by physical examination) and decompensated (hypotensive) septic shock. Moreover, emphasis is also placed on the use of individualized patient evaluation before the administration of IV fluid boluses, including physical examination by a clinician and frequent reassessment to determine the appropriate volume of fluid resuscitation. The clinician should also integrate clinical signs with patient and locality-specific information about prevalent diseases, vulnerabilities (such as severe anemia and malnutrition), and available critical care resources.

When caring for children with severe febrile illness (such as those included in the FEAST trial) in settings with limited access to critical care resources (ie, mechanical ventilation and inotropic support), use extreme caution when administering bolus intravenous fluids because it may be harmful (refer to Maitland et al. *N Engl J Med*. 2011;364[26]:2483-2495 for details).

The other scenario in which you should exercise caution with fluid boluses is in patients with evidence of fluid overload. Signs and symptoms of fluid overload include hepatomegaly as well as increased respiratory distress and pulmonary edema, such as rales/crackles, which may result from increased vascular permeability. If such signs develop, stop rapid bolus fluid administration. The child requires supplemental O₂ and may require endotracheal intubation and mechanical ventilation with positive end-expiratory pressure. If additional fluid boluses are needed, give smaller volumes more slowly (eg, 5-10 mL/kg over 10-20 minutes).

Pulmonary edema may also develop with cardiogenic shock, which often manifests because of hepatomegaly, cardiomegaly, and poor myocardial contractility. In both instances, you may need to reduce the volume and rate of

fluid administration.

Draw blood samples for blood culture. Administer the first dose of broad-spectrum antibiotics within 1 hour after first medical contact. Do not delay antimicrobial therapy to wait for blood cultures or to perform other diagnostic assessments such as lumbar puncture.

Anticipate the possible need for vasopressors and stress-dose hydrocortisone. Order these drugs from the pharmacy early so that they will be at the bedside. They should be immediately available if the shock is fluid refractory or if adrenal insufficiency is suspected. Vasopressors can be administered via peripheral venous or IO access, so initiation of vasopressor therapy should not await transfer to a critical care unit. Because delaying initiation of vasoactive support has been associated with increased length of stay and mortality in pediatric septic shock, it is important to begin vasoactive drug therapy within the first hour if shock persists despite bolus fluid administration.

Identify and correct metabolic derangements immediately. Hypoglycemia and ionized hypocalcemia are commonly seen in septic shock and may contribute to myocardial dysfunction.

Use diagnostic assessments (eg, lactate concentration, base deficit, ScvO₂, echocardiogram, point-of-care ultrasound) to identify the severity of shock and monitor the response to fluid therapy.

Managing Fluid-Refractory Septic Shock

Vasoactive medications are appropriate to initiate after 40 to 60 mL/kg of fluid has been administered and the patient continues to have signs/symptoms of compromised perfusion. Vasoactive medications may be appropriate to administer before this fluid threshold if there is evidence of or concern about fluid overload (eg, rales, hepatomegaly).

- Administer vasoactive therapy to improve tissue perfusion and blood pressure.
- Either epinephrine or norepinephrine are appropriate as the first vasoactive agent being initiated; there is no evidence to support one over the other based on the patient being in cold shock vs warm shock.
- Despite the lack of evidence, the majority of the expert authors of the 2020 Surviving Sepsis Campaign International Guidelines preferred epinephrine as the initial vasoactive agent in patients with low cardiac output, myocardial dysfunction, or both. For patients in septic shock with low SVR, norepinephrine was the initial vasoactive agent of choice among this panel of experts.
- Both epinephrine and norepinephrine are preferable to dopamine in pediatric septic shock.
- If the patient continues with hypotension, inadequate perfusion, or both, despite high-dose catecholamine infusion(s), consider initiating a vasopressin infusion.
- Typically, epinephrine is administered to the child who has fluid-refractory septic shock and cold extremities.
- Administer additional fluid boluses of 10 to 20 mL/kg isotonic crystalloid as deemed clinically appropriate on the basis of the patient's clinical assessment.
- Consider endotracheal intubation and early assisted ventilation with supplemental O₂ and positive end-expiratory pressure as needed.

After initial stabilization/resuscitation, including administration of antibiotics, fluid boluses, and vasoactive therapy within the first hour of care, evaluate heart rate, blood pressure, and peripheral perfusion to determine the next intervention. If heart rate, blood pressure, and perfusion are returning to normal, arrange admission or transfer to an appropriate pediatric facility. Initiate consultation with a pediatric critical care unit or transport team. If the child remains hypotensive or poorly perfused, proceed to the next level in the algorithm. Continue fluid and vasoactive therapy while preparing for admission or transfer.

Critical Care of Septic Shock

Once critical care expertise is available for the child with persistent shock despite initial fluid bolus and vasoactive drug therapy, additional hemodynamic monitoring can be established and additional vasoactive drugs and fluid boluses can be provided.

- Establish central venous and intra-arterial pressure monitoring.
- Assess the child's heart rate, blood pressure, and perfusion and consider advanced hemodynamic monitoring, including invasive arterial blood pressure monitoring, central venous pressure monitoring, and central venous oxygen saturation.
- Analyze laboratory data (eg, serum glucose, calcium, lactate).
- Continue epinephrine/norepinephrine and add additional vasoactive drugs as needed.
- Provide fluid boluses as needed to treat shock; assess the child frequently and after each fluid bolus.

- Consider stress-dose hydrocortisone (particularly if the child has been on chronic systemic steroids). Cortisol levels are not routinely indicated or recommended.

It is not always clear from the physical examination whether a child has vasodilation or vasoconstriction. For example, some children with cool extremities may have vasodilation but be poorly perfused because of low stroke volume and poor cardiac function. The reasons for specific drug selection according to type of shock are described later in this manual.

While managing septic shock, it is important to support organ function, serum electrolytes, acid-base balance, and lactate, glucose, and calcium concentrations. In titrating drug and fluid therapy, goals of care are to improve the child's lactate, blood pressure, and organ perfusion.

Correction of Adrenal Insufficiency

A child in septic shock with fluid-refractory and epinephrine- or norepinephrine-resistant shock may have adrenal insufficiency. If you suspect adrenal insufficiency, or a patient is at known risk for adrenal insufficiency (ie, history of steroid use), give hydrocortisone 1 to 2 mg/kg IV bolus early. If possible, obtain a baseline cortisol concentration before administration, and obtain expert consultation for any additional evaluation and management.

Therapeutic End Points

Titrate vasoactive agents in septic shock to therapeutic end points, including

- Improving lactate concentration
- Normalized heart rate (declining from very rapid rate toward normal)
- Adequate blood pressure
- Good distal pulses and perfusion with capillary refill 2 seconds or less
- Improved level of consciousness/responsiveness
- Appropriate urine output

Strict adherence to end points is recommended to avoid excessive vasoconstriction in key organs.

Poststabilization Care

Once signs of shock are resolved, the child requires continued monitoring and support of organ function. The source infection must be treated appropriately. In addition, the healthcare team should evaluate the cause of the infection to determine if it could have been prevented (and how) and the speed of the recognition of the septic shock (could it have been improved), and identify any potential improvements that could be made to increase the efficiency and effectiveness of shock resuscitation for future patients with septic shock.

Managing Anaphylactic Shock

Managing anaphylactic shock focuses on treating life-threatening cardiopulmonary problems and reversing or blocking the mediators released as part of the uncontrolled allergic response. Primary therapy is administration of epinephrine to reverse the effects of histamine and other allergic mediators. The epinephrine dose may be repeated if needed. Angioedema (tissue swelling resulting from a marked increase in capillary permeability) may result in complete upper airway obstruction, so providers should anticipate the need for very early airway intervention with assisted ventilation and intubation. The epinephrine may prevent or reverse hypotension, and fluid resuscitation may also be effective in restoring blood pressure and supporting effective perfusion.

Specific Treatment Considerations

Consider the initial general management of shock outlined in the [Summary: Initial Management Principles](#) section in addition to the following specific treatments for anaphylactic shock:

- Place the patient supine, administer oxygen, and maintain airway.
- Epinephrine: first-line treatment
 - IM epinephrine or epinephrine by autoinjector (pediatric or adult, depending on the child's size) is the most important agent for the treatment of anaphylaxis.
 - A second dose or an epinephrine infusion may be needed after 10-15 minutes in severe anaphylaxis, frequently a low-dose infusion (<0.05 mcg/kg per minute) is effective.
- Administer isotonic crystalloid fluid boluses as needed to support circulation.
- Albuterol

- Administer albuterol as needed for bronchospasm by metered dose inhaler, intermittent nebulizer, or continuous nebulizer.

Adjunct Treatments

- Antihistamines
 - H₁ blocker (ie, diphenhydramine)
 - Consider an H₂ blocker (eg, famotidine)
- Corticosteroids
 - Methylprednisolone or equivalent corticosteroid

Observation is indicated for identification and treatment of late-phase symptoms. Late-phase symptoms may occur in 25% to 30% of children several hours after acute-phase symptoms. The likelihood of late-phase symptoms increases in proportion to the severity of acute-phase symptoms.

Managing Neurogenic Shock

Children with neurogenic shock typically present with hypotension, bradycardia, and sometimes hypothermia. Minimal response to fluid resuscitation is commonly observed. Blood pressure is characterized by a low diastolic blood pressure with a wide pulse pressure because of loss of vascular tone. Children with spinal shock may be more sensitive to variations in ambient temperature and may require supplemental warming or cooling.

Specific Treatment Considerations

The initial management principles for shock outlined in the Summary: Initial Management Principles section may be considered in addition to the following specific treatments for neurogenic shock:

- Position the child flat or head down to improve venous return.
- Administer a trial of fluid therapy (isotonic crystalloid) and assess response.
- *For fluid-refractory hypotension*, use vasopressors (eg, norepinephrine, epinephrine) as indicated.
- Provide supplemental warming or cooling as needed.

Managing Cardiogenic Shock

Cardiogenic shock is a condition of inadequate tissue perfusion that results from myocardial dysfunction. Initially, cardiogenic shock may resemble hypovolemic shock, so identifying a cardiogenic etiology may be difficult. If you suspect cardiogenic shock, consider a slow administration (10-20 minutes) of a relatively small fluid bolus (ie, 5-10 mL/kg) while carefully monitoring the child for response. Cardiogenic shock is probably present if the child does not improve, the child's respiratory function deteriorates, or the child develops signs of pulmonary edema. Evidence of venous congestion (eg, elevated central venous pressure, distended jugular veins, or hepatomegaly) and cardiomegaly (on chest x-ray) are also suggestive of a cardiac etiology of shock.

Main Objectives

A main objective in managing cardiogenic shock is to improve the effectiveness of cardiac function and cardiac output by increasing the efficiency of ventricular ejection. Another main objective is to minimize metabolic demand.

Many children with cardiogenic shock have high preload and do not require additional fluid therapy. Others may require a cautious fluid bolus to increase preload. The most effective way to increase stroke volume in these patients is to reduce afterload (SVR) rather than give an inotropic agent. Inotropes may increase cardiac contractility but will also increase myocardial O₂ demand. However, children who are already hypotensive may require fluid therapy and inotropic support before they will tolerate afterload reduction. Specific management includes

- Cautious fluid administration and monitoring
- Laboratory and other diagnostic studies
- Medications
- Mechanical circulatory support

Consult a pediatric critical care or pediatric cardiology specialist at the earliest opportunity. This will help to establish a diagnosis (eg, echocardiogram), guide ongoing therapy, and facilitate direct transfer to definitive care.

Cautious Fluid Administration and Monitoring

Large heart size on the chest x-ray in a child with evidence of shock and poor cardiac output is the hallmark of cardiogenic shock with adequate intravascular volume. Obtain an echocardiogram for more objective and accurate data about preload and cardiac function.

Do not give large or rapid fluid boluses to a child with cardiogenic shock. The boluses can worsen heart function and increase fluid in the lungs. If objective data or the child's history (eg, vomiting and poor intake) are consistent with inadequate preload, you may give a small fluid bolus cautiously. Remember the following:

- Give small isotonic fluid boluses (5-10 mL/kg)
- Give fluid boluses over relatively longer periods of time (eg, 10-20 minutes instead of 5-10 minutes)
- Monitor the child closely during fluid infusion
 - Assess for respiratory function frequently
 - Watch for development of pulmonary edema and deterioration in pulmonary function

Give supplemental O₂. Be prepared to provide assisted ventilation. Noninvasive positive pressure support may reduce the need for mechanical ventilation by decreasing the work of breathing and improving oxygenation.

Consider establishing central venous access to facilitate measurement of central venous pressure as an index of preload status and to assess right ventricular end-diastolic pressure and to provide access for multiple infusions. Central venous access also allows monitoring of ScvO₂ as an objective measurement of the adequacy of O₂ delivery relative to metabolic demand. Invasive monitoring with a pulmonary artery catheter, an option in the pediatric intensive care unit, is not critical to diagnosing cardiogenic shock.

Laboratory and Other Diagnostic Studies

Obtain laboratory studies to assess the impact of shock on end-organ function. No single laboratory study is completely sensitive or specific for cardiogenic shock. Appropriate studies often include

- An arterial blood gas to determine the magnitude of metabolic acidosis and adequacy of oxygenation and ventilation
- Hemoglobin concentration to ensure that O₂-carrying capacity is adequate
- Lactate concentration and ScvO₂ as indicators of the adequacy of O₂ delivery relative to metabolic demand
- Cardiac enzymes (eg, creatine kinase, myocardial, troponin) and thyroid function tests

Other useful studies include the following:

- Chest x-ray: provides information about cardiac size, pulmonary vascular markings, pulmonary edema, and coexistent pulmonary pathology
- Electrocardiogram: may detect arrhythmia, myocardial injury, ischemic heart disease, or evidence of drug toxicity
- Echocardiogram: may be diagnostic, revealing congenital heart disease, akinetic or dyskinetic ventricular wall motion or valvular dysfunction; also provides objective measurement of ventricular chamber volume (ie, preload) and function

Medications

If the child is normotensive, medication therapy consists of diuretics and vasodilators or inodilators. Diuretics are indicated when the child has evidence of pulmonary edema or systemic venous congestion. Vasodilators and inodilators are typically given by continuous infusion.

Children with cardiogenic shock may require medications to increase cardiac output by improving contractility. Most also require agents to reduce peripheral vascular resistance. This includes vasodilators, inotropes, and phosphodiesterase enzyme inhibitors (ie, inodilators). Milrinone is the preferred drug in many centers. For a detailed discussion of these agents, refer to Medication Therapy in this Part.

Increased metabolic demand, particularly increased myocardial O₂ demand, plays a role in the vicious cycle of cardiogenic shock. Reducing metabolic demand is a critical component in managing cardiogenic shock. Use ventilatory support and antipyretics to reduce metabolic demand. Analgesics and sedatives reduce O₂ consumption but also reduce the endogenous stress response that is helping redistribute blood flow to compensate for low cardiac output. Give these agents in small doses. Monitor the child carefully for evidence of potential respiratory depression or hypotension.

Mechanical Circulatory Support

Children with cardiogenic shock who do not respond to medical management may benefit from mechanical circulatory support if the cause of shock is potentially reversible. Extracorporeal life support can provide temporary maintenance of cardiac output, oxygenation, and ventilation while the underlying cause of cardiopulmonary failure is treated. Forms of extracorporeal life support include extracorporeal membrane oxygenation and ventricular assist devices. One specific etiology of cardiogenic shock for which extracorporeal membrane oxygenation may be considered is acute fulminant myocarditis with high risk of imminent cardiac arrest. Extracorporeal life support is usually available only in tertiary pediatric centers that have the resources and expertise to manage children with acute cardiopulmonary failure.

Specific Treatment Considerations

Follow the initial management principles for shock outlined in the [Summary: Initial Management Principles](#) section in addition to the following considerations specific to cardiogenic shock:

- Administer supplemental O₂ and consider need for noninvasive positive pressure or mechanical ventilation.
- Give 5 to 10 mL/kg isotonic crystalloid infusion slowly (over 10-20 minutes); repeat PRN.
- Assess frequently for pulmonary edema.
- Be prepared to assist ventilation.
- Obtain expert consultation early.
- Order laboratory and other studies to determine underlying cause and degree of cardiac and end-organ dysfunction.
- Administer pharmacologic support (eg, vasodilators, phosphodiesterase enzyme inhibitors, inotropes, analgesics, antipyretics).
- Consider mechanical circulatory support.

Managing Obstructive Shock

Managing obstructive shock is specific to the type of obstruction. This section discusses managing

- Cardiac tamponade
- Tension pneumothorax
- Ductal-dependent congenital heart lesions
- Massive pulmonary embolism

Main Objectives

The early clinical presentation of obstructive shock may resemble hypovolemic shock. A reasonable initial approach may include administering a fluid challenge (10-20 mL/kg isotonic crystalloid). Because children with obstructive shock can rapidly progress to cardiopulmonary failure and then cardiac arrest, immediately identifying obstructive shock by using the secondary assessment and diagnostic assessments is critical to effective treatment. The main objectives in managing obstructive shock are to correct the cause of cardiac output obstruction and restore tissue perfusion.

General Management Principles

In addition to considerations specific to the etiology of the obstruction, follow the initial management principles outlined in the section [Fundamentals of Shock Management](#).

Specific Management of Cardiac Tamponade

Cardiac tamponade is caused by accumulating fluid, blood, or air in the pericardial space, limiting systemic venous return, impairing ventricular filling, and reducing cardiac output. Favorable outcome requires rapidly identifying and immediately treating tamponade. Children with cardiac tamponade may improve temporarily with fluid administration to augment cardiac output and tissue perfusion until pericardial drainage can be performed.

Consult appropriate specialists (eg, pediatric critical care, pediatric cardiology, pediatric surgery) early. Elective pericardial drainage (pericardiocentesis), often guided by echocardiography or fluoroscopy, should be performed by specialists who are trained and skilled in the procedure. Emergency pericardiocentesis may be performed in the setting of impending or actual pulseless arrest when there is a strong suspicion of pericardial tamponade.

Specific Management of Tension Pneumothorax

Tension pneumothorax is characterized by accumulating air under pressure in the pleural space with compression of the lung on the involved side and mediastinal shift to the other side of the chest. This prevents the lungs from

expanding properly and applies pressure on the heart and great veins, compromising venous return and cardiac output. Favorable outcome depends on immediate diagnosis and urgent treatment. A tension pneumothorax should be identified through clinical examination, and treatment should not await confirmation with a chest radiograph.

Treatment of a tension pneumothorax is immediate needle decompression followed by thoracostomy for chest tube placement as soon as possible. A trained provider can quickly perform an emergency needle decompression by inserting an 18- to 20-gauge over-the-needle catheter over the top of the child's third rib (second intercostal space) in the midclavicular line. A gush of air is a sign that needle decompression has been successful. This indicates relief of pressure buildup in the pleural space.

Specific Management of Ductal-Dependent Lesions

Ductal-dependent lesions are a group of congenital cardiac abnormalities. These abnormalities result in pulmonary or systemic blood flow that must pass through an open/patent ductus arteriosus. The infant can deteriorate rapidly once the ductus begins to close during the first days or weeks of life.

Congenital heart defects with ductal-dependent pulmonary blood flow usually include a severe obstruction to pulmonary blood flow from the right ventricle, so all pulmonary blood flow then must pass from the aorta through the ductus arteriosus and into the lungs. When the ductus begins to close, the infant initially becomes hypoxemic and cyanotic, which can progress to shock if not recognized and intervened upon early.

Congenital heart defects with ductal-dependent systemic blood flow usually consist of an obstruction to outflow through or from the left side of the heart into the aorta. In such patients, systemic blood flow must pass from the right ventricle and pulmonary artery through the ductus into the aorta. In these patients, when the ductus starts to close, signs of shock develop with severe deterioration in systemic perfusion.

For any infant with ductal-dependent pulmonary or systemic blood flow, immediate treatment with continuous infusion of prostaglandin E₁ (PGE₁) can restore ductal openness/patency and may be lifesaving.

Other management actions for ductal-dependent obstructive lesions are

- Ventilatory support with O₂ administration
- Expert consultation to direct therapy
- Echocardiography to establish diagnosis and guide therapy
- Administration of inotropic agents to improve myocardial contractility
- Judicious administration of fluids to improve cardiac output
- Correction of metabolic derangements, including metabolic acidosis

Specific Management of Massive Pulmonary Embolism

Massive pulmonary embolism is a sudden block in the main or large-branch pulmonary artery. This block is usually caused by a blood clot that has traveled to the lungs from another part of the body. The block also can result from other substances, including fat, air, amniotic fluid, catheter fragment, or injected matter. Blood flow through the pulmonary circulation to the left side of the heart is obstructed, resulting in decreased left ventricular filling and inadequate cardiac output.

Initial treatment is supportive, including O₂ administration, ventilatory assistance, and fluid therapy if the child is poorly perfused. Consult a specialist who can perform echocardiography, a computed tomography scan with IV contrast, or angiography to confirm the diagnosis. Anticoagulants (eg, heparin, enoxaparin) are the definitive treatment for most children with pulmonary embolism who are not in shock. Because anticoagulants do not act immediately to relieve obstruction, consider fibrinolytic agents (eg, recombinant tissue plasminogen activator, systemically or by image-guided direct thrombolysis) in children with severe cardiovascular compromise.

Computed tomography angiography is the diagnostic assessment of choice because it can be rapidly obtained and does not require an invasive angiogram. Additional diagnostic studies that might be useful are an arterial blood gas, CBC, D-dimer, electrocardiography, chest x-ray, ventilation-perfusion scan, and echocardiography.

Managing Shock Flowchart

Table 55 summarizes general management of shock and specific management by etiology.

Table 55. Managing Shock Flowchart

Managing shock flowchart	

<ul style="list-style-type: none">• Oxygen• Pulse oximetry• ECG monitor	<ul style="list-style-type: none">• IV/IO access• BLS as indicated• Point-of-care glucose testing		
Hypovolemic shock: Specific management for selected conditions			
Nonhemorrhagic		Hemorrhagic	
<ul style="list-style-type: none">• 20 mL/kg NS/LR bolus, repeat as needed• Consider colloid		<ul style="list-style-type: none">• Control external bleeding• 20 mL/kg NS/LR bolus, repeat 2 or 3x as needed• Transfuse PRBCs as indicated	
Distributive shock: Specific management for selected conditions			
Septic	Anaphylactic		Neurogenic
Management algorithm: <ul style="list-style-type: none">• Septic Shock	<ul style="list-style-type: none">• IM epinephrine (or autoinjector)• Fluid boluses (10-20 mL/kg NS/LR)• Albuterol• Antihistamines, corticosteroids• Epinephrine infusion		<ul style="list-style-type: none">• 20 mL/kg NS/LR bolus, repeat PRN• Vasopressor
Cardiogenic shock: Specific management for selected conditions			
Bradyarrhythmia/tachyarrhythmia		Other (eg, CHD, myocarditis, cardiomyopathy, poisoning)	
Management algorithms: <ul style="list-style-type: none">• Bradycardia• Tachycardia		<ul style="list-style-type: none">• 5 to 10 mL/kg NS/LR bolus, repeat PRN• Inotropic and/or vasoactive infusion• Consider expert consultation• Antidote for poisoning	
Obstructive shock: Specific management for selected conditions			
Ductal-dependent (LV outflow obstruction)	Tension pneumothorax	Cardiac tamponade	Pulmonary embolism
<ul style="list-style-type: none">• Prostaglandin E1• Expert consultation	<ul style="list-style-type: none">• Needle decompression• Tube thoracostomy	<ul style="list-style-type: none">• Pericardiocentesis• 20 mL/kg NS/LR bolus	<ul style="list-style-type: none">• 20 mL/kg NS/LR bolus, repeat PRN• Consider thrombolytics, anticoagulants• Expert consultation

Resources for Managing Circulatory Emergencies

Intraosseous Access

Intraosseous (IO) cannulation is a relatively simple and effective method of rapidly establishing vascular access for emergency fluids or medications. It provides access to a noncollapsible marrow venous plexus, serving as a safe and reliable route for administering drugs, crystalloids, colloids, and blood during resuscitation. One can achieve IO access in children of all ages, often in about 30 to 60 seconds. In certain circumstances (eg, cardiac arrest or severe shock with severe vasoconstriction), it may be the initial vascular access attempted. Any medication that can be administered IV can be given by the IO route. Fluids and medications delivered via an IO catheter can reach the central circulation within seconds. If you cannot readily obtain peripheral vascular access in a child with compensated or hypotensive shock, be prepared to establish IO access as soon as it is needed.

Sites for IO Access

Many sites, such as the proximal tibia (just below the growth plate), are appropriate for IO infusion. Other sites include the distal tibia (just above the medial malleolus), the distal femur, and the anterior-superior iliac spine. Newer devices, such as the IO drill, are approved for use in the proximal humerus in older children, adolescents, and adults.

Contraindications

Contraindications to IO access include

- Fractures and crush injuries near the access site
- Conditions with fragile bones (eg, osteogenesis imperfecta)
- Previous attempts to establish IO access in the same bone

Avoid IO cannulation if an infection is present in the overlying tissues.

Procedure (Proximal Tibia)

Use the following procedure to establish IO access:

1. To establish access in the proximal tibia, position the leg with slight external rotation. Identify the tibial tuberosity just below the knee joint; the insertion site is the flat part of the tibia, about 1 to 3 cm (about 1 finger's width) below and medial to this bony prominence ([Figure 42](#)). Always use universal precautions when attempting vascular access. Disinfect the overlying skin and surrounding area.
2. Leave the stylet in the needle during insertion to prevent the needle from becoming clogged with bone or tissue. Stabilize the leg on a firm surface; do not place your hand behind the leg. *Note:* If a standard IO needle or bone marrow needle is not available, a large-bore (at least 18-gauge) standard hypodermic needle can be substituted. But the lumen may become clogged with bone or bone marrow during insertion. Short, wide-gauge spinal needles with internal stylets can be used in an emergency, but they tend to bend easily. Use a hemostat to help stabilize the needle during insertion.
3. Insert the needle through the skin over the anteromedial surface of the tibia perpendicular to the tibia; this avoids injury to the growth plate. Use a twisting motion with gentle but firm pressure. Continue inserting the needle through the cortical bone until there is a sudden decrease in resistance as the needle enters the marrow space. If the needle is placed correctly, it should stand easily without support and should not wiggle with gentle prodding.
4. Remove the stylet and attach a syringe. Aspiration of bone marrow and blood into the hub of the needle confirms correct placement; you may send blood to the lab for analysis. (*Note:* Blood or bone marrow may not always be aspirated despite correct needle placement.) Infuse a small volume of saline. It should infuse easily. Check for swelling at the insertion site or posteriorly, behind the insertion site. (Swelling occurs if the needle was inserted too deeply and penetrated through the posterior cortical bone.)
5. To stabilize the needle, place tape over the flange; you also may place gauze padding on both sides of the needle for support.
6. Tape IV tubing to the skin to avoid tension on the tubing that might displace the needle.
7. Infuse fluid with a syringe attached to a 3-way stopcock or by pressure infusion. When using a pressurized fluid bag, make sure no air gets into the system.
8. Any medication that can be administered IV can be given by the IO route, including vasoactive drug infusions such as epinephrine. Follow all bolus medications with a saline flush.




Figure 42A. Locations for intraosseous insertion. A, General landmarks for IO insertion in the leg of an infant.




Figure 42B. Locations for IO insertion in the proximal

tibia and distal femur in older children.




Figure 42C. Location for IO insertion in the iliac crest.




Figure 42D. Location for IO insertion in the distal tibia.




Figure 42E. Technique for immobilizing the leg while twisting the IO needle into the leg of an infant.

After IO Insertion

After inserting an IO needle/catheter, do the following:

- Check the site and the underside of the leg for signs of swelling and IO needle displacement; fluids or drugs delivered via a displaced needle can cause severe complications (eg, tissue necrosis, compartment syndrome).
- Because IO needles are intended for short-term use (generally less than 24 hours) and are often displaced early in a resuscitation, prioritize ongoing attempts to establish vascular access.

Color-Coded Length-Based Resuscitation Tape

Use a color-coded length-based tape ([Table 56](#)) to select the correct size of resuscitation supplies and to determine the child's weight (if not known) for calculating drug doses.

Table 56. Color-Coded Length-Based Resuscitation Tape

Zone	3 kg	4 kg	5 kg	Pink	Red	Purple	Yellow	White	Blue	Orange	Green
ETT uncuffed (mm)	3.5	3.5	3.5	3.5	3.5	4.0	4.5	5.0	5.5	N/A	N/A
ETT cuffed (mm)	3.0	3.0	3.0	3.0	3.0	3.5	4.0	4.5	5.0	5.5	6.0
Lip-tip (cm)	9-9.5	9.5-10	10-10.5	10-10.5	10.5-11	11-12	12.5-13.5	14-15	15.5-16.5	17-18	18.5-19.5
Suction (F)	8	8	8	8	8	8	10	10	10	10	12
L-scope blade	1 straight	1 straight	1 straight	1 straight	1 straight	1-1.5 straight	2 straight/curved	2 straight/curved	2 straight/curved	2-3 straight/curved	2-3 straight/curved
Stylet	6 F	6 F	6 F	6 F	6 F	6 F	10 F	10 F	10 F	14 F	14 F
OPA (mm)	50	50	50	50	50	60	60	60	70	80	80
NPA (F)	14	14	14	14	14	18	20	22	24	26	26
Bag-mask device (minimum mL)	450	450	450	450	450	450	450	450-750	750-1000	750-1000	1000
ETCO₂ detector	Ped	Ped	Ped	Ped	Ped	Ped	Ped	Adult	Adult	Adult	Adult
LMA	1	1	1	1.5	1.5	2	2	2	2-2.5	2.5	3
Tidal volume (mL)	20-30	24-40	30-50	40-65	50-85	65-105	80-130	100-165	125-210	160-265	200-330
Frequency	20-25/min	20-25/min	20-25/min	20-25/min	20-25/min	15-25/min	15-25/min	15-25/min	12-20/min	12-20/min	12-20/min

Abbreviations: ETT, endotracheal tube; F, French; LMA, laryngeal mask airway; NPA, nasopharyngeal airway; OPA, oropharyngeal airway; Ped, pediatric.

The Broselow-Luten System Point of Care Guide is © 2020 Vyaire Medical, Inc.; used with permission.

Part 11: Recognizing Arrhythmias

This Part discusses recognizing arrhythmias, such as bradycardia (slow heart rate) and tachycardia (fast heart rate) with a palpable pulse and adequate or inadequate perfusion in infants and children.

Learning Objective

After completing this Part, you should be able to differentiate between unstable and stable patients with arrhythmias.

You will be expected to identify bradycardic and tachycardic rhythms in an infant or child with and without cardiopulmonary compromise.

Bradycardia Definitions

Bradycardia is a heart rate that is slow in comparison with a normal heart rate range for the child's age, level of activity, and clinical condition. Refer to [Table 14: Normal Heart Rates](#) in [Part 4](#).



Critical Concepts: Symptomatic Bradycardia and Cardiopulmonary Compromise

Symptomatic bradycardia is a heart rate slower than normal for the child's age (usually less than 60/min) associated with cardiopulmonary compromise.

Cardiopulmonary compromise is defined as hypotension, acutely altered mental status (ie, decreased level of consciousness), and signs of shock.

Bradycardia is an ominous sign of impending cardiac arrest in infants and children, especially if it is associated with hypotension or evidence of poor tissue perfusion. If, despite adequate oxygenation and ventilation, the heart rate is less than 60/min in an infant or child with signs of cardiopulmonary compromise, begin CPR.



Critical Concepts: Evaluating Heart Rate and Rhythm

Consider the following when evaluating the heart rate and rhythm in any seriously ill or injured child:

- The child's typical heart rate and baseline rhythm
- The child's activity level and clinical condition (including baseline cardiac function)

Children with congenital heart disease may have underlying conduction abnormalities. Interpret the child's heart rate and rhythm by comparing them to the child's baseline heart rate and rhythm. Children with poor baseline cardiac function are more likely to become symptomatic from arrhythmias than children with normal cardiac function.

Tissue hypoxia is the leading cause of symptomatic bradycardia in children. Therefore, symptomatic bradycardia in children is usually the result of (rather than the reason for) progressive hypoxemia and respiratory failure or shock. Priorities in initial assessment and management should be to support the airway and adequately oxygenate and ventilate.

Bradycardia may be classified as

- Primary bradycardia
- Secondary bradycardia

Primary bradycardia is the result of congenital or acquired heart conditions that slow the spontaneous depolarization rate of the heart's normal pacemaker cells or slow conduction through the heart's conduction system. Causes of primary bradycardia include

- Congenital abnormality of the heart pacemaker or conduction system
- Surgical injury to the pacemaker or conduction system

- Cardiomyopathy
- Myocarditis

Secondary bradycardia is the result of noncardiac conditions that alter the normal function of the heart (ie, slow the sinus node pacemaker or slow conduction through the atrioventricular [AV] junction). Causes of secondary bradycardia include

- Hypoxia
- Acidosis
- Hypotension
- Hypothermia
- Drug effects
- Increased vagal tone (suctioning, gagging, vomiting)

Recognizing Bradycardia

Signs and Symptoms of Bradycardia

Cardiac output (the volume of blood pumped by the heart per minute) is the product of stroke volume (the volume of blood pumped with each ventricular contraction) and heart rate (number of times the ventricles contract per minute):

Cardiac Output = Stroke Volume × Heart Rate

When heart rate decreases, cardiac output can only be maintained by an increase in stroke volume. Because the heart's ability to increase stroke volume is limited during infancy and early childhood, cardiac output typically declines with bradycardia. An extremely slow heart rate results in critically low cardiac output that can be life threatening and lead to cardiopulmonary compromise. The signs of cardiopulmonary compromise associated with bradycardia are

- Hypotension
- Altered mental status: decreased level of consciousness or irritability
- Shock: poor end-organ perfusion with or without hypotension

Additional signs can include

- Respiratory distress or failure
- Chest pain or vague feeling of discomfort in older children
- Sudden collapse

ECG Characteristics of Bradycardia

The electrocardiographic (ECG) characteristics of bradycardia include

- Heart rate: slow compared with normal heart rate for age
- P waves: may or may not be visible
- QRS complex: narrow or wide (depending on the origin of the rhythm and/or location of injury to the conduction system)
- P wave and QRS complex: may be unrelated (ie, AV dissociation)

Refer to [Rhythm Recognition Review](#) in the [Appendix](#) for examples.

Types of Bradyarrhythmias

Bradycardia, defined earlier in this Part, is a nonspecific term. If a patient has bradycardia due to slowing of the sinus node depolarization rate, the term *sinus bradycardia* is used.

Bradyarrhythmia is the appropriate term to use when the heart rate is abnormally low *and* the rhythm is abnormal. Types of bradyarrhythmias include AV blocks, sinus node arrest, junctional escape rhythms, and ventricular escape rhythms. Only AV blocks will be reviewed; the other bradyarrhythmias are more complex and are not discussed in the PALS Provider Course.

Sinus Bradycardia

Sinus bradycardia is a sinus node depolarization rate that is slower than normal for the child's age (refer to [Table 14: Normal Heart Rates](#) in [Part 4](#)). Sinus bradycardia is not necessarily problematic. It is often present in healthy children at rest when metabolic demands of the body are relatively low (eg, during sleep). Well-conditioned athletes often have sinus bradycardia because they have high stroke volume and increased vagal tone. However, sinus bradycardia can also develop in response to hypoxia, hypotension, and acidosis. As discussed above, it is often the result of progressive respiratory failure or shock and indicates impending cardiac arrest. Sinus bradycardia also may result from drug effects. Therefore, evaluating sinus bradycardia must always involve assessing the clinical status of the child.

Rarely, children have a primary bradycardia with an intrinsic disorder of the sinus node that impairs the sinus node to depolarize at an adequate rate. These children usually have a history of surgery for complex congenital heart disease. Additional causes of sinus node disorders include congenital abnormalities of the conduction system, cardiomyopathy, and myocarditis.

AV Block

AV block is a disturbance of electrical conduction through the AV node. AV block is classified as follows and in [Table 57](#):

- *First degree*: A prolonged PR interval representing slowed conduction through the AV node ([Figure 43A](#))
- *Second degree*: Block of some, but not all, atrial impulses before they reach the ventricles. This block can be further classified as Mobitz type I or Mobitz type II second-degree AV block.
 - In Mobitz type I or Wenckebach-type second-degree AV block, there is successive prolongation of the PR interval preceding non-conducted P waves.
 - Mobitz type II is identified by intermittently nonconducted P waves, with a constant PR interval conducted on beats.
- *Third degree*: None of the atrial impulses conduct to the ventricles. This block may also be referred to as *complete heart block* or *complete AV block* ([Figure 43D](#)).

Table 57. AV Block Classification

Type	Causes	Characteristics	Symptoms
First degree	<ul style="list-style-type: none"> • <i>Note</i>: May be present in healthy children • Enhanced vagal tone • Myocarditis • Electrolyte disturbances (eg, hyperkalemia) • Hypoxemia • Myocardial infarction • Cardiac surgery • Drugs (eg, calcium channel blockers, β-adrenergic blockers, digoxin) • Acute rheumatic fever • Intrinsic AV nodal disease 	Prolonged PR interval with 1:1 P: QRS	Asymptomatic
Second-degree Mobitz type I (Wenckebach phenomenon)	<ul style="list-style-type: none"> • <i>Note</i>: May be present in healthy children • Drugs (eg, calcium channel and β-adrenergic blockers, digoxin) • Any condition that stimulates vagal (parasympathetic) tone • Myocardial infarction • Lyme disease 	In Mobitz type I or Wenckebach-type second-degree AV block there is successive prolongation of the PR interval preceding nonconducted P waves.	Occasionally, may cause presyncope (light-headedness)
Second-degree Mobitz type II	<ul style="list-style-type: none"> • Typically results from intrinsic conduction system abnormalities • Rarely caused by increased 	Mobitz type II is identified by intermittently nonconducted P waves with a constant PR interval on conducted beats.	May cause <ul style="list-style-type: none"> • Sensed irregularities of heartbeat (palpitations) • Presyncope (light-headedness) • Syncope

	parasympathetic tone or drugs <ul style="list-style-type: none"> • Cardiac surgery • Myocardial infarction • Lyme disease 		
Third degree	<ul style="list-style-type: none"> • Extensive conduction system disease or injury, including myocarditis • Cardiac surgery • Congenital complete heart block • Myocardial infarction • Can also result from increased parasympathetic tone, toxic drug effects, or severe hypoxia/acidosis • Lyme disease 	<ul style="list-style-type: none"> • No relationship between P waves and QRS complexes • No atrial impulses reach ventricles • Ventricular rhythm maintained by a slower pacemaker 	Most frequent symptoms are <ul style="list-style-type: none"> • Fatigue • Light-headedness • Syncope



Figure 43A. Examples of AV block. A, Sinus bradycardia with first-degree AV block.



Figure 43B. Second-degree AV block Mobitz type I (Wenckebach phenomenon).



Figure 43C. Second-degree AV block Mobitz type II.



Figure 43D. Third-degree AV block.

Tachyarrhythmias

Tachycardia is a heart rate that is fast compared with the normal heart rate for the child's age. Refer to [Table 14: Normal Heart Rates](#) in [Part 4](#). Sinus tachycardia is a normal response to anxiety, pain, fever, or other physiologic stressors.

Tachyarrhythmias are rapid abnormal rhythms originating either in the atria or the ventricles of the heart. Tachyarrhythmias can be tolerated without symptoms for a variable period of time, especially if cardiac function is good. However, tachyarrhythmias can also cause acute hemodynamic compromise, such as shock or deterioration to cardiac arrest. Rapid deterioration is more likely to occur if cardiac function is poor when the arrhythmia develops.

Recognizing Tachyarrhythmias

Signs and Symptoms

Tachyarrhythmias may cause nonspecific signs and symptoms that differ according to the child's age. Clinical findings may include palpitations, light-headedness, and syncope. In infants who are at home, the tachyarrhythmia may be undetected for long periods (eg, for hours or days) until cardiac output is significantly compromised and the infant develops signs of congestive heart failure (CHF) such as irritability, poor feeding, and rapid breathing. Episodes of extremely rapid heart rate may be life threatening if cardiac output is compromised. When a tachyarrhythmia develops in a child with poor baseline cardiovascular function, clinical signs can develop quickly and deterioration can be rapid.

Signs of hemodynamic instability associated with tachyarrhythmias are

- Hypotension
- Altered mental status (ie, decreased level of consciousness)
- Signs of shock (poor end-organ perfusion) with or without hypotension

Additional signs may include

- Sudden collapse with rapid, weak pulses
- Respiratory distress/failure

Effect on Cardiac Output

An increased heart rate can produce increased cardiac output, up to a point. If that point is exceeded (ie, the heart rate is extremely rapid), stroke volume decreases because diastole is shortened and there is insufficient time for the ventricles to fill during diastole. Cardiac output then decreases substantially. In addition, coronary perfusion (blood flow to the heart muscle) occurs chiefly during diastole; the decrease in duration of diastole that occurs with a very rapid heart rate reduces coronary perfusion. Finally, a fast heart rate increases myocardial O_2 demand. In infants, prolonged episodes of rapid heart rate (supraventricular tachycardia [SVT]) can cause myocardial dysfunction, leading to CHF. In any child, an extremely rapid heart rate can result in inadequate cardiac output and, ultimately, cardiogenic shock.

Classifying Tachycardia and Tachyarrhythmias

Tachycardia and tachyarrhythmias are classified according to the width of the QRS complex; the arrhythmias are divided into those with narrow (0.09 second or less) vs wide (greater than 0.09 second) QRS complexes. Sinus tachycardia, SVT, and atrial flutter are classified as narrow complex, and ventricular tachycardia and SVT with aberrant interventricular conduction are classified as wide complex.

Sinus Tachycardia

Sinus tachycardia (ST) is a sinus node depolarization rate faster than normal for the child's age. It typically develops in response to the body's need for increased cardiac output or O_2 delivery. ST is a normal physiologic response and is not considered an arrhythmia (Figure 44). In ST, the heart rate is not fixed but varies with activity and other factors (eg, the child's sleep/awake state, temperature) that influence O_2 demand.



Figure 44. Sinus tachycardia (heart rate 180/min) in a febrile 10-month-old infant.

Common causes of ST include exercise, pain, anxiety, tissue hypoxia, hypovolemia (hemorrhagic and nonhemorrhagic fluid loss), shock, fever, metabolic stress, injury, toxins, poisons, drugs, and anemia. Cardiac tamponade, tension pneumothorax, and thromboembolism are less common causes of ST.

ECG Characteristics of ST

ECG characteristics of ST include the following:

- Heart rate: beat-to-beat variability with changes in activity or stress level
 - Usually less than 220/min in infants
 - Usually less than 180/min in children
- P waves: present/normal
- PR interval: constant, normal duration

- R-R interval: variable
- QRS complex: narrow (0.09 second or less)


SVT

SVT is an abnormally fast rhythm originating above the ventricles. In infants and children, the most common cause is a reentry mechanism that occurs through an accessory pathway or within the AV node. SVT is the most common tachyarrhythmia that causes cardiovascular compromise during infancy. In addition to an accessory pathway reentry or AV nodal reentry, other mechanisms that can cause SVT are atrial flutter and ectopic atrial focus.

Two outdated terms for SVT are *paroxysmal atrial tachycardia* and *paroxysmal SVT*. SVT was labeled *paroxysmal* because it occurs episodically (in paroxysms). The rapid rhythm starts and stops suddenly, often without warning.

Clinical Presentation of SVT

SVT ([Figure 45](#)) is a rapid, regular rhythm that often appears abruptly and may be episodic. During episodes of SVT, cardiopulmonary function is influenced by the child's age, duration of the tachycardia, baseline ventricular function, and ventricular rate. In infants with normal ventricular function, SVT may be present but undetected for long periods (hours or days) until cardiac output is significantly impaired. However, if baseline myocardial function is impaired (eg, in a child with congenital heart disease or cardiomyopathy), SVT can produce signs of shock in a short time.



In infants, SVT is often diagnosed when symptoms of CHF develop. *Common signs and symptoms of SVT in infants* include irritability; poor feeding; rapid breathing; unusual sleepiness; vomiting; and pale, mottled, gray, or cyanotic skin. *Common signs and symptoms of SVT in older children* include palpitations, shortness of breath, chest pain or discomfort, light-headedness, and fainting.

SVT is initially well tolerated in most infants and older children. However, it can lead to CHF and clinical evidence of shock when baseline myocardial function is impaired (eg, in a child with congenital heart disease or cardiomyopathy) or in an infant having prolonged episodes over hours to days. Ultimately, SVT can cause cardiovascular collapse.

Signs



SVT may be identified as the result of its effect on systemic perfusion. SVT with cardiopulmonary compromise can produce the signs and symptoms listed in [Table 58](#).

Table 58. Signs and Symptoms of SVT With Cardiopulmonary Compromise

Evaluate	Signs and symptoms
Airway	Usually open/patent unless level of consciousness is significantly impaired
Breathing	<ul style="list-style-type: none"> • Tachypnea • Increased work of breathing • Crackles (or “wheezing” in infants) if CHF develops • Grunting if CHF develops
Circulation	<ul style="list-style-type: none"> • Tachycardia beyond the typical range for ST and characterized by fixed rate and/or abrupt onset • Delayed capillary refill time • Weak peripheral pulses • Cool extremities • Diaphoretic, pale, mottled, gray, or cyanotic skin

	<ul style="list-style-type: none"> Hypotension Jugular venous distension (difficult to observe in young children) or hepatomegaly if CHF develops
Disability	<ul style="list-style-type: none"> Altered mental status Sleepiness or lethargy Irritability
Exposure	Defer evaluation of temperature until ABCs are supported

ECG Characteristics of SVT

ECG characteristics of SVT include the following:

- Heart rate: no beat-to-beat variability with activity
 - Usually 220/min or greater in infants
 - Usually 180/min or greater in children
- P waves: absent or abnormal (may appear after the QRS complex)
- PR interval: because P waves are usually absent, PR interval cannot be determined; in ectopic atrial tachycardia, a short PR interval may be seen
- R-R interval: often constant
- QRS complex: usually narrow; wide complex uncommon

Narrow-QRS-complex SVT: In more than 90% of children with SVT, the QRS complex is narrow ([Figure 45](#)), ie, 0.09 second or less.

Wide-QRS-complex SVT: SVT with aberrant conduction (uncommon in the pediatric age group) produces a wide QRS complex (ie, greater than 0.09 second). This form of SVT most often occurs as a result of rate-related bundle branch block (within the ventricles) or preexisting bundle branch block. It also may be caused by an accessory pathway in which electrical impulses are conducted from the atria to the ventricles through the accessory pathway rather than through the AV node. The impulse then returns to the atria through the AV node (or through a different accessory pathway).

It can be difficult to differentiate SVT with aberrant conduction from ventricular tachycardia (VT). This usually requires careful analysis of at least one 12-lead ECG. Both SVT with aberrant conduction and VT can cause similar hemodynamic instability, have similar rates, and have wide QRS complexes (greater than 0.09 second). In the pediatric age group, unless patient history or previous ECGs suggest the likelihood of SVT with aberrant conduction (eg, preexisting bundle branch block), assume that a tachycardia with a wide QRS complex is due to VT.

Comparison of ST and SVT

It may be difficult to differentiate SVT with shock from shock resulting from another etiology with compensatory ST. The characteristics in [Table 59](#) may aid in differentiating ST from SVT. Note that signs of heart failure and other signs and symptoms of poor perfusion may be absent early after the onset of SVT.

Table 59. Characteristics of ST and SVT

Characteristic	ST	SVT
History	Gradual onset Compatible with ST (eg, history of fever, pain, dehydration, hemorrhage)	Abrupt onset or termination or both Infant: Symptoms of CHF Child: Sudden onset of palpitations
Physical examination	Signs of underlying cause of ST (eg, crackles, fever, hypovolemia, anemia)	Infant: Signs of CHF (eg, rales, hepatomegaly, edema)
Heart rate	Infant: Usually <220/min Child: Usually <180/min	Infant: Usually ≥220/min Child: Usually ≥180/min
Monitor	Variability in heart rate with changes in level of activity or stimulation; slowing of heart rate with rest or treatment of underlying cause (eg, administering IV fluids for hypovolemia)	Minimal or no variability in heart rate with changes in level of activity or stimulation
ECG	P waves present/normal/upright in leads I/aVF	P waves absent/abnormal/inverted (negative) in leads II/III/aVF; if present, they usually follow the QRS complex

Chest x-ray	Usually small heart and clear lungs unless ST is caused by pneumonia, pericarditis, or underlying heart disease	Signs of CHF (eg, enlarged heart, pulmonary edema) may be present
--------------------	---	---

P waves may be difficult to identify in both ST and SVT once the ventricular rate exceeds 200/min.

Atrial Flutter

Atrial flutter is a narrow-complex tachyarrhythmia that can develop in newborn infants with normal hearts ([Figure 46](#)). It also can develop in children with congenital heart disease, especially after cardiac surgery. A reentrant mechanism typically is present in children with enlarged atria or with anatomic barriers resulting from cardiac surgery (eg, atriotomy scars or surgical anastomoses). A reentry circuit within the atria allows a wave of depolarization to travel in a circle within the atria. Because the AV node is not part of the circuit, AV conduction may be variable. The atrial rate can exceed 300/min, whereas the ventricular rate is slower and may be irregular. Classically, a “sawtooth” pattern of the P waves is present on the ECG.




Figure 46. Atrial flutter in an adolescent (atrial rate about 270/min; ventricular rate approximately 70/min).

Ventricular Tachycardia

VT is a wide-QRS-complex tachyarrhythmia generated within the ventricles ([Figure 47](#)). VT is uncommon in children. When VT with pulses is present, the ventricular rate may vary from near normal to greater than 200/min. Rapid ventricular rates compromise ventricular filling, stroke volume, and cardiac output and may deteriorate into pVT or ventricular fibrillation (VF).



Figure 47A. Ventricular tachycardia; A, Monomorphic.



Figure 47B. Polymorphic (torsades de pointes).

Most children who develop VT have underlying heart disease (or have had surgery for heart disease), long QT syndrome, or myocarditis/cardiomyopathy. They may have a family history of a sudden, unexplained death in a child or young adult, suggesting cardiomyopathy or an inherited cardiac ion “channelopathy.” Other causes of VT in children include electrolyte disturbances (eg, hyperkalemia, hypocalcemia, hypomagnesemia), drug toxicity (eg, tricyclic antidepressants, cocaine, methamphetamines), and acute systemic illnesses (eg, severe sepsis) or coronary artery abnormalities (eg, Kawasaki syndrome) causing myocardial ischemia.

ECG Characteristics of VT

ECG characteristics of VT include the following:

- Ventricular rate: at least 120/min and regular
- QRS complex: wide (greater than 0.09 second)
- P waves: often not identifiable; when present, may not be related to QRS (AV dissociation); at slower rates, atria may be depolarized in a retrograde manner, resulting in a 1:1 ventricular-to-atrial association
- T waves: typically opposite in polarity from QRS

It may be difficult to differentiate SVT with aberrant conduction from VT. Fortunately, aberrant conduction is present in less than 10% of children with SVT. In general, the healthcare provider should initially assume that a wide-complex

rhythm is VT unless the child is known to have aberrant conduction or previous episodes of wide-QRS-complex SVT.

Polymorphic VT, Including Torsades de Pointes

VT may be monomorphic (QRS complexes are uniform in appearance) or polymorphic (QRS complexes vary in appearance). If the ventricular rate is slow enough, the patient with monomorphic VT can maintain pulses. In contrast, polymorphic VT is typically associated with loss of pulses at onset or within a very short time after onset. Torsades de pointes is a distinctive form of polymorphic VT. The term *torsades de pointes* is French and means “turning on a point.” In torsades de pointes, the QRS complexes change in polarity and amplitude, appearing to rotate around the ECG isoelectric line ([Figure 47B](#)). The ventricular rate can range from 150 to 250/min. Torsades de pointes can be seen in conditions associated with a prolonged QT interval, including congenital long QT syndrome, electrolyte abnormalities, and drug toxicity. The prolonged QT interval is identified during sinus rhythm; it cannot be evaluated during the tachycardia. A rhythm strip may show the child’s baseline QT prolongation because torsades de pointes sometimes occurs in bursts that convert spontaneously to sinus rhythm.

Conditions and agents that predispose to torsades de pointes include

- Long QT syndromes (often congenital and inherited)
- Hypomagnesemia
- Hypokalemia
- Antiarrhythmic drug toxicity (ie, Class IA, quinidine, procainamide, disopyramide; Class IC, flecainide; Class III, sotalol, amiodarone)
- Other drug toxicities (eg, tricyclic antidepressants, calcium channel blockers, phenothiazines)

It is important to recognize that VT, especially polymorphic VT (including torsades de pointes), can rapidly deteriorate to VF. The long QT syndromes and other inherited arrhythmia syndromes (ie, channelopathies) are associated with sudden death from either primary VF or torsades de pointes. Polymorphic VT not associated with a prolonged QT interval during sinus rhythm is treated as generic VT.

Part 12: Managing Arrhythmias

This Part discusses managing bradycardia (slow heart rate) with a pulse and cardiopulmonary compromise and tachycardia (fast heart rate) with a palpable pulse and cardiopulmonary compromise. Providers should quickly treat symptomatic arrhythmias before they result in shock or cardiac arrest.

Learning Objective

After completing this Part, you should be able to describe clinical characteristics of instability in patients with arrhythmias.

You will also be expected to manage a child as outlined in the algorithms for bradycardia with a pulse and cardiopulmonary compromise and tachycardia with a pulse and cardiopulmonary compromise.

Principles of Managing Pediatric Arrhythmias

Whenever the child has an abnormal heart rate or rhythm, you must quickly determine if the arrhythmia is causing hemodynamic instability or other signs of deterioration. The signs of instability in a patient with arrhythmia include the following:

- Respiratory distress or failure
- Shock with poor end-organ perfusion, which may occur with or without hypotension
- Irritability or a decreased level of consciousness
- Chest pain or a vague feeling of discomfort in older children
- Sudden collapse

Priorities in initially managing arrhythmias are the same as they are for all critically ill children: support the ABCs—airway, breathing, and circulation—and treat the underlying cause.

Management: Pediatric Bradycardia With a Pulse

The Pediatric Bradycardia With a Pulse Algorithm (Figure 48) outlines the steps for evaluating and managing the child who presents with symptomatic bradycardia (bradycardia with a pulse). Refer to the Critical Concepts box Symptomatic Bradycardia and Cardiopulmonary Compromise in [Part 11](#) for more information about symptomatic bradycardia and cardiopulmonary compromise. In the text that follows, numbers refer to the corresponding steps in this algorithm.




Figure 48. Pediatric Bradycardia With a Pulse Algorithm.

Patient with bradycardia (Step 1)

Once you identify symptomatic bradycardia, assess for signs of cardiopulmonary compromise.



Critical Concepts: Assess for Cardiopulmonary Compromise (Step 2)

Assess the child for signs of cardiopulmonary compromise, including

- Hypotension
- Acutely altered mental status: acutely decreased level of consciousness
- Signs of shock

A child with primary bradycardia may benefit from evaluation by a pediatric cardiologist. However, do not delay initiating emergency treatment, including high-quality CPR, if symptoms are present.

If there are signs of cardiopulmonary compromise, go to assessment and support (Step 3):

- Maintain a patent airway.
- Assist breathing and positive pressure ventilation and oxygen as necessary.
- Use a cardiac monitor to identify rhythm; monitor pulse, blood pressure, and oximetry.

If there are no signs of cardiopulmonary compromise (Step 9),

- Support ABCs as needed.
- Consider oxygen.
- Observe and perform frequent reassessments.
- Obtain a 12-lead ECG.
- Identify and treat underlying causes.

Start CPR if heart rate is less than 60/min despite oxygenation and ventilation (Step 4).

Determine if bradycardia persists (Step 5).

If pulses, perfusion, and respirations are adequate, no emergency treatment is needed. Monitor and continue evaluating (Step 9).

If bradycardia and cardiopulmonary compromise persist (Step 6),

- Continue CPR.
- Establish IV/IO access.
- Administer epinephrine.
- Administer atropine for increased vagal tone or primary AV block.
- Consider transthoracic/transvenous pacing.
- Identify and treat underlying causes.

If bradycardia is associated with cardiopulmonary compromise (Critical Concepts box Reassess for Cardiopulmonary Compromise) and if heart rate is less than 60/min despite effective oxygenation and ventilation, continue CPR. If the bradycardia persists, proceed with medication therapy and possible pacing. Reassess the child frequently in response to each therapy provided ([Table 60](#)).

Table 60. Managing Symptomatic Bradycardia With Cardiopulmonary Compromise

Evaluate	Interventions
Airway	Support the airway (position the child or allow the child to assume a position of comfort) or open the airway (perform manual airway maneuver) if needed.
Breathing	<ul style="list-style-type: none">• Give O₂ in high concentration—use a nonrebreathing mask if available.• Assist ventilation as indicated (eg, bag-mask ventilation).• Attach a pulse oximeter to assess oxygenation.

Circulation	<ul style="list-style-type: none"> • Monitor blood pressure and assess perfusion. • Attach a monitor/defibrillator (with transcutaneous pacing capability if available). • Establish vascular access (IV or IO). • Check electrode pad position and skin contact to ensure that there are no artifacts and that the ECG tracing is accurate. • Record a 12-lead ECG if available (do not delay therapy). • Obtain appropriate laboratory studies (eg, potassium, glucose, ionized calcium, magnesium, blood gas for pH, toxicology screen).
--------------------	---

A child with primary bradycardia may benefit from evaluation by a pediatric cardiologist. However, do not delay initiating emergency treatment, including high-quality CPR, if symptoms are present.

Check pulse and reassess rhythm (Step 7).

Check the pulse every 2 minutes. If a pulse is not present, go to the Pediatric Cardiac Arrest Algorithm (Step 8) in Part 5: Recognizing and Managing Cardiac Arrest.

If a pulse is present, reassess to determine if bradycardia and cardiopulmonary compromise continue despite oxygenation, ventilation, and CPR (Step 5). If bradycardia and cardiopulmonary compromise have improved, support ABCs as needed, consider oxygen, observe and perform frequent reassessment, obtain a 12-lead ECG, and identify and treat underlying causes (Step 9). If the child is still bradycardic with cardiopulmonary compromise, administer medications, consider transthoracic/transvenous pacing, and identify and treat underlying causes.

Medications

If bradycardia and cardiopulmonary compromise continue despite oxygenation, ventilation, and CPR, administer epinephrine. Consider atropine for increased vagal tone or primary AV block.

Epinephrine

Epinephrine is indicated for symptomatic bradycardia that persists despite effective oxygenation and ventilation. Epinephrine has both α - and β -adrenergic activity. β -adrenergic activity increases heart rate and cardiac contractility, and α -adrenergic activity causes vasoconstriction. The effects of epinephrine and other catecholamines may be reduced by acidosis and hypoxia. This makes support of the airway, ventilation, oxygenation, and perfusion (with chest compressions) essential.

- For IV/IO route, give 0.01 mg/kg (0.1 mL/kg of 0.1 mg/mL concentration)
- For endotracheal route, give 0.1 mg/kg (0.1 mL/kg of 1 mg/mL concentration)
- Repeat every 3 to 5 minutes as needed

For persistent bradycardia, consider a continuous infusion of epinephrine (0.1 to 0.3 mcg/kg per minute). A continuous epinephrine infusion may be useful, particularly if the child has responded to a bolus of epinephrine. Titrate the infusion dose to clinical response.

Atropine

Atropine sulfate is a parasympatholytic (or anticholinergic) drug that accelerates sinus or atrial pacemakers and enhances atrioventricular (AV) conduction. Administer atropine instead of epinephrine for bradycardia caused by increased vagal tone, cholinergic drug toxicity (eg, organophosphates), or complete AV block. Atropine (and pacing) are preferred over epinephrine as the first-choice treatment of symptomatic AV block due to primary bradycardia. Atropine is not indicated for AV block from secondary bradycardia (ie, treatable causes such as hypoxia or acidosis). The rationale for using atropine rather than epinephrine in these situations is that epinephrine can cause ventricular arrhythmias if the myocardium is chronically abnormal or hypoxic/ischemic. If the child does not respond to atropine, use epinephrine.

Atropine may be used for the treatment of second-degree AV block (Mobitz types I and II) and third-degree AV block. The healthcare provider should recognize, however, that symptomatic AV block may not respond to atropine and the child may require pacing.

- For IV/IO route, give 0.02 mg/kg; minimum 0.1 mg, maximum 0.5 mg
 - May repeat dose once, in 5 minutes.
 - *Note:* Larger doses may be required for organophosphate poisoning.
- For endotracheal route, give 0.04 to 0.06 mg/kg.

- *Note:* IV/IO administration is preferred, but if it is not available, atropine can be administered by ET tube. Because absorption of atropine given by the endotracheal route is unreliable, a larger dose (2-3 times the IV dose) may be required.

Tachycardia may follow atropine administration, but it is generally well tolerated in the pediatric patient.

Consider Transthoracic/Transvenous Pacing

Temporary transthoracic/transvenous pacing may be lifesaving in selected cases of bradycardia caused by complete heart block or abnormal sinus node function. For example, pacing is indicated for AV block after surgical correction of congenital heart disease.

Identify and Treat Underlying Causes

Identify and treat potentially reversible causes and special circumstances that can cause bradycardia. The 2 most common potentially reversible causes of bradycardia are hypoxia and increased vagal tone. Be aware that after heart transplantation, sympathetic nerve fibers are no longer attached to the heart, so the response to sympathomimetic drugs may be unpredictable. For the same reason, anticholinergic drugs, such as atropine, may be ineffective. Early cardiac pacing may be indicated in such patients.

Treat potentially reversible causes of bradycardia as shown in [Table 61](#).

Table 61. Treatment of Bradycardia Causes

Reversible cause	Treatment
Hypoxia	Give high-concentration supplemental O ₂ with assisted ventilation as necessary.
Hydrogen ion (acidosis)	Provide ventilation to treat respiratory acidosis secondary to hypercarbia; consider sodium bicarbonate in severe metabolic acidosis.
Hyperkalemia	Restore normal potassium concentration. <ul style="list-style-type: none"> • Stop all fluids that contain potassium • Stabilize myocardium: calcium (chloride or gluconate) • Shift intracellularly: albuterol, sodium bicarbonate, insulin with glucose administration • Excrete: loop diuretics, kayexalate • Remove: dialysis
Hypothermia	Warm the child as needed, but avoid hyperthermia if the patient has experienced a cardiac arrest.
Heart block	For AV block, consider atropine, chronotropic drugs, and electrical pacing; obtain expert consultation.
Toxins/poisons/drugs	Treat with a specific antidote and provide supportive care. Some toxicologic causes of bradyarrhythmias are <ul style="list-style-type: none"> • Cholinesterase inhibitors (organophosphates, carbamates, and nerve agents) • Calcium channel blockers • β-adrenergic blockers • Digoxin and other cardiac glycosides • Clonidine and other centrally acting α_2-adrenergic agonists • Opioids • Succinylcholine
Trauma	Head trauma: Bradycardia in a child with head trauma is an ominous sign of high intracranial pressure. Provide oxygenation and ventilation. A brief period of mild hyperventilation may occasionally be used as temporizing rescue therapy in response to signs of impending herniation (eg, irregular respirations or apnea, bradycardia, hypertension, unequal or dilated pupil[s] not responsive to light, decerebrate or decorticate posturing). Obtain immediate expert assistance for relief of increased intracranial pressure.

Pulseless Arrest

If pulseless cardiac arrest develops, start CPR. Proceed according to the [Pediatric Cardiac Arrest Algorithm](#) (refer to [Part 5: Recognizing and Managing Cardiac Arrest](#)).

Managing Tachyarrhythmias

Pulseless Arrest Initial Management Questions

Answer the following questions to direct your initial management of a critically ill or injured child with a rapid heart rate:

Does the child have a pulse (or signs of circulation)?

- If no, initiate the Pediatric Cardiac Arrest Algorithm (refer to Part 5: Recognizing and Managing Cardiac Arrest).
 - *Note:* Because the accuracy of a pulse check is poor, recognizing cardiac arrest may require that you identify the absence of signs of circulation (ie, the child is unresponsive, is not breathing or only gasping). With invasive monitoring of arterial pressure, absence of arterial waveform is observed.
- If yes, proceed with the Tachycardia Algorithm.

Initial Management Priorities

As soon as you recognize a tachyarrhythmia in an infant or child, assess and look for signs of hypotension, altered mental status, shock (ie, poor perfusion), or life-threatening hemodynamic instability. Initial management priorities include the following:

- Maintain a patent airway; assist breathing as necessary.
- Use a cardiac monitor to identify the rhythm, and monitor pulse, blood pressure, and oximetry.
- Establish IV/IO access.
- Obtain a 12-lead ECG if available (but do not delay urgent intervention).
- Obtain laboratory studies (eg, potassium, glucose, ionized calcium, magnesium, blood gas to assess pH and cause of pH changes) as appropriate. *Note:* Do not delay urgent intervention for these studies.
- Assess neurologic status.
- Anticipate the need for medications depending on the type of rhythm disturbance (ie, supraventricular vs ventricular).
- Simultaneously try to identify and treat reversible causes.

Emergency Interventions

Specific emergency interventions used to treat tachyarrhythmias with a pulse are dictated by the severity of the child's condition. Treatments also vary based on the width of the observed QRS complex (narrow vs wide). Interventions may include the following:

- Vagal maneuvers (if the child with a narrow-complex tachycardia is stable)
- Medication therapy
- Synchronized cardioversion


Vagal Maneuvers

In normal infants and children, the heart rate decreases when the vagus nerve is stimulated. In patients with SVT, vagal stimulation may terminate the tachycardia by slowing conduction through the AV node. Several maneuvers can stimulate vagal activity. The success rates of these maneuvers in terminating tachyarrhythmias vary, depending on the child's age, level of cooperation, and underlying condition.



Critical Concepts: Vagal Maneuvers

Ice to the face is a vagal maneuver that can be performed in infants and children of all ages (Figure 49). Fill a small plastic bag with a mixture of ice and water. Apply it to the upper half of the child's face for 15 to 20 seconds. Do not occlude the nose or mouth.






Figure 49. Vagal maneuvers.
Note that the bag of ice water does not cover the nose or mouth and does not obstruct ventilation.

- Children old enough to cooperate can perform a Valsalva maneuver by blowing through a narrow straw
- Do not use ocular pressure because it may produce retinal injury

Support the child's airway, breathing, and circulation. If possible, obtain a 12-lead ECG before and after the maneuver; record and monitor the ECG continuously during the maneuver. *If the child is stable* and the rhythm does not convert, you may repeat the attempt. If the second attempt fails, select another method or provide medication therapy. *If the child is unstable*, attempt vagal maneuvers only while making preparations for pharmacologic or electrical cardioversion. Do not delay definitive treatment to perform vagal maneuvers.

Medication Therapy

Table 62 reviews common agents used in managing tachyarrhythmias.

Table 62. Medication Therapy Used in the Pediatric Tachycardia With a Pulse Algorithm

Drug	Indications/precautions	Dosage/administration
Adenosine	<p>Indications</p> <ul style="list-style-type: none"> • Drug of choice for treating SVT • Effective for SVT caused by reentry at the AV node (both accessory pathway and AV nodal reentry mechanisms) • May be helpful in distinguishing atrial flutter from SVT • Not effective for treating atrial flutter, atrial fibrillation, or tachycardias caused by mechanisms other than reentry through the AV node <p>Mechanism of Action</p> <ul style="list-style-type: none"> • Blocks conduction through the AV node temporarily (for about 10 seconds) <p>Precautions</p> <ul style="list-style-type: none"> • A common cause of adenosine cardioversion "failure" is that the drug is administered too slowly or with inadequate IV flush. • A brief period (10-15 seconds) of bradycardia (asystole or third-degree heart block) may ensue after administration of adenosine (Figure 50); consider warning caregiver and patient, if age appropriate, that bradycardia can be very uncomfortable. 	<p>Dose</p> <ul style="list-style-type: none"> • With continuous ECG monitoring, administer 0.1 mg/kg (maximum initial dose 6 mg) as a rapid IV bolus. • If the drug is effective, the rhythm will convert to sinus rhythm within 15 to 30 seconds after administration (Figure 50). • If there is no effect, give 1 dose of 0.2 mg/kg (maximum second dose 12 mg); this dose is more likely to be needed when the drug is administered into a peripheral (rather than central) vein. • Decrease the initial dose by approximately 75% for patients receiving carbamazepine or dipyridamole or those with transplanted hearts. <p>Administration</p> <ul style="list-style-type: none"> • Because adenosine has a short half-life (<10 seconds), administer as rapidly as possible. • The drug is rapidly taken up by vascular endothelial cells and red blood cells and metabolized by an enzyme on the surface of red blood cells (adenosine deaminase). • To enhance delivery to the site of action in the heart, use a rapid flush technique (5-10 mL normal saline). • Adenosine may be given by the IO route.
Amiodarone	<p>Indications</p> <ul style="list-style-type: none"> • May be considered when treating hemodynamically unstable SVT refractory to vagal maneuvers, adenosine, and cardioversion and for whom expert consultation is not available • Can be used for treating a wide variety of atrial and ventricular tachyarrhythmias in children • Safe and effective for hemodynamically unstable VT in children 	<p>Dose</p> <ul style="list-style-type: none"> • For supraventricular and ventricular arrhythmias with poor perfusion, a loading dose of 5 mg/kg infused over 20 to 60 minutes is recommended (maximum single dose: 300 mg). Because this drug can cause hypotension and decrease cardiac contractility, a slower rate of delivery is recommended for treating a perfusing rhythm than for cardiac arrest. Providers must weigh the potential for causing hypotension against the need to achieve a rapid drug effect.

	<p>Mechanism of Action</p> <ul style="list-style-type: none"> • Inhibits α- and β-adrenergic receptors, producing vasodilation and AV nodal suppression (this slows conduction through the AV node) • Inhibits the outward potassium current so it prolongs the QT duration • Inhibits sodium channels, which slows conduction in the ventricles and prolongs QRS duration <p>Precautions</p> <ul style="list-style-type: none"> • Drug effects may be beneficial in some patients but may also increase the risk for polymorphic VT (torsades de pointes) by prolonging the QT interval. • Potential significant acute side effects of amiodarone include bradycardia, hypotension, and polymorphic VT. • Use with caution if hepatic failure is present. • Because the pharmacology of amiodarone is complex, and it has slow and incomplete oral absorption, long half-life, and potential for long-term adverse effects, a pediatric cardiologist or similarly experienced provider should direct long-term amiodarone therapy. 	<ul style="list-style-type: none"> • Repeat doses of 5 mg/kg may be given up to a maximum of 15 mg/kg per day as needed (should not exceed the maximum recommended adult cumulative daily dose of 2.2 g over 24 hours). <p>Administration</p> <ul style="list-style-type: none"> • Rapid administration of amiodarone may cause vasodilation and hypotension; it may also cause heart block or polymorphic VT. • Monitor blood pressure frequently during administration. • Seek expert consultation when using amiodarone. • Routine use of amiodarone and another agent that prolongs the QT interval (eg, procainamide) is not recommended.
--	---	--



Figure 50. SVT converting to sinus rhythm with administration of adenosine.

Synchronized Cardioversion

Electrical cardioversion is painful. Whenever possible, establish vascular access and provide procedural sedation and analgesia before cardioversion, especially in a hemodynamically stable infant or child. If the child's condition is unstable, do not delay synchronized cardioversion to achieve vascular access. Sedation in the setting of an arrhythmia carries increased risk. When providers give procedural sedation in this setting, they must carefully select medications to minimize hemodynamic effects. If your patient is stable and you are considering cardioversion, seek expert consultation.

The next section discusses the following important concepts about cardioversion:

- Definition of synchronized cardioversion
- Potential problems with synchronized shocks
- Indications for the use of synchronized cardioversion
- Energy doses

Delivering Synchronized Shocks

Manual defibrillators are capable of delivering both unsynchronized and synchronized shocks. If the shock is unsynchronized, it is delivered at any time in the cardiac cycle. Synchronized shocks are used for cardioversion from SVT and VT with a pulse. If the shock is synchronized, shock delivery is timed to coincide with the R wave of the patient's QRS complex. The goal is to prevent VF that could result from delivery of the shock during the vulnerable period of the T wave. When you press the Shock button to deliver a synchronized shock, the defibrillator/cardioverter may seem to pause before it delivers a shock because it is waiting to synchronize shock delivery with the next QRS complex. For pulseless rhythms, it clinically does not matter whether the shock is synchronized; accordingly, unsynchronized shocks (eg, defibrillations) are used for both VF and pVT. Refer to the [Critical Concepts box Cardioversion \(for Unstable SVT or VT With a Pulse\)](#) later in this Part for a description of the procedure.

Important considerations: In theory, synchronization is simple. The operator pushes the Sync button on the defibrillator, charges the device, and delivers the shock. However, in practice there can be potential problems, such as

- In most units, the Sync button must be activated each time synchronized cardioversion is attempted. Most devices will default to an unsynchronized shock immediately after delivery of a synchronized shock.
- If the R waves of a tachycardia are undifferentiated or of low amplitude, the monitor sensors may be unable to identify them and therefore will not deliver the shock. If this occurs, increase the gain of the ECG lead being monitored or select a different ECG lead.
- Synchronization may take extra time (eg, if it is necessary to attach separate ECG electrodes or if the operator is unfamiliar with the equipment).

Indications: Use synchronized cardioversion for

- Hemodynamically unstable patients (poor perfusion, hypotension, or heart failure) with tachyarrhythmias (SVT, atrial flutter, VT), but with palpable pulses
- Elective cardioversion, under the direction of a pediatric cardiologist, for children with hemodynamically stable SVT, atrial flutter, or VT with a pulse

Energy dose: In general, cardioversion requires less energy than defibrillation does. Start with an energy dose of 0.5 to 1 J/kg for cardioversion of SVT or VT with a pulse. If the initial dose is ineffective, increase the dose to 2 J/kg. The experienced provider may increase the shock dose more gradually (eg, 0.5 J/kg, and then 1 J/kg, followed by 2 J/kg for subsequent doses). If the rhythm does not convert to sinus rhythm, reevaluate the diagnosis of SVT vs ST. Sedate if needed, but don't delay cardioversion.



Critical Concepts: Cardioversion (for Unstable SVT or VT With a Pulse)

Consider expert consultation for suspected VT.

1. Turn on the defibrillator.
2. Set lead switch to paddles (or lead I, II, or III if monitor leads are used).
3. Select adhesive pads or paddles. Use the largest pads or paddles that can fit on the patient's chest without touching each other.
4. If using paddles, apply conductive gel or paste. Be sure cables are attached to the defibrillator.
5. Consider sedation.
6. Select synchronized mode.
7. Look for markers on R waves indicating that sync mode is operative. If necessary, adjust monitor gain until sync markers occur with each R wave.
8. Select energy dose:
Initial dose: 0.5 to 1 J/kg
Subsequent doses: 2 J/kg
9. Announce "Charging defibrillator," and press charge on the defibrillator controls or apex paddle.
10. When the defibrillator is fully charged, firmly state, "I am going to shock on 3." Then count and say, "All clear!"
11. After confirming all personnel are clear of the patient, press the Shock button on the defibrillator or press both paddle discharge buttons simultaneously. Hold paddles in place until the shock is delivered.
12. Check the monitor. If tachycardia persists, prepare to attempt cardioversion again.
13. Reset the sync mode and increase the energy dose. You must reset the sync mode after each synchronized cardioversion because most defibrillators default back to unsynchronized mode after synchronized shock delivery. This default allows an immediate defibrillation (nonsynchronized) shock if the cardioversion produces VF.

Note: If VF develops, immediately begin CPR and prepare to deliver an unsynchronized shock as soon as possible. Refer to the Critical Concepts box Manual Defibrillation (for VF or pVT), in Part 5.

Other Emergency Interventions

Many other interventions (eg, digoxin, short-acting β -blockers, overdrive pacing) have been used to treat children with SVT but should be reserved for expert consultation.

Verapamil, a calcium channel blocking agent, should not be used routinely to treat SVT in infants because refractory hypotension and cardiac arrest have been reported after administration. Use verapamil with caution in children because it may cause hypotension and myocardial depression. If using verapamil in children 1 year or older, infuse the drug in a dose of 0.1 mg/kg (up to 5 mg) over at least 2 minutes with continuous ECG monitoring.

Summary of Emergency Interventions


The specific emergency interventions listed in [Table 63](#) are used to treat tachyarrhythmias with pulses, based on the width of the observed QRS complex (narrow vs wide).

Table 63. Emergency Interventions for Tachyarrhythmias With Pulses

Intervention	Narrow-complex tachyarrhythmia	Wide-complex tachyarrhythmia
Vagal maneuvers	Used for SVT	Used for SVT with aberrancy
Medication therapy	<p>Used for SVT:</p> <ul style="list-style-type: none"> Adenosine Amiodarone (seek expert consultation) Procainamide (seek expert consultation) Verapamil for children <1 year of age (seek expert consultation) <p>Used for other SVT with a pulse (eg, atrial flutter): Seek expert consultation</p>	<p>Used for VT with palpable pulses:</p> <ul style="list-style-type: none"> Amiodarone (seek expert consultation) Procainamide (seek expert consultation) <p>Used for torsades de pointes:</p> <ul style="list-style-type: none"> Magnesium <p>Used for SVT with abnormal/aberrant intraventricular conduction:</p> <ul style="list-style-type: none"> Adenosine Amiodarone (seek expert consultation) Procainamide (seek expert consultation)
Synchronized cardioversion	<p>Used for</p> <ul style="list-style-type: none"> SVT Atrial flutter (seek expert consultation) 	Used for VT with palpable pulses

Pediatric Tachycardia With a Pulse Algorithm

The [Pediatric Tachycardia With a Pulse Algorithm](#) ([Figure 51](#)) outlines the steps for assessing and managing a child presenting with symptomatic tachycardia with or without cardiopulmonary compromise.



Initial assessment and support (Step 1):

- Maintain a patent airway; assist breathing as necessary.
- Administer oxygen.
- Use a cardiac monitor to identify the rhythm; monitor pulse, blood pressure, and oximetry.
- Establish IV/IO access.
- Obtain 12-lead ECG if available.

Evaluate rhythm with 12-lead ECG or monitor (Step 2):

- Determine if this is sinus tachycardia or a tachyarrhythmia.

Probable sinus tachycardia if (Step 3):

- P waves present/normal
- Variable RR interval

- Infant rate usually less than 220/min
- Child rate usually less than 180/min

Search for and treat cause (Step 4).

Assess for cardiopulmonary compromise (Step 5):

- Acutely altered mental status
- Signs of Shock
- Hypotension

If there are signs of cardiopulmonary compromise, evaluate QRS duration (Step 6):

- This is to characterize if the QRS complex is narrow (0.09 sec or less) or wide (greater than 0.09).

If narrow QRS complex (0.09 seconds or less), treat as probable supraventricular tachycardia (Step 7):

- P waves are absent/abnormal.
- Respiratory rate interval is not variable.
- Infant rate is usually 220/min or greater.
- Child rate is usually 180/min or greater.
- History of abrupt rate change.

Administer adenosine or perform synchronized cardioversion (Step 8):

- If IV/IO access is present, give adenosine. Adenosine is the drug of choice for most common forms of SVT caused by a reentrant pathway involving the AV node. For IV/IO route, give 0.1 mg/kg (maximum first dose 6 mg). If the first dose is ineffective, you may give 1 dose of 0.2 mg/kg (maximum second dose 12 mg). Use a rapid bolus with a *rapid flush* 2-syringe technique (5-10 mL normal saline).
- If IV/IO access is not available, or if adenosine is ineffective, perform synchronized cardioversion. In general, cardioversion requires less energy than defibrillation does. Start with an energy dose of 0.5 to 1 J/kg for cardioversion of SVT. If the initial dose is ineffective, increase the dose to 2 J/kg. The experienced provider may increase the shock dose more gradually (eg, 0.5 J/kg, and then 1 J/kg, followed by 2 J/kg for subsequent doses). If the rhythm does not convert to sinus rhythm, reevaluate the diagnosis of SVT vs sinus tachycardia (ST). Sedate if needed but don't delay cardioversion

If wide complex (greater than 0.09 seconds), treat as possible VT (Step 9).

Perform synchronized cardioversion (Step 10):

- Expert consultation is advised before any additional drug therapies are administered for treatment of a wide-complex tachycardia.

If there are no signs of cardiopulmonary compromise, evaluate QRS duration (Step 11):

- This is to characterize if the QRS complex is narrow (0.09 sec or less) or wide (greater than 0.09).

If narrow complex (0.09 seconds or less), treat as probable supraventricular tachycardia (Step 12):

- P waves are absent/abnormal.
- Respiratory rate interval is not variable.
- Infant rate is usually 220/min or greater.
- Child rate is usually 180/min greater.
- History of abrupt rate change

Consider vagal maneuvers (Step 13):

In the stable patient with SVT, try the following:

- Place a bag with ice water over the upper half of an infant's face (without obstructing the airway).
- Ask older children to bear down, blow through an obstructed straw, or try to push the plunger out of a 10 mL syringe by blowing into the tip.
- Monitor and record the ECG continuously before, during, and after attempted vagal maneuvers. If the maneuvers fail, you may attempt them a second time. Do not apply ocular pressure. For more information, refer to the Critical Concepts box Vagal Maneuvers earlier in this Part.

If IV/IO access is present, give adenosine (Step 14):

- For SVT resistant to vagal maneuvers, establish vascular access and administer adenosine. Adenosine is the drug of choice for most common forms of SVT caused by a reentrant pathway involving the AV node. For IV/IO route, give 0.1 mg/kg (maximum first dose 6 mg). If the first dose is ineffective, you may give 1 dose of 0.2 mg/kg (maximum second dose 12 mg). Use a rapid bolus with a *rapid flush* 2-syringe technique (5-10 mL normal saline).

If wide complex (greater than 0.09 second), treat as possible VT (Step 15).

If rhythm is regular and QRS monomorphic, consider adenosine (Step 16):

- Avoid adenosine if rhythm is irregular as this may result in an unstable rhythm.

Expert consultation is recommended. If a child with a wide-complex tachycardia is hemodynamically stable, early consultation with a pediatric cardiologist or other provider with appropriate expertise is strongly recommended (Step 17).

Pharmacologic Conversion

Establish vascular access and consider administering one of the following medications:

- Amiodarone: For IV/IO route, give 5 mg/kg over 20 to 60 minutes.
- Procainamide: For IV/IO route, give 15 mg/kg over 30 to 60 minutes.

Seek expert consultation when giving amiodarone or procainamide. Do not routinely administer amiodarone and procainamide together or with other medications that prolong the QT interval. If these initial efforts do not terminate the rapid rhythm, reevaluate the rhythm.

If not already administered, consider adenosine, because a wide-complex tachycardia could be SVT with aberrant ventricular conduction (Step 12).

Part 13: Post–Cardiac Arrest Care

As soon as ROSC develops after cardiac arrest or resuscitation from severe shock or respiratory failure, a systematic approach to assessment and support of the respiratory, cardiovascular, and neurologic systems, including providing targeted temperature management, is critical. Although effective resuscitation is a major focus of the PALS Provider Course, ultimate outcome is often determined by the subsequent care the child receives. This includes safe transport to a center with expertise in caring for seriously ill or injured children.

One objective of optimal post–cardiac arrest care is to avoid common causes of both early and late morbidity and mortality. Early mortality can be caused by hemodynamic instability and respiratory complications. Late morbidity and mortality can result from multiorgan failure, including brain injury.

The extent of post–cardiac arrest evaluation and management is influenced by the PALS provider’s scope of practice and available resources.

Learning Objective

After completing this Part, you should be able to implement post–cardiac arrest care.

During the course you will learn about the phases of post–cardiac arrest care. This will include optimizing oxygenation, ventilation, and perfusion, stabilizing cardiopulmonary function, and providing neurologic care, including targeted temperature management.

Goals of Therapy

For optimal post–cardiac arrest care, identify and treat organ system dysfunction. This includes

- Providing adequate oxygenation and ventilation
- Supporting tissue perfusion and cardiovascular function
- Avoiding hypotension
- Correcting acid-base and electrolyte imbalances
- Maintaining appropriate glucose concentration
- Providing targeted temperature management: avoiding hyperthermia and considering need for therapeutic hypothermia
- Ensuring adequate analgesia and sedation

Post–cardiac arrest management consists of 2 general phases to stabilize the child.

The first phase is immediate post–cardiac arrest management. During this phase you will continue to provide advanced life support for immediate life-threatening conditions and focus on the ABCs:

- **Airway and Breathing:** Assess and support airway, oxygenation, and ventilation. At this time, you will typically use diagnostic equipment and assessments—such as monitoring end-tidal CO₂ by capnography, arterial blood gas analysis, and chest x-ray—to further establish the adequacy of oxygenation and ventilation and to confirm endotracheal (ET) tube position in the mid-trachea.
- **Circulation:** Assess and maintain adequate blood pressure and perfusion. Treat arrhythmias. Diagnostic assessments—such as lactate concentration, central venous O₂ saturation, and base deficit—provide information on adequacy of tissue perfusion. As you proceed with evaluation, identify and treat any reversible or contributing causes of the arrest or critical illness.

In the second phase of post–cardiac arrest management, provide broader multiorgan supportive care, including targeted temperature management. After the child is stabilized, coordinate transfer or transport to a tertiary care setting as appropriate.

Primary Goals

The primary goals of post–cardiac arrest management are to

- Optimize and stabilize airway, oxygenation, ventilation, and cardiopulmonary function with emphasis on restoring and maintaining vital organ perfusion and function (especially of the brain)
- Prevent secondary organ injury
- Identify and treat the cause of acute illness
- Institute measures that may improve long-term, neurologically intact survival
- Minimize the risk of deterioration of the child during transport to a higher level of care

Post–Cardiac Arrest Care Checklist

This checklist (Figure 52) includes a systematic approach to assessing and caring for the child after cardiac arrest. Assess the child by using the systematic approach (refer to [Part 4: Systematic Approach to the Seriously Ill or Injured Child](#)). In addition to the primary assessment, your evaluation will often include the secondary assessment as well as diagnostic evaluation. The secondary assessment is a review of patient history and a focused physical examination. Diagnostic evaluation includes invasive and noninvasive monitoring and appropriate laboratory and nonlaboratory tests.

This Part discusses evaluation and management of the following systems during the post–cardiac arrest period:

- Respiratory system
- Cardiovascular system
- Neurologic system

Components of Post-Cardiac Arrest Care	Check
Oxygenation and ventilation	
Measure oxygenation and target normoxemia 94%-99% (or child's normal/appropriate oxygen saturation).	<input type="checkbox"/>
Measure and target PaO ₂ , appropriate to the patient's underlying condition and limit exposure to severe hyperoxia or hypoxemia.	<input type="checkbox"/>
Hemodynamic monitoring	
Set specific hemodynamic goals during post-cardiac arrest care and review daily.	<input type="checkbox"/>
Monitor vital cardiac telemetry.	<input type="checkbox"/>
Monitor arterial blood pressure.	<input type="checkbox"/>
Monitor central venous pressure, urine output, and central venous oxygen saturation to help guide therapies.	<input type="checkbox"/>
Use peripheral arterial lines with or without noninvasive measurements to maintain a systolic blood pressure greater than the 5th percentile for age and sex.	<input type="checkbox"/>
Targeted temperature management (TTM)	
Measure and continuously monitor core temperature.	<input type="checkbox"/>
Prevent and treat fever immediately after arrest and during rewarming.	<input type="checkbox"/>
If patient is comatose apply TTM (32°C–34°C) followed by 36°C–37.5°C or only TTM (36°C–37.5°C).	<input type="checkbox"/>
Prevent shivering.	<input type="checkbox"/>
Monitor blood pressure and treat hypotension during rewarming.	<input type="checkbox"/>
Neuromonitoring	
If patient has encephalopathy and resources are available, monitor with continuous electroencephalogram.	<input type="checkbox"/>
Treat seizures.	<input type="checkbox"/>
Consider early brain imaging to diagnose treatable causes of cardiac arrest.	<input type="checkbox"/>
Electrolytes and glucose	
Measure blood glucose and avoid hypoglycemia.	<input type="checkbox"/>
Maintain electrolytes within normal ranges to avoid possible life-threatening arrhythmias.	<input type="checkbox"/>
Medication	
Treat with sedatives and analgesics.	<input type="checkbox"/>
Prognosis	
Always consider multiple modalities (clinical and other) over any single predictive factor.	<input type="checkbox"/>
Remember that assessments may be modified by TTM or induced hypothermia.	<input type="checkbox"/>
Consider electroencephalogram in conjunction with other factors after the first 7 days after cardiac arrest.	<input type="checkbox"/>
Consider neuroimaging such as magnetic resonance imaging during the first 7 days.	<input type="checkbox"/>

Figure 52. Post–Cardiac Arrest Care Checklist.

Respiratory System

Management Priorities

Continue to monitor and support the child's airway, oxygenation, and ventilation. Look for clinical signs and objective measurements of adequate oxygenation and ventilation. (Refer to [Part 7: Recognizing Respiratory Distress and Failure](#) for more information on assessment of the respiratory system.) During resuscitation, high-flow O₂, inhaled medications, and ET intubation may be required. In the post–cardiac arrest phase, elective intubation may be appropriate to achieve airway control and support the child during diagnostic studies, such as a computed tomography scan. If the child is being manually ventilated, transition to mechanical ventilation.

The goals of respiratory management in the immediate post–cardiac arrest period are listed in [Table 64](#).

Table 64. Goals of Respiratory Management in Immediate Post–Cardiac Arrest Period

Goal	Considerations
Maintain adequate oxygenation (generally an O₂ saturation of 94%-99%) to reduce the risk of reperfusion injury	<p>Once ROSC is achieved, titrate O₂ administration to target normoxemia while ensuring that hypoxemia is strictly avoided. Maintain an O₂ saturation of 94%-99%, avoiding hypoxemia and hyperoxia (an O₂ saturation of 100% can correspond to a PaO₂ of anywhere between 80 and approximately 500 mm Hg).</p> <p>Determining optimal PaO₂ and O₂ saturation requires evaluation of the child's arterial O₂ content because it is an important determinant of tissue O₂ delivery. If the child is anemic, tissue O₂ delivery may be better maintained by achieving a high</p>

	<p>PaO₂ and O₂ saturation. In comparison, an O₂ saturation of 94%-100% is typically adequate in a child with a normal hemoglobin concentration and normal O₂ consumption and no cyanotic heart disease. Thus, oxygen is titrated to a value appropriate to the specific patient condition.</p>
<p>Maintain adequate ventilation and PaCO₂ appropriate to the patient</p>	<p>It is reasonable to target a PaCO₂ (or end-tidal CO₂ if PaCO₂ is unavailable) that is appropriate to each child's clinical condition and to limit exposure to severe hypercapnia or hypocapnia. For example, for most patients with neurologic injury, a normal PaCO₂ is desirable to avoid hypocapnia (hypocarbica) or hypercapnia (hypercarbica). However, in children with asthma and respiratory failure, rapid correction of hypercapnia is unnecessary. Conversely, in children with congenital heart disease and pulmonary hypertension, hypercapnia must be avoided.</p> <p>Efforts to achieve normocarbica with mechanical ventilation in a child with asthma could result in complications such as pneumothorax.</p>

General Recommendations

General recommendations for assessment and management of the respiratory system may include those listed in [Table 65](#).

Table 65. General Recommendations for Assessing and Managing the Respiratory System

Assessment	
<p>Monitoring</p>	<ul style="list-style-type: none"> Continuously monitor the following parameters (at a minimum): SpO₂ and heart rate by pulse oximetry (compare pulse oximetry heart rate with ECG and pulse rate to ensure that pulse oximeter values are accurate) Heart rate and rhythm If the patient is intubated, monitor end-tidal CO₂ by capnography if equipment and expertise are available, or intermittently confirm exhaled CO₂ by colorimetric device. Always monitor exhaled CO₂ by either capnography or colorimetric device during intrahospital and interhospital transport to aid in immediate detection of inadvertent extubation. If the child is already intubated, verify tube position, openness/patency, and security. After proper tube position is confirmed, ensure that the tube is well taped and that tube position at the lip or gum is documented. <p><i>Providers must use both clinical assessment and confirmatory devices (such as monitoring of exhaled CO₂) to verify proper tube placement immediately after intubation, during transport, and when the child is moved (eg, from gurney to bed).</i></p>
<p>Physical examination</p>	<ul style="list-style-type: none"> Observe for adequate and equal chest rise bilaterally and auscultate for abnormal or asymmetric breath sounds. Monitor for evidence of respiratory compromise (eg, tachypnea, increased work of breathing, agitation, decreased responsiveness, poor air exchange, cyanosis) or inadequate respiratory effort.
<p>Laboratory tests</p>	<ul style="list-style-type: none"> Obtain an arterial sample for arterial blood gas analysis if possible. If the child is mechanically ventilated, obtain the arterial blood gas 10-15 minutes after establishing initial ventilator settings; ideally, correlate blood gases with capnographic end-tidal CO₂ to enable noninvasive monitoring of ventilation.
<p>Other tests</p>	<ul style="list-style-type: none"> Obtain a chest x-ray to verify correct depth of the ET tube insertion and position in the mid-trachea, and to identify pulmonary conditions that may require specific treatment (eg, pneumothorax, aspiration).
Management	
<p>Oxygenation</p>	<ul style="list-style-type: none"> If the child is not intubated, provide supplemental O₂ with a partial or a nonrebreathing mask until you confirm adequate SpO₂. After ROSC, adjust inspired O₂ concentration to achieve an SpO₂ of 94%-99%. If the child has an SpO₂ of <90% while receiving 100% inspired O₂, consider noninvasive ventilatory support or ET intubation with mechanical ventilation and positive end-expiratory pressure. If the child has a cyanotic cardiac lesion, adjust the O₂ saturation goal to the child's baseline SpO₂ and clinical status.
<p>Ventilation</p>	<ul style="list-style-type: none"> Assist ventilation as needed, targeting a normal PaCO₂ (or ET-CO₂ if PaCO₂ is unavailable) (ie, 35-45 mm Hg) if the child's lung function was previously normal. Remember that normalizing PaCO₂ may not be appropriate in all situations. Avoid routine hyperventilation in children with neurologic problems unless there are signs of impending cerebral herniation. Limit exposure to severe hypercapnia or hypocapnia.
<p>Respiratory failure</p>	

	<ul style="list-style-type: none"> Intubate the trachea if O₂ administration and other interventions do not achieve adequate oxygenation and ventilation. Intubate if needed to maintain an open/patent airway and adequate oxygenation and ventilation in the child with decreased level of consciousness. In some patients, CPAP or noninvasive ventilation may be adequate. Use age- and weight-appropriate ventilator settings. Verify ET tube position, openness/patency, and security; retape, if needed, before transport. Assess for a large glottic air leak. Consider reintubation with a cuffed tube or a larger tube if the glottic air leak prevents adequate chest rise, oxygenation, or ventilation. Weigh the risk of removing the advanced airway against the benefit of improving tidal volume, oxygenation, and ventilation. If a cuffed ET tube is in place and is inflated, check the cuff pressure (goal for most tubes is <20 to 25 cm H₂O; follow the manufacturer's recommendations) or assess for the presence of a minimal glottic air leak at an inspiratory pressure of <20 to 25 cm H₂O. Insert a gastric tube to relieve and help prevent gastric inflation. <p><i>Use the "DOPE" mnemonic to troubleshoot acute deterioration in a mechanically ventilated patient. (Refer to the Sudden Deterioration in an Intubated Patient section in the Resources for Managing Respiratory Emergencies in Part 8 of this manual.)</i></p>
Analgesia and sedation	<ul style="list-style-type: none"> Control pain with analgesics (eg, fentanyl or morphine) and anxiety with sedatives (eg, lorazepam or midazolam) as needed. Administer sedation and analgesia to all responsive intubated patients. <p><i>Use lower doses of sedatives or analgesics if the child is hemodynamically unstable; titrate the dose while stabilizing hemodynamic function. When used in equipotent doses, morphine is more likely than fentanyl to cause hypotension because morphine causes histamine release.</i></p>
Neuromuscular blockade	<ul style="list-style-type: none"> For the intubated patient with poor oxygenation and ventilation despite adequate sedation and analgesia, assess for acute causes of deterioration by using the DOPE mnemonic. Then consider neuromuscular blocking agents (eg, vecuronium, pancuronium) with sedation. Indications for use of neuromuscular blocking agents include <ul style="list-style-type: none"> High peak or mean airway pressure caused by high airway resistance or reduced lung compliance Patient ventilator asynchrony Difficult airway <p><i>Neuromuscular blockade may reduce the risk of ET tube displacement. Be aware that neuromuscular blockers do not provide sedation or analgesia and will mask seizures. Neuromuscular blockers will also eliminate many signs of agitation that may signal inadequate oxygenation and ventilation. When using neuromuscular blockers, always ensure that the child is adequately sedated by evaluating for signs of stress, such as tachycardia, hypertension, pupil dilation, or tearing.</i></p>

Cardiovascular System

Management Priorities

Ischemia resulting from cardiac arrest and subsequent reperfusion can cause circulatory dysfunction that can last for hours after ROSC. Compromised tissue perfusion and oxygenation from shock and respiratory failure can have secondary adverse effects on cardiovascular function. Providers should maintain adequate blood pressure, cardiac output, and distribution of blood flow to restore or maintain tissue oxygenation and delivery of metabolic substrates. Circulatory management priorities include

- Restoring and maintaining intravascular volume (preload)
- Treating myocardial dysfunction
- Controlling arrhythmias
- Maintaining normal blood pressure and adequate systemic perfusion
- Maintaining adequate SpO₂ and PaO₂
- Maintaining adequate hemoglobin concentration
- Considering therapies to reduce metabolic demand (eg, support ventilation and reduce temperature)

This section includes

- General recommendations for advanced evaluation and management of the cardiovascular system
- The PALS Management of Shock After ROSC Algorithm
- Information about administering maintenance fluids

Review [Part 9: Recognizing Shock](#) and [Part 10: Managing Shock](#) for more information about the pathophysiology of shock and the use of fluid therapy and medications to maintain cardiac output and tissue perfusion.

General Recommendations

Refer to [Table 66](#) for general recommendations for assessment and management of the cardiovascular system.

Table 66. General Recommendations for Assessing and Managing the Cardiovascular System

Assessment	
Monitoring	<ul style="list-style-type: none"> Monitor the following frequently or continuously: <ul style="list-style-type: none"> Heart rate and rhythm by cardiac monitor Blood pressure and pulse pressure (noninvasively or invasively) SpO₂ by pulse oximetry Urine output by urinary catheter Temperature In the critical care setting, also consider monitoring <ul style="list-style-type: none"> Central venous pressure by central venous catheter Central venous O₂ saturation by catheter providing continuous ScvO₂ or by intermittent blood draw Trends in regional venous oxygenation by near infrared spectroscopy Cardiac function (eg, echocardiogram) and/or cardiac output by noninvasive monitoring <p><i>Noninvasive blood pressure monitoring (ie, by automated blood pressure devices) is often unreliable in children with poor perfusion or frequent arrhythmias. Blood pressure monitoring with an indwelling arterial catheter and monitoring system is more reliable in these children, provided the catheter is open/patent and the transducer is appropriately zeroed and leveled.</i></p>
Physical examination	<ul style="list-style-type: none"> Repeat the physical examination (eg, evaluate quality of central and peripheral pulses, heart rate, capillary refill, blood pressure, extremity temperature and color) frequently until the child is stable. Monitor end-organ function (eg, neurologic function, renal function, skin perfusion) to assess circulatory function.
Laboratory tests	<ul style="list-style-type: none"> Arterial or venous blood gas Hemoglobin and hematocrit Serum glucose, electrolytes, blood urea nitrogen (BUN), creatinine, calcium Consider monitoring lactate and central venous O₂ saturation <p>In addition to pH, note the magnitude of any metabolic acidosis (base deficit). A persistent metabolic (lactic) acidosis suggests inadequate cardiac output and O₂ delivery. Serum electrolytes can help identify an anion gap acidosis. If the child has an elevated anion gap but normal lactate, consider other causes of acidosis, such as toxins or uremia.</p> <p>The difference in O₂ saturation between an arterial and superior vena caval blood sample [S(a-v)O₂] provides information about the balance of oxygen supply vs demand. Assuming that O₂ consumption remains constant, a high S(a-v)O₂ difference (>35-40) suggests low oxygen delivery. This may be caused by a fall in cardiac output, or arterial oxygen content. When oxygen delivery falls, there will be increased O₂ extraction in the tissues (ie, blood flow and O₂ delivery are decreased, so O₂ extraction must increase), producing a fall in the superior vena caval oxygen saturation. Alternatively, increased O₂ consumption with stable cardiac output can result in a widened a-vO₂ difference as well.</p> <p>Troponin concentrations are frequently elevated after cardiac arrest, especially if defibrillation was performed.</p>
Nonlaboratory tests	<ul style="list-style-type: none"> Perform a chest x-ray to evaluate ET tube insertion depth and position in the mid-trachea, assess heart size, and identify pulmonary edema or other pathology. Evaluate a 12-lead ECG for arrhythmias or evidence of myocardial ischemia. Consider echocardiography if there is concern about pericardial tamponade or myocardial dysfunction. <p><i>A small heart is often present with reduced cardiac preload or severe lung hyperinflation. A large heart may be associated with normal or increased cardiac preload, pericardial effusion, CHF, or when the patient is unable to take a deep breath (eg, with severe abdominal distension).</i></p>
Management	
Intravascular volume	<ul style="list-style-type: none"> Establish secure vascular access (if possible, 2 catheters, either IV or IO). Administer fluid boluses (10-20 mL/kg of isotonic crystalloid over 5-20 minutes) as needed to establish adequate intravascular volume. Smaller boluses of fluid (5-10 mL/kg) administered over 10-20 minutes may be appropriate in the setting of heart failure. Adjust the fluid administration rate to replace fluid deficits and meet ongoing requirements. Avoid excessive fluid administration in the presence of myocardial dysfunction, heart failure, or respiratory failure. Consider the need for colloid or blood administration. Calculate maintenance fluid requirements and administer as appropriate.

	<p>Do not use boluses of hypotonic or dextrose-containing fluids for volume resuscitation.</p> <p>Refer to <u>Administering Maintenance Fluids</u> later in this Part.</p>
Blood pressure	<ul style="list-style-type: none"> • Treat hypotension aggressively, titrating volume and vasoactive medications as appropriate. • Post–cardiac arrest systolic blood pressure must be maintained above the fifth percentile for age (for children 1-10 years of age, estimated with the following formula: 70 mm Hg + (age in years × 2)). • If hypotension is due to excessive vasodilation (eg, sepsis), early use of a vasopressor may be indicated. • The use of adrenergic agents during resuscitation may produce elevation in systemic vascular resistance and cause hypertension. Because the half-life of these agents is relatively short, assess for other causes of hypertension in the post–cardiac arrest phase (eg, pain, anxiety, seizures). <p><i>Treatment of hypotension is crucial to avoid secondary multisystem injury. Refer to the PALS Management of Shock After ROSC Algorithm for more information about management of hypotensive and normotensive shock.</i></p>
Tissue oxygenation	<ul style="list-style-type: none"> • Provide supplemental O₂ in a sufficient concentration to ensure adequate oxygenation. • After ROSC following cardiac arrest, titrate O₂ to maintain adequate SpO₂ (94%-99%). • Support adequate perfusion. • Consider transfusion with packed red blood cells for patients with low hematocrit and signs of inadequate O₂ delivery.
Metabolic demand	<ul style="list-style-type: none"> • Consider ET intubation and assisted ventilation to reduce the work of breathing. • Control pain with analgesia (eg, morphine, fentanyl). • Control agitation as needed with sedation (eg, lorazepam, midazolam); rule out hypoxemia, hypercarbia, or poor perfusion as potential causes of agitation. • Control fever with antipyretics and a cooling blanket if needed (neuromuscular blockade may be appropriate to control shivering). <p><i>Caution: Sedatives or analgesics may cause hypotension. Consider expert consultation before elective ET intubation. The use of sedatives or analgesics, ET intubation, and initiating positive-pressure ventilation can all precipitate cardiovascular collapse in a child with poor myocardial function.</i></p>
Arrhythmias	<ul style="list-style-type: none"> • Monitor for tachyarrhythmias and bradyarrhythmias and treat aggressively. • If bradycardia develops, first ensure adequate oxygenation and ventilation; if the heart rate remains <60/min with signs of poor perfusion despite adequate oxygenation and ventilation, initiate CPR. Bradycardia can also occur in the setting of hypothermia; consider whether perfusion and blood pressure are adequate. • If arrhythmia persists, control with drugs or electrical therapy per algorithms. • Seek expert consultation for arrhythmia management. <p><i>Refer to <u>Part 11: Recognizing Arrhythmias</u> and <u>Part 12: Managing Arrhythmias</u> for more information.</i></p>
Postarrest myocardial dysfunction	<ul style="list-style-type: none"> • Anticipate postarrest myocardial dysfunction in the first 24 hours after ROSC. • Consider vasoactive agents to improve contractility and/or decrease afterload if blood pressure is adequate. • Correct metabolic abnormalities that can contribute to poor myocardial function (eg, acidosis, hypocalcemia, hypoglycemia). • Consider positive-pressure ventilation (noninvasive ventilation or via ET tube) to improve left ventricular function. <p><i>Myocardial dysfunction is common in children after resuscitation from cardiac arrest. Postarrest myocardial dysfunction can produce hemodynamic instability and secondary organ injury and may precipitate another cardiac arrest.</i></p>

Treatment of Shock

After resuscitation from cardiac arrest or shock, hemodynamic compromise may result from a combination of

- Inadequate intravascular volume
- Decreased cardiac contractility
- Either increased or decreased systemic vascular resistance (SVR) or pulmonary vascular resistance

Children with cardiogenic shock typically have poor myocardial function and a compensatory increase in systemic and pulmonary vascular resistances in an attempt to maintain an adequate blood pressure. The increased SVR may become detrimental because it increases left ventricular afterload. Very low SVR most commonly occurs in children with early septic shock. When children with septic shock do not respond to bolus fluid administration (ie, the shock is fluid refractory), they may have high rather than low SVR and poor myocardial function, similar to cardiogenic shock.

Support of Systemic Perfusion

The parameters listed in [Table 67](#) can be manipulated to optimize systemic perfusion.

Table 67. Parameters to Optimize Systemic Perfusion

Parameters to optimize	Action (when needed)
Preload	<ul style="list-style-type: none"> Administer fluid bolus.
Contractility	<ul style="list-style-type: none"> Administer inotropes or inodilators. Correct hypoxia, electrolyte and acid-base imbalances, and hypoglycemia/hypocalcemia. Treat poisonings (eg, administer antidotes if available).
Afterload (SVR)	<ul style="list-style-type: none"> Administer vasopressors or vasodilators as appropriate.
Heart rate	<ul style="list-style-type: none"> Administer chronotropes for bradycardia (eg, epinephrine). Administer antiarrhythmics. Correct hypoxia. Consider pacing.

Refer to the [Pathophysiology of Shock](#) section in [Part 9](#) for a discussion of preload, afterload, and contractility.

Managing Shock After ROSC

The [PALS Management of Shock After ROSC Algorithm](#) ([Figure 53](#)) outlines evaluation and management steps after cardiac arrest. Numbers in the text refer to the corresponding steps in the algorithm.




Figure 53. PALS Management of Shock After ROSC Algorithm.

Optimize oxygenation and ventilation (Step 1)

Achieving adequate oxygenation and ventilation is important for supporting cardiovascular function. Titrate FiO_2 to maintain an O_2 saturation of 94% to 99%; wean O_2 concentration if saturation is 100%. Consider placing an advanced airway and using waveform capnography if not yet established. If possible, target a PCO_2 that is appropriate for the patient's conditions, and limit exposure to severe hypercapnia or hypocapnia.

Fluid therapy (Step 2)

The first intervention to consider for treating shock is administering a 20 mL/kg IV/IO bolus of isotonic crystalloid. If you suspect the commonly occurring postarrest myocardial dysfunction, consider administering a smaller fluid bolus (5-10 mL/kg) over 10 to 20 minutes, and then reassess. If the child demonstrates signs of poor cardiac function (eg, large liver, pulmonary edema, jugular venous distention, large heart on chest x-ray), carefully evaluate the need for fluid administration. Excessive fluid administration can worsen cardiopulmonary function. Consider the need for inotropic and/or vasopressor support for fluid-refractory shock.

Reassess the child frequently and after each fluid bolus to determine response to therapy.

Possible contributing factors (Step 3)

Consider factors that may contribute to postarrest shock, including metabolic derangements and conditions such as hypovolemia and cardiac tamponade (H's and T's).

Hypotensive shock (Step 4)

If the child remains hypotensive after bolus fluid administration, consider infusion of one or a combination of the following drugs:

- Epinephrine: for IV/IO route, give 0.03 to 0.2 mcg/kg per minute* *and/or*
- Norepinephrine: for IV/IO route, give 0.03 to 0.5 mcg/kg per minute*

*Upper limit dosing range can be highly variable and should be based on clinical scenarios.

Ensure that cardiac preload is adequate; base your choice of drug on the most likely cause of hypotension (inadequate heart rate, poor contractility, excessive vasodilation, or a combination of factors). If the heart rate is abnormally low, administering catecholamine may increase the heart rate and cardiac output. However, when catecholamines cause extreme tachycardia, they increase myocardial O₂ demand.

Epinephrine

Epinephrine is a potent vasoactive agent that can either lower or increase SVR depending on the infusion dose. Low-dose infusions generally produce β -adrenergic effects (increased heart rate and contractility and vasodilation); higher doses generally produce α -adrenergic effects (vasoconstriction). Because there is great interpatient variability, titrate the drug to the desired clinical effect. Epinephrine may be preferable to dopamine in children (especially infants) with marked circulatory instability and hypotensive shock.

Norepinephrine

Norepinephrine is a potent inotropic and peripheral vasoconstricting agent. Titrate the infusion to treat shock with low SVR (septic, anaphylactic, spinal) that is unresponsive to bolus fluid administration.

Normotensive shock (Step 5)

If the child is normotensive but remains poorly perfused after bolus fluid administration, consider administering one or a combination of the following drugs:

- Low-dose epinephrine: for IV/IO route, give 0.03 to 0.05 mcg/kg per minute *and/or*
- Milrinone: for IV/IO route, load with 50 mcg/kg over 10 to 60 minutes. Loading doses may cause hypotension; infusion: 0.25 to 0.75 mcg/kg per minute

Low-Dose Epinephrine

Refer to [Epinephrine](#) in [Hypotensive Shock \(Step 4\)](#), earlier in this Part.

Milrinone

Milrinone is an inodilator that augments cardiac output with little effect on heart rate and myocardial O₂ demand. Use an inodilator for treatment of myocardial dysfunction with increased systemic or pulmonary vascular resistance. You may need to administer additional fluids because the vasodilatory effects will expand the vascular space and could lead to hypotension.

Compared with drugs like dopamine and norepinephrine, inodilators have a long half-life. In addition, whenever the infusion rate is changed, there will be a long delay in reaching a new steady-state hemodynamic effect (4.5 hours with milrinone). The adverse effects may persist for several hours after you stop the infusion. Milrinone is renally excreted, and the dose may need to be decreased in patients with renal disease.

Other post–cardiac arrest considerations (Step 6)

ROSC affects multiple organ systems. Monitor for and treat agitation and seizures with appropriate medications. Monitor for and treat hypoglycemia. Assess blood gas, serum, electrolytes, and calcium. Correct metabolic derangements, noting that treating metabolic acidosis is best accomplished by treating the underlying cause of the acidosis (ie, restore perfusion in shock). Provide targeted temperature management: consider therapeutic hypothermia for children who remain comatose and prevent or aggressively treat fever. Arrange for transfer to an appropriate pediatric critical care unit.

Administer Maintenance Fluids

Maintenance Fluid Composition

After initial stabilization, adjust the rate and composition of IV fluids based on the patient's condition. If cardiovascular function is adequate, consider administering maintenance fluids once intravascular volume has been restored and fluid deficits have been replaced. When you calculate the maintenance fluid requirements and plan for fluid administration, include the fluid administered with vasoactive drug infusions.

In the first hours after resuscitation, the appropriate composition of IV fluids is an isotonic crystalloid (0.9% NaCl or lactated Ringer's), with or without dextrose, based on the child's condition and age. Avoid hypotonic fluids in critically ill children in the post-cardiac arrest phase. Recent evidence suggests that buffered solutions (eg, lactated Ringer's or other electrolyte-containing isotonic solution) may be preferred because hyperchloremia has been associated with poorer outcomes. However, if you use one of these solutions, you must ensure that the child is receiving some source of dextrose because hypoglycemia will likely occur.

You may add specific components to maintenance fluids based on the clinical condition:

- Include dextrose in IV fluids for infants and for children who are hypoglycemic or at risk for hypoglycemia.
- Add potassium chloride (KCl) 10 to 20 mEq/L for children with adequate renal function and documented urine output once periodic monitoring of potassium is available. Do not add KCl to maintenance fluid in children with hyperkalemia, renal failure, muscle injury, or severe acidosis.

Maintenance Fluid Calculation by 4-2-1 Method

Use the 4-2-1 method as a practical approach to estimating hourly maintenance fluid requirements in children ([Table 68](#)).

Table 68. Estimating Maintenance Fluid Requirements

Weight (kg)	Estimated hourly fluid requirements	Sample collection
<10	4 mL/kg per hour	8-kg infant: 4 mL/kg per hour × 8 kg = 32 mL/h
10-20	40 mL/h + 2 mL/kg per hour for each kilogram between 10 and 20 kg*	15-kg child: 40 mL/h + 2 mL/kg per hour × 5 kg = 50 mL/h
>20	60 mL/h + 1 mL/kg per hour for each kilogram above 20 kg	30-kg child: 60 mL/h + 1 mL/kg per hour × 10 kg = 70 mL/h

*An alternate calculation of maintenance hourly fluid rate for patients weighing more than 20 kg is weight in kilograms + 40 mL/h.

Once you have calculated the estimated maintenance fluid requirements, adjust the actual rate of fluid administration to the child's clinical condition (eg, pulse, blood pressure, systemic perfusion, urine output) and hydration level.

Neurologic System

Management Priorities

During the post-cardiac arrest period, providers should aim to preserve brain function and prevent secondary neuronal injury. Neurologic management priorities include

- Maintaining adequate brain perfusion
- Maintaining normoglycemia
- Providing targeted temperature management: prevent or aggressively treat fever and consider therapeutic hypothermia if indicated
- Treating increased intracranial pressure
- Treating seizures; searching for and treating cause

General Recommendations

Refer to [Table 69](#) for general recommendations for assessment and management of the neurologic system.

Table 69. General Recommendations for Assessing and Managing the Neurologic System

Assessment	
Monitoring	<ul style="list-style-type: none"> • Monitor temperature.

	<ul style="list-style-type: none"> Monitor heart rate and systemic blood pressure. <p><i>In children with poor peripheral perfusion, reliable monitoring of core temperature requires invasive devices (rectal, bladder, esophageal thermometer).</i></p>
Physical examination	<ul style="list-style-type: none"> Perform frequent, brief neurologic assessments (eg, Glasgow Coma Scale, pupil responses, gag reflex, corneal reflexes, oculoccephalic reflexes). Identify signs of impending cerebral herniation. Identify seizure activity. Identify abnormal neurologic findings, including abnormal movements (posturing/myoclonus/hyperreflexia). <p><i>Signs of impending cerebral herniation include unequal or dilated unresponsive pupils, posturing, hypertension, bradycardia, respiratory irregularities or apnea, and reduced response to stimulation. A sudden increase in intracranial pressure (ICP) (if monitoring is in place) is often observed. Other causes of central nervous system dysfunction are hypoxic-ischemic brain injury, hypoglycemia, convulsive or nonconvulsive seizures, toxins/drugs, electrolyte abnormalities, hypothermia, traumatic brain injury, stroke or intracranial hemorrhage, and central nervous system infection.</i></p> <p><i>Refer to the section <u>Disability</u>, in <u>Part 4</u> for more information on neurologic assessment.</i></p>
Laboratory tests	<ul style="list-style-type: none"> Perform point-of-care glucose testing; repeat measurement after treatment of hyperglycemia or hypoglycemia. Obtain serum electrolytes, point-of-care glucose, and serum ionized calcium concentration if seizure activity is present; measure concentrations of anticonvulsant medications if the child was receiving these agents. Consider toxicologic studies if poisoning or overdose is suspected. Consider cerebral spinal fluid studies if central nervous system infection is suspected, but defer a lumbar puncture if the patient's cardiopulmonary status is not stable.
Nonlaboratory tests	<ul style="list-style-type: none"> Consider a computed tomography scan if central nervous system dysfunction or neurologic deterioration is present. Consider an electroencephalogram (EEG) if nonconvulsive status epilepticus is suspected or seizures are a concern during administration or duration of effect of neuromuscular blockers. EEGs performed within the first 7 days after pediatric cardiac arrest may be considered in prognosticating neurologic outcome at the time of hospital discharge, but should not be used as the sole criterion.
Management	
Brain perfusion	<ul style="list-style-type: none"> Optimize brain perfusion by supporting cardiac output and arterial O₂ content. Avoid hyperventilation unless there are signs of impending cerebral herniation. <p><i>Support cardiac output by optimizing heart rate, preload, afterload, and contractility. Refer to the section <u>Support of Systemic Perfusion</u> earlier in this Part for more information.</i></p>
Blood glucose	<ul style="list-style-type: none"> Treat hypoglycemia. Monitor glucose concentration. In general, try to avoid causing or worsening hyperglycemia. In the critical care setting, consider treating persistent hyperglycemia; careful monitoring is needed to prevent hypoglycemia. <p><i>Although hyperglycemia is associated with poor outcome in critically ill children, the relative benefits of active treatment of hyperglycemia vs risks of hypoglycemia in critically ill children remains uncertain. In most animal studies, hyperglycemia at the time of cerebral ischemia produces a worse outcome, but the effect of hyperglycemia occurring after ROSC is less clear.</i></p>
Targeted temperature management	<p>Provide targeted temperature management. This includes preventing or aggressively treating fever and considering therapeutic hypothermia.</p> <p>Prevent/Treat Fever</p> <ul style="list-style-type: none"> Prevent fever; adjust environmental temperature as needed (cooling devices may be used to maintain a target temperature and prevent fever). Aggressively treat fever (temperature 38°C or higher) with antipyretics and cooling devices (eg, cooling blanket) or procedures. Do not actively rewarm a post–cardiac arrest patient who has a temperature between 32°C and 37°C after ROSC unless hypothermia is contributing to hemodynamic instability. <p><i>Fever adversely influences recovery from ischemic brain injury and is associated with poor outcome after resuscitation from cardiac arrest. Metabolic O₂ demand increases by 10% to 13% for each degree Celsius elevation of temperature above normal. Increased metabolic demand may worsen neurologic injury. Furthermore, fever increases the release of inflammatory mediators, cytotoxic enzymes, and neurotransmitters, which increase brain injury.</i></p> <p>Therapeutic Hypothermia</p> <ul style="list-style-type: none"> For infants and children remaining comatose after out-of-hospital cardiac arrest, it is reasonable either to maintain 5 days of continuous normothermia (36°-37.5°C) or to maintain 2 days of initial

	<p>continuous hypothermia (32°-34°C), followed by 3 days of continuous normothermia.</p> <ul style="list-style-type: none"> For infants and children remaining comatose after in-hospital cardiac arrest, there is insufficient evidence to recommend cooling over normothermia, but all patients should receive targeted temperature management that prevents or aggressively treats fever. During cooling, treatment or prevention of shivering is often required. Monitor for and treat complications of hypothermia, including diminished cardiac output, arrhythmia, infection, pancreatitis, coagulopathy, thrombocytopenia, hypophosphatemia, and hypomagnesemia.
Increased ICP	<ul style="list-style-type: none"> Elevate the head of bed to 30° if blood pressure is adequate and no spinal precautions are in place. Keep head in midline. Support adequate ventilation to maintain normocapnia. If signs of impending cerebral herniation develop (eg, irregular respirations or apnea, bradycardia, hypertension, unequal or dilated pupil[s] not responsive to light, decerebrate or decorticate posturing), a brief period of mild hyperventilation may occasionally be used as temporizing rescue therapy. Consider mannitol or hypertonic saline for acute herniation syndrome. For children with neurosurgical conditions (eg, traumatic brain injury, intracranial hemorrhage), obtain expert consultation about indications for monitoring of ICP and/or neurosurgical intervention. <p><i>Prolonged hyperventilation is not effective to treat increased ICP, and excessive hyperventilation may worsen neurologic outcome. Hypocarbica results in cerebral vasoconstriction, reducing cerebral blood flow. Hyperventilation also reduces venous return and cardiac output, contributing to cerebral ischemia.</i></p>
Seizures	<ul style="list-style-type: none"> Treat seizures aggressively. Therapeutic options include a benzodiazepine (eg, lorazepam, midazolam), fosphenytoin/phenytoin, levetiracetam or a barbiturate (eg, phenobarbital). Monitor blood pressure carefully if you use phenytoin or phenobarbital, because these drugs may cause hypotension. Search for a correctable metabolic cause, such as hypoglycemia, hyponatremia, or hypocalcemia. Consider toxins or metabolic disease as the etiology. Consult a neurologist if available.

Appendix

BLS Competency Testing

BLS Skills Testing Checklists

The Child CPR and AED Skills Testing Checklist and the Infant CPR Skills Testing Checklist provide detailed descriptions of the CPR skills that you will be expected to perform. Your instructor will evaluate your CPR skills during the skills test on the basis of these descriptions.

If you perform a specific skill exactly as described in the critical performance criteria details, the instructor will check that specific skill as “passing.” If you do not perform a specific skill exactly as it is described, the skill will not be checked off and you will require remediation in that skill.

Study the BLS skills testing checklists so that you will be able to perform each skill correctly.

Pediatric Advanced Life Support
**Child CPR and AED
Skills Testing Checklist**



American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Student Name _____ Date of Test _____

Hospital Scenario: "You are working in a hospital or clinic, and you see a child who has suddenly collapsed in the hallway. You check that the scene is safe and then approach the patient. Demonstrate what you would do next."

Prehospital Scenario: "You arrive on the scene for a child who is not breathing. No bystander CPR has been provided. You approach the scene and ensure that it is safe. Demonstrate what you would do next."

Assessment and Activation

- ☐ Checks responsiveness
- ☐ Shouts for help/Activates emergency response system/Sends for AED
- ☐ Checks breathing
- ☐ Checks pulse

Once student shouts for help, instructor says, "Here's the barrier device. I am going to get the AED."

Cycle 1 of CPR (30:2) *CPR feedback devices preferred for accuracy

Child Compressions

- ☐ Performs high-quality compressions*:
 - Hand placement on lower half of sternum
 - 30 compressions in no less than 15 and no more than 18 seconds
 - Compresses at least one third the depth of the chest, about 2 inches (5 cm)
 - Complete recoil after each compression

Child Breaths

- ☐ Gives 2 breaths with a barrier device:
 - Each breath given over 1 second
 - Visible chest rise with each breath
 - Resumes compressions in less than 10 seconds

Cycle 2 of CPR (repeats steps in Cycle 1) Only check box if step is successfully performed

- ☐ Compressions
- ☐ Breaths
- ☐ Resumes compressions in less than 10 seconds

Rescuer 2 says, "Here is the AED. I'll take over compressions, and you use the AED."

AED (follows prompts of AED)

- ☐ Powers on AED
- ☐ Correctly attaches pads
- ☐ Clears for analysis
- ☐ Clears to safely deliver a shock
- ☐ Safely delivers a shock

Resumes Compressions

- ☐ Ensures compressions are resumed immediately after shock delivery
 - Student directs instructor to resume compressions or
 - Student resumes compressions

STOP TEST

Instructor Notes

- Place a check in the box next to each step the student completes successfully.
- If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation).

Test Results Circle **PASS** or **NR** to indicate pass or needs remediation:

PASS

NR

Instructor Initials _____ Instructor Number _____ Date _____

Pediatric Advanced Life Support Child CPR and AED Skills Testing Critical Skills Descriptors

- 1. Assesses victim and activates emergency response system (this must precede starting compressions) within a maximum of 30 seconds. After determining that the scene is safe:**
 - Checks for responsiveness by tapping and shouting
 - Shouts for help/directs someone to call for help *and* get AED/defibrillator
 - Checks for no breathing or no normal breathing (only gasping)
 - Scans from the head to the chest for a minimum of 5 seconds and no more than 10 seconds
 - Checks carotid pulse
 - Can be done simultaneously with check for breathing
 - Checks for a minimum of 5 seconds and no more than 10 seconds
- 2. Performs high-quality chest compressions (initiates compressions immediately after recognition of cardiac arrest)**
 - Correct hand placement
 - Lower half of sternum
 - 2-handed (second hand on top of the first or grasping the wrist of the first hand) or 1-handed
 - Compression rate of 100 to 120/min
 - Delivers 30 compressions in 15 to 18 seconds
 - Compression depth and recoil—compress at least one third the depth of the chest, approximately 2 inches (5 cm)
 - Use of a commercial feedback device or high-fidelity manikin is required
 - Complete chest recoil after each compression
 - Minimizes interruptions in compressions
 - Delivers 2 breaths so less than 10 seconds elapses between last compression of one cycle and first compression of next cycle
 - Compressions resumed immediately after shock/no shock indicated
- 3. Provides 2 breaths by using a barrier device**
 - Opens airway adequately
 - Uses a head tilt–chin lift maneuver or jaw thrust
 - Delivers each breath over 1 second
 - Delivers breaths that produce visible chest rise
 - Avoids excessive ventilation
 - Resumes chest compressions in less than 10 seconds
- 4. Performs same steps for compressions and breaths for Cycle 2**
- 5. AED use**
 - Powers on AED
 - Turns AED on by pushing button or lifting lid as soon as it arrives
 - Correctly attaches pads
 - Places proper-sized pads for victim's age in correct location
 - Clears for analysis
 - Clears rescuers from victim for AED to analyze rhythm (pushes analyze button if required by device)
 - Communicates clearly to all other rescuers to stop touching victim
 - Clears to safely deliver shock
 - Communicates clearly to all other rescuers to stop touching victim
 - Delivers a shock
 - Resumes chest compressions immediately after shock delivery
 - Does *not* turn off AED during CPR
- 6. Resumes compressions**
 - Ensures that high-quality chest compressions are resumed immediately after shock delivery
 - Performs same steps for compressions



Infant CPR Skills Testing Checklist



American
Heart
Association.

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Student Name _____ Date of Test _____

Hospital Scenario: "You are working in a hospital or clinic when a woman runs through the door, carrying an infant. She shouts, 'Help me! My baby's not breathing.' You have gloves and a pocket mask. You send your coworker to activate the emergency response system and to get the emergency equipment."

Prehospital Scenario: "You arrive on the scene for an infant who is not breathing. No bystander CPR has been provided. You approach the scene and ensure that it is safe. Demonstrate what you would do next."

Assessment and Activation

- | | |
|--|--|
| <input type="checkbox"/> Checks responsiveness | <input type="checkbox"/> Shouts for help/Activates emergency response system |
| <input type="checkbox"/> Checks breathing | <input type="checkbox"/> Checks pulse |

Once student shouts for help, instructor says, "Here's the barrier device."

Cycle 1 of CPR (30:2) *CPR feedback devices preferred for accuracy

Infant Compressions

- ☐ Performs high-quality compressions*:
 - Placement of 2 fingers or 2 thumbs in the center of the chest, just below the nipple line
 - 30 compressions in no less than 15 and no more than 18 seconds
 - Compresses at least one third the depth of the chest, about 1½ inches (4 cm)
 - Complete recoil after each compression

Infant Breaths

- ☐ Gives 2 breaths with a barrier device:
 - Each breath given over 1 second
 - Visible chest rise with each breath
 - Resumes compressions in less than 10 seconds

Cycle 2 of CPR (repeats steps in Cycle 1) Only check box if step is successfully performed

- | | | |
|---------------------------------------|----------------------------------|---|
| <input type="checkbox"/> Compressions | <input type="checkbox"/> Breaths | <input type="checkbox"/> Resumes compressions in less than 10 seconds |
|---------------------------------------|----------------------------------|---|

Rescuer 2 arrives with bag-mask device and begins ventilation while Rescuer 1 continues compressions with 2 thumb-encircling hands technique.

Cycle 3 of CPR

- | | | | |
|--|--|--|---|
| <input type="checkbox"/> Powers on AED | <input type="checkbox"/> Correctly attaches pads | <input type="checkbox"/> Clears for analysis | <input type="checkbox"/> Clears to safely deliver a shock |
| <input type="checkbox"/> Safely delivers a shock | | | |

Rescuer 1: Infant Compressions

- ☐ Performs high-quality compressions*:
 - 15 compressions with 2 thumb-encircling hands technique
 - 15 compressions in no less than 7 and no more than 9 seconds
 - Compress at least one third the depth of the chest, about 1½ inches (4 cm)
 - Complete recoil after each compression

Rescuer 2: Infant Breaths

This rescuer is not evaluated.

Cycle 4 of CPR

Rescuer 2: Infant Compressions

This rescuer is not evaluated.

Rescuer 1: Infant Breaths

- ☐ Gives 2 breaths with a bag-mask device:
 - Each breath given over 1 second
 - Visible chest rise with each breath
 - Resumes compressions in less than 10 seconds

STOP TEST

Instructor Notes

- Place a check in the box next to each step the student completes successfully.
- If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation).

Test Results Circle **PASS** or **NR** to indicate pass or needs remediation:

PASS

NR

Instructor Initials _____ Instructor Number _____ Date _____

© 2020 American Heart Association

Pediatric Advanced Life Support Infant CPR Skills Testing Critical Skills Descriptors

- Assesses victim and activates emergency response system (this must precede starting compressions) within a maximum of 30 seconds. After determining that the scene is safe:**
 - Checks for responsiveness by tapping and shouting
 - Shouts for help/directs someone to call for help *and* get emergency equipment
 - Checks for no breathing or no normal breathing (only gasping)
 - Scans from the head to the chest for a minimum of 5 seconds and no more than 10 seconds
 - Checks brachial pulse
 - Can be done simultaneously with check for breathing
 - Checks for a minimum of 5 seconds and no more than 10 seconds
- Performs high-quality chest compressions during 1-rescuer CPR (initiates compressions within 10 seconds of identifying cardiac arrest)**
 - Correct placement of hands/fingers in center of chest
 - 1 rescuer: 2 fingers just below the nipple line or 2 thumb encircling technique
 - Compression rate of 100 to 120/min
 - Delivers 30 compressions in 15 to 18 seconds
 - Adequate depth for age
 - Infant: at least one third the depth of the chest (approximately 1½ inches [4 cm])
 - Use of a commercial feedback device or high-fidelity manikin is required
 - Complete chest recoil after each compression
 - Appropriate ratio for age and number of rescuers
 - 1 rescuer: 30 compressions to 2 breaths
 - Minimizes interruptions in compressions
 - Delivers 2 breaths so less than 10 seconds elapses between last compression of one cycle and first compression of next cycle
- Provides effective breaths with bag-mask device during 2-rescuer CPR**
 - Opens airway adequately
 - Delivers each breath over 1 second
 - Delivers breaths that produce visible chest rise
 - Avoids excessive ventilation
 - Resumes chest compressions in less than 10 seconds
- Switches compression technique at appropriate interval as prompted by the instructor (for purposes of this evaluation). Switch should take no more than 5 seconds.**
- Performs high-quality chest compressions during 2-rescuer CPR**
 - Correct placement of hands/fingers in center of chest
 - 2 rescuers: 2 thumb–encircling hands just below the nipple line
 - Compression rate of 100 to 120/min
 - Delivers 15 compressions in 7 to 9 seconds
 - Adequate depth for age
 - Infant: at least one third the depth of the chest (approximately 1½ inches [4 cm])

- Complete chest recoil after each compression
- Appropriate ratio for age and number of rescuers
 - 2 rescuers: 15 compressions to 2 breaths
- Minimizes interruptions in compressions
 - Delivers 2 breaths so less than 10 seconds elapses between last compression of one cycle and first compression of next cycle

Initial Assessment—Pediatric Assessment Triangle*



Appearance

From the doorway, your initial observation of the patient. Observe for

- Abnormal tone
- Decreased interaction
- Inconsolable
- Abnormal look/gaze
- Abnormal speech/cry

Work of Breathing

- Abnormal sounds
- Abnormal positioning
- Retractions
- Flaring
- Apnea/gasping

Circulation to the Skin

- Pallor
- Mottling
- Dusky
- Cyanosis

*If the patient is unresponsive, not breathing, or only gasping, initiate the Pediatric BLS Algorithm for Healthcare Providers—Single Rescuer or Pediatric BLS Algorithm for Healthcare Providers—2 or More Rescuers in [Part 2](#).

Primary Assessment



A: Airway

Assessment

- Is the airway maintainable?
- Is the airway clear?
- If no to any of these, below are the interventions*

Interventions

- Maintain airway patency by positioning, using OPA
- Suction as indicated
- Advanced airway (eg, supraglottic airway or endotracheal tube)
- If inserting an advanced airway, verify correct placement with waveform capnography

B: Breathing

Assessment

- Adequate depth and rate of respirations
- Chest rise
- Noisy breathing (eg, grunting, stridor, wheezing)
- Use of accessory muscles, nasal flaring
- Pulse oximetry*

Interventions

- Provide high-flow O₂
- Bag-mask device with or without OPA
- Advanced airway
- Avoid excessive ventilation

C: Circulation

Assessment

- Adequate peripheral and/or central pulse
- Heart rate
- Blood pressure*
- Capillary refill—peripheral and/or central
- Skin color and temperature
- Level of consciousness

Interventions

- Obtain IV/IO access
- Consider fluid resuscitation

D: Disability

Assessment

- Quickly assess for responsiveness, level of consciousness, and pupillary response to light
- AVPU: Alert, Voice, Pain, and Unresponsive
- Check point-of-care glucose

Interventions

- Spinal motion restrictions
- Correct hypoglycemia
- Consider naloxone for acute opioid toxicity

E: Exposure

Assessment

- Remove clothing to perform a physical examination (anterior and posterior), looking for obvious signs of trauma, bleeding, burns, unusual markings, rashes, or medical alert bracelets

- Temperature

Interventions

- Ensure normothermia
- Control bleeding
- Decontamination

*If at any part of this sequence you find that a patient has a life-threatening condition, correction of that condition takes precedence over establishing baseline vital sign measures, such as blood pressure or pulse oximetry. When the primary assessment is completed and after life-threatening problems have been addressed, the healthcare provider proceeds to the secondary assessment (Consensus Statement: Emergency Medical Services for Children—Definitions and Pediatric Assessment Approaches. April 2005. Updated July 2015).

Remediation

Any provider who does not pass both skills tests will practice and undergo remediation during the remediation lesson at the end of the course.

Providers who require remediation and retesting will be tested in the entire skill.

Skills Station Competency Checklists

Airway Management Skills Station Competency Checklist

Critical Performance Steps

- Verbalizes difference between high-flow and low-flow O₂ delivery systems
 - High flow: O₂ flow exceeds patient inspiratory flow, preventing entrainment of room air if system is tight-fitting; delivers nearly 1.00 FIO₂ (eg, nonrebreathing mask with reservoir high-flow nasal cannula)
 - Low flow (≤ 10 L/min): patient inspiratory flow exceeds O₂ flow, allowing entrainment of room air; delivers 0.22 to 0.60 FIO₂ (eg, nasal cannula, simple O₂ mask)
- Verbalizes maximum nasal cannula flow rate for standard nasal cannula (4 L/min)
- Opens airway by using head tilt–chin lift maneuver while keeping mouth open (jaw thrust for trauma victim)
 - For more information, see instructor demonstration
- Verbalizes different indications for OPA and NPA
 - OPA only for unconscious victim without a gag reflex
 - NPA for conscious or semiconscious victim
- Selects correctly sized airway by measuring
 - OPA from corner of mouth to angle of mandible
- Inserts OPA correctly
- Verbalizes assessment for adequate breathing after insertion of OPA
- Suctions with OPA in place; states suctioning not to exceed 10 seconds
- Selects correct mask size for ventilation
 - For more information, see [Part 8: Managing Respiratory Distress and Failure](#)
- Assembles bag-mask device, opens airway, and creates seal by using E-C clamp technique
- With bag-mask device, gives 1 breath every 2 to 3 seconds for about 30 seconds. Gives each breath in approximately 1 second; each breath should cause chest rise
- Endotracheal Intubation
 - States equipment needed for endotracheal (ET) tube intubation procedure
 - Demonstrates technique to confirm proper ET tube placement by physical exam and by using an exhaled CO₂ device
 - Secures ET tube
 - Suctions with ET tube in place
 - For more information, see [Table 38. Pre-event Equipment Checklist for Endotracheal Intubation](#)

The following steps are optional. They are demonstrated and evaluated only when the student's scope of practice involves ET intubation.

- Endotracheal intubation

- Prepares equipment for ET intubation
- Inserts ET tube correctly

Rhythm Disturbances/Electrical Therapy Skills Station Competency Checklist

Critical Performance Steps

- Applies 3 ECG leads correctly (or local equipment if >3 leads are used)
 - Negative (white) lead: to right shoulder
 - Positive (red) lead: to left lower ribs
 - Ground (black, green, brown) lead: to left shoulder
 - For more information, see instructor demonstration
- Demonstrates correct operation of monitor
 - Turns monitor on
 - Adjusts device to manual mode (not AED mode) to display rhythm in standard limb leads (I, II, III) or paddles/electrode pads
 - For more information, see instructor demonstration
- Verbalizes correct electrical therapy for appropriate core rhythms
 - Synchronized cardioversion for unstable SVT, VT with pulses
 - Defibrillation for pulseless VT, VF
 - For more information, see [Part 11: Recognizing Arrhythmias](#) and [Part 12: Managing Arrhythmias](#)
- Selects correct paddle/electrode pad for infant or child; places paddles/electrode pads in correct position
 - For more information, see [Part 12: Managing Arrhythmias](#)
- Demonstrates correct and safe synchronized cardioversion
 - Places device in synchronized mode
 - Selects appropriate energy (0.5 to 1.0 J/kg for initial shock)
 - Charges, clears, and delivers current
 - For more information, see [Part 12: Managing Arrhythmias](#)
- Demonstrates correct and safe manual defibrillation
 - Places device in unsynchronized mode
 - Selects energy (2 to 4 J/kg for initial shock)
 - Charges, clears, and delivers current
 - For more information, see [Part 12: Managing Arrhythmias](#)

Vascular Access Skills Station Competency Checklist

Critical Performance Steps

- Verbalizes indications for IO insertion
 - For more information, see [Resources for Managing Circulatory Emergencies](#) in [Part 10](#)
- Verbalizes sites for IO insertion (anterior tibia, distal femur, medial malleolus, anterior-superior iliac spine)
- Verbalizes contraindications for IO placement
 - Fracture in extremity
 - Previous insertion attempt in the same bone
 - Infection overlying bone
- Inserts IO catheter safely
- Verbalizes how to confirm IO catheter is in correct position; verbalizes how to secure IO catheter
- Attaches IV line to IO catheter; demonstrates giving IO fluid bolus by using 3-way stopcock and syringe
 - For more information, see Instructor demonstration
- Shows how to determine correct drug doses by using a color-coded length-based tape or other resource
 - For more information, see [Table 56. Color-Coded Length-Based Resuscitation Tape](#)
- *Optional:* Verbalizes correct procedure for establishing IV access

Rhythm Recognition Review

--



Figure 54. Rhythm Strip 1:
Normal sinus rhythm, rate
100/min.

Note that every P wave in [Figure 54](#) conducts to the ventricle, resulting in a QRS complex. Be aware that normal heart rates are age dependent in the pediatric population. For example, a heart rate of 75/min would be normal for a 10-year-old but bradycardic for a neonate. Likewise, a rate of 140/min would be normal for an infant but tachycardic for an adolescent.



Figure 55. Rhythm Strip 2:
Sinus bradycardia.

Note that the P waves in [Figure 55](#) result in a QRS complex (conducted to the ventricle). The rate is very slow (approximately 45/min). Sinus bradycardia is often a manifestation of hypoxemia and acidosis. It may be seen in healthy children, particularly during sleep.



Figure 56. Rhythm Strip 3:
Sinus tachycardia, 180/min.

Note that P waves in [Figure 56](#) are visible preceding every QRS. The rate for sinus tachycardia may vary according to age. In an infant, sinus tachycardia could be as high as 220/min.



Figure 57. Rhythm Strip 4:
Sinus rhythm with first-
degree heart block.

Note that the PR interval in [Figure 57](#) is prolonged (0.3 second). This is often a reflection of increased vagal tone and may be seen in healthy children. Less often, it can be a sign of intrinsic atrioventricular node disease, myocarditis, electrolyte disturbances (such as hyperkalemia), hypoxemia, drug toxicity (such as digoxin, β -blocker, or calcium channel blocker), or acute rheumatic fever.



Figure 58. Rhythm Strip 5:
Second-degree heart block,
Mobitz type I, or
Wenckebach.

Note that the PR interval in [Figure 58](#) progressively prolongs until a P wave fails to conduct to the ventricle. Like first-degree heart block, this is often seen in healthy children, especially during sleep. It may also be a manifestation of drug toxicity, such as digoxin, β -blocker, or calcium channel blocker.



Figure 59. Rhythm Strip 6:
Second-degree heart block,
Mobitz type II.

Note that some, but not all, of the P waves in [Figure 59](#) do not conduct to the ventricle. There is no progressive prolongation of the PR interval. This is a sign of intrinsic conduction system disease, typically related to cardiac surgery or myocardial inflammation or infarction.



**Figure 60. Rhythm Strip 7:
Third-degree (complete)
heart block with ventricular
escape rhythm.**

Note that none of the P wave in [Figure 60](#) conducts to the ventricle. Often the QRS complex “marches” at a constant interval because of junctional or ventricular escape rhythm. There is no relation between P waves and QRS complexes. Occasionally this is a result of severe hypoxemia and acidosis. This may also be a manifestation of damage to the atrioventricular node or extensive conduction system disease, such as that seen after cardiac surgery, or myocarditis, or with congenital complete heart block.



**Figure 61. Rhythm Strip 8:
Supraventricular
tachycardia, 230/min.**

Note that the QRS complexes in [Figure 61](#) are narrow and regular, the rate is very fast (greater than 200/min), and P waves are not obvious.



**Figure 62. Rhythm Strip 9:
Atrial flutter.**

Note that there is a “sawtooth” pattern to the P waves in [Figure 62](#), reflecting an extremely rapid atrial rate. Conduction of the P waves to the ventricle may be variable, resulting in an irregular QRS rate.



**Figure 63. Rhythm Strip 10:
Ventricular tachycardia,
150/min.**

Note that the QRS complexes in [Figure 63](#) are wide (greater than 0.09 second), regular, and fast. The QRS morphologies are all identical, characterizing it as monomorphic ventricular tachycardia. P waves are not identifiable.



**Figure 64. Rhythm Strip 11:
Polymorphic ventricular
tachycardia.**

Note that the QRS complexes in [Figure 64](#) are wide, irregular, and very fast (greater than 200/min). The QRS complexes vary in appearance, characterizing it as polymorphic. In fact, there is a phase change during this recording where the QRS complexes are initially positive and then become negative before returning to positive polarity. This is a torsades de pointes (“turning of points”) type of polymorphic ventricular tachycardia, seen most often in states where the QT interval is prolonged during the baseline.



**Figure 65. Rhythm Strip 12:
Ventricular fibrillation.**

Note that there are no definable QRS complexes in Figure 65, just irregular disorganized electrical activity.



**Figure 66. Rhythm Strip 13:
Sinus rhythm with peaked T
waves.**

Note how the T waves in Figure 66 have increased amplitude and are even larger than the QRS complexes. This may be indicative of hyperkalemia.

Learning Station Competency Checklists and PALS Case Scenario Testing Checklists

The learning station competency checklists provide detailed descriptions of the critical performance steps that you will perform at each learning/skills station.

Additionally, the case scenario testing checklists provide the steps you will perform during each case scenario test. Your instructor will evaluate your skills on the basis of these descriptions.

If you perform a specific skill exactly as described in the critical performance criteria details, the instructor will check that specific skill as “passing.” If you do not perform a specific skill exactly as it is described, the skill will not be checked off and you will require remediation in that skill.

Study these checklists so that you will be able to perform each skill correctly.

Airway Management Skills Station Competency Checklist



American Heart Association

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Verbalizes difference between high-flow and low-flow O ₂ delivery systems <ul style="list-style-type: none"> High flow: O₂ flow exceeds patient inspiratory flow, preventing entrainment of room air if system is tight-fitting; delivers nearly 1.00 FIO₂, eg, nonrebreathing mask with reservoir, high-flow nasal cannula Low flow (≤10 L/min): patient inspiratory flow exceeds O₂ flow, allowing entrainment of room air; delivers 0.22 to 0.60 FIO₂, eg, standard nasal cannula, simple O₂ mask 	
Verbalizes maximum nasal cannula flow rate for standard nasal cannula (4 L/min)	
Opens airway by using head tilt–chin lift maneuver while keeping mouth open (jaw thrust for trauma victim)	
Verbalizes different indications for OPA and NPA <ul style="list-style-type: none"> OPA only for unconscious victim without a gag reflex NPA for conscious or semiconscious victim 	
Selects correctly sized airway by measuring <ul style="list-style-type: none"> OPA from corner of mouth to angle of mandible 	
Inserts OPA correctly	
Verbalizes assessment for adequate breathing after insertion of OPA	
Suctions with OPA in place; states suctioning not to exceed 10 seconds	
Selects correct mask size for ventilation	
Assembles bag-mask device, opens airway, and creates seal by using E-C clamp technique	
With bag-mask device, gives 1 breath every 2 to 3 seconds for 30 seconds. Gives each breath in approximately 1 second; each breath should cause chest rise	
Endotracheal Intubation <ul style="list-style-type: none"> States equipment needed for endotracheal (ET) tube intubation procedure Demonstrates technique to confirm proper ET tube placement by physical exam and by using an exhaled CO₂ device Secures ET tube Suctions with ET tube in place 	
The following steps are optional. They are demonstrated and evaluated only when the student's scope of practice involves ET intubation.	
Endotracheal Intubation <ul style="list-style-type: none"> Prepares equipment for ET intubation Inserts ET tube correctly 	

STOP TEST

Instructor Notes

- Place a check in the box next to each step the student completes successfully.
- If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation).

Test Results	Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____	Instructor Number _____	Date _____	

Rhythm Disturbances/ Electrical Therapy Skills Station Competency Checklist



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Applies 3 ECG leads correctly (or local equipment if >3 leads are used) <ul style="list-style-type: none"> Negative (white) lead: to right shoulder Positive (red) lead: to left lower ribs Ground (black, green, brown) lead: to left shoulder 	
Demonstrates correct operation of monitor <ul style="list-style-type: none"> Turns monitor on Adjusts device to manual mode (not AED mode) to display rhythm in standard limb leads (I, II, III) or paddles/electrode pads 	
Verbalizes correct electrical therapy for appropriate core rhythms <ul style="list-style-type: none"> Synchronized cardioversion for unstable SVT, VT with pulses Defibrillation for pulseless VT, VF 	
Selects correct paddle/electrode pad for infant or child; places paddles/electrode pads in correct position	
Demonstrates correct and safe synchronized cardioversion <ul style="list-style-type: none"> Places device in synchronized mode Selects appropriate energy (0.5 to 1 J/kg for initial shock) Charges, clears, delivers current 	
Demonstrates correct and safe manual defibrillation <ul style="list-style-type: none"> Places device in unsynchronized mode Selects energy (2 to 4 J/kg for initial shock) Charges, clears, delivers current 	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

Vascular Access Skills Station Competency Checklist



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Verbalizes indications for IO insertion	
Verbalizes sites for IO insertion (anterior tibia, distal femur, medial malleolus, anterior-superior iliac spine)	
Verbalizes contraindications for IO placement <ul style="list-style-type: none"> • Fracture in extremity • Previous insertion attempt in the same bone • Infection overlying bone 	
Inserts IO catheter safely	
Verbalizes how to confirm IO catheter is in correct position; verbalizes how to secure IO catheter	
Attaches IV line to IO catheter; demonstrates giving IO fluid bolus by using 3-way stopcock and syringe	
Shows how to determine correct drug doses by using a color-coded length-based tape or other resource	
The following is optional:	
Verbalizes correct procedure for establishing IV access	

STOP TEST

Instructor Notes

- Place a check in the box next to each step the student completes successfully.
- If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation).

Test Results Circle **PASS** or **NR** to indicate pass or needs remediation: **PASS** **NR**

Instructor Initials _____ Instructor Number _____ Date _____

PALS Case Scenario Testing Checklist Respiratory Case Scenario Upper Airway Obstruction



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs	
Directs administration of 100% oxygen or supplemental oxygen as needed to support oxygenation	
Directs application of cardiac monitor and pulse oximetry	
Identifies signs and symptoms of upper airway obstruction	
Categorizes as respiratory distress or failure	
Directs administration of nebulized epinephrine and corticosteroid (for croup), or IM epinephrine and IV corticosteroid (for anaphylaxis)	
States indications for bag-mask ventilation and/or other airway or ventilation support	
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the indications for bag-mask ventilation and/or other airway or ventilation support?"</i>	
Directs establishment of IV or IO access, if indicated	
Directs reassessment of patient in response to treatment	
Case Conclusion/Debriefing	
<i>The following step is evaluated only if the student's scope of practice applies</i>	
Describes how to estimate correct endotracheal tube size for this patient	
<i>If the student does not verbalize the above, prompt the student with the following question: "How would you estimate the endotracheal tube size for this infant with upper airway obstruction?"</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Respiratory Case Scenario Lower Airway Obstruction



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs	
Directs administration of 100% oxygen or supplemental oxygen as needed to support oxygenation	
Directs application of cardiac monitor and pulse oximetry	
Identifies signs and symptoms of lower airway obstruction	
Categorizes as respiratory distress or failure	
Directs administration of albuterol and corticosteroids (for asthma) or suctioning or possible additional laboratory studies (for bronchiolitis)	
States indications for bag-mask ventilation and/or other airway or ventilation support	
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the indications for bag-mask ventilation and/or other airway or ventilation support?"</i>	
Directs establishment of IV or IO access, if appropriate	
Directs reassessment of patient in response to treatment	
Case Conclusion/Debriefing	
<i>The following step is evaluated only if the student's scope of practice applies</i>	
States indications for endotracheal intubation	
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the indications for endotracheal intubation?"</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Respiratory Case Scenario Lung Tissue Disease



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs	
Directs administration of 100% oxygen (or supplemental oxygen as needed to support oxygenation) and evaluates response	
Identifies indications for bag-mask ventilation and/or additional airway or ventilation support	
Describes methods to verify that bag-mask ventilation is effective	
Directs application of cardiac monitor and pulse oximetry	
Identifies signs and symptoms of lung tissue disease	
Categorizes as respiratory distress or failure	
Directs establishment of IV or IO access	
Directs reassessment of patient in response to treatment	
Identifies need for involvement of advanced provider with expertise in pediatric intubation and mechanical ventilation	
Case Conclusion/Debriefing	
<i>The following step is evaluated only if the student's scope of practice applies</i>	
States indications for endotracheal intubation	
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the indications for endotracheal intubation?"</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario **Testing Checklist** **Respiratory Case Scenario** **Disordered Control of Breathing**



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs	
Directs administration of 100% oxygen (or supplemental oxygen as needed to support oxygenation) and evaluates response	
Identifies indications for bag-mask ventilation and/or additional airway or ventilation support	
Describes methods to verify that bag-mask ventilation is effective	
Directs application of cardiac monitor and pulse oximetry	
Identifies signs of disordered control of breathing	
Categorizes as respiratory distress or failure	
Directs establishment of IV or IO access	
Directs reassessment of patient in response to treatment	
Identifies need for involvement of advanced provider with expertise in pediatric intubation and mechanical ventilation	
Case Conclusion/Debriefing	
<i>The following step is evaluated only if the student's scope of practice applies</i>	
States indications for endotracheal intubation	
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the indications for endotracheal intubation?"</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Shock Case Scenario Hypovolemic Shock



American Academy
of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™

Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs	
Directs administration of 100% oxygen	
Directs application of cardiac monitor and pulse oximetry	
Identifies signs and symptoms of hypovolemic shock	
Categorizes as compensated or hypotensive shock	
Directs establishment of IV or IO access	
Directs rapid administration of a 20 mL/kg fluid bolus of isotonic crystalloid; repeats as needed to treat signs of shock	
Reassesses patient during and after each fluid bolus. Stops fluid bolus if signs of heart failure (worsening respiratory distress, development of hepatomegaly or rales/crackles) develop	
Directs reassessment of patient in response to each treatment	
Case Conclusion/Debriefing	
States therapeutic end points during shock management	
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the therapeutic end points during shock management?"</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Shock Case Scenario Obstructive Shock



American Academy
of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™

Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs	
Directs application of cardiac monitor and pulse oximetry	
Verbalizes DOPE mnemonic for intubated patient who deteriorates	
<i>If the student does not verbalize the above, prompt the student with the following questions: "What mnemonic is helpful to recall when the intubated patient deteriorates? What does this mnemonic mean?"</i>	
Identifies signs and symptoms of obstructive shock	
States at least 2 causes of obstructive shock	
<i>If the student does not state the above, prompt the student with the following statement: "Tell me at least 2 causes of obstructive shock."</i>	
Categorizes as compensated or hypotensive shock	
Directs establishment of IV or IO access, if needed	
Directs rapid administration of a fluid bolus of isotonic crystalloid, if needed (ie, for cardiac tamponade, massive pulmonary embolus)	
Directs appropriate treatment for obstructive shock (needle decompression for tension pneumothorax; fluid bolus, and pericardiocentesis for cardiac tamponade; oxygen, ventilatory support, fluid bolus, and expert consultation for massive pulmonary embolus; prostaglandin infusion and expert consultation for neonate with ductal-dependent congenital heart disease and constriction/closure of the ductus arteriosus)	
Directs reassessment of patient in response to treatment	
Case Conclusion/Debriefing	
States therapeutic end points during shock management	
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the therapeutic end points during shock management?"</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Shock Case Scenario Distributive Shock



American Academy
of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™

Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs	
Directs administration of 100% oxygen	
Directs application of cardiac monitor and pulse oximetry	
Identifies signs and symptoms of distributive (septic) shock	
Categorizes as compensated or hypotensive shock	
Directs establishment of IV or IO access	
Directs rapid administration of a 10-20 mL/kg fluid bolus of isotonic crystalloid for septic shock and 20 mL/kg fluid bolus of isotonic crystalloid for anaphylactic shock; repeats as needed (with careful reassessment) to treat shock	
Reassesses patient during and after each fluid bolus. Stops fluid bolus if signs of heart failure (worsening respiratory distress, development of hepatomegaly or rales/crackles) develop	
Directs initiation of vasoactive drug therapy within first hour of care for fluid-refractory shock	
Directs reassessment of patient in response to treatment	
Directs early administration of antibiotics (within first hour after shock is identified)	
Case Conclusion/Debriefing	
States therapeutic end points during shock management	
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the therapeutic end points during shock management?"</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Shock Case Scenario Cardiogenic Shock



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs	
Directs administration of 100% oxygen	
Directs application of cardiac monitor and pulse oximetry	
Identifies signs and symptoms of cardiogenic shock	
Categorizes as compensated or hypotensive shock	
Directs establishment of IV or IO access	
Directs slow administration of a 5 to 10 mL/kg fluid bolus of isotonic crystalloid over 10 to 20 minutes and reassesses patient during and after fluid bolus. Stops fluid bolus if signs of heart failure worsen	
Directs reassessment of patient in response to treatment	
Recognizes the need to obtain expert consultation from pediatric cardiologist	
Identifies need for inotropic/vasoactive drugs during treatment of cardiogenic shock	
<i>If the student does not indicate the above, prompt the student with the following question: "What are the indications for inotropic/vasoactive drugs during cardiogenic shock?"</i>	
Case Conclusion/Debriefing	
States therapeutic end points during shock management	
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the therapeutic end points during shock management?"</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Cardiac Case Scenario Supraventricular Tachycardia



Student Name _____ Date of Test _____

Critical Performance Steps		Check if done correctly
Team Leader		
Assigns team member roles		
Uses effective communication throughout		
Patient Management		
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs		
Directs application of cardiac monitor and pulse oximetry		
Directs administration of supplemental oxygen		
Identifies narrow-complex tachycardia (ie, SVT with adequate perfusion) and verbalizes how to distinguish between ST and SVT		
<i>If the student does not verbalize the above, prompt the student with the following question: "How do you distinguish between ST and SVT?"</i>		
Directs performance of appropriate vagal maneuvers		
Directs establishment of IV or IO access		
Directs preparation and administration of appropriate doses (first and, if needed, second) of adenosine		
States the rationale for the strong recommendation for expert consultation before providing synchronized cardioversion if the stable child with SVT fails to respond to vagal maneuvers and adenosine		
Directs or describes appropriate indications for and safe delivery of attempted cardioversion at 0.5 to 1 J/kg (subsequent doses increased by 0.5 to 1 J/kg, not to exceed 2 J/kg)		
Performs reassessment of patient in response to treatment		
Case Conclusion/Debriefing		
Discusses indications and appropriate energy doses for synchronized cardioversion		
<i>If the student does not verbalize the above, prompt the student with the following question: "What are the indications and appropriate energy doses for synchronized cardioversion?"</i>		

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Cardiac Case Scenario Bradycardia



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Directs assessment of airway, breathing, circulation, disability, and exposure, including vital signs	
Identifies bradycardia associated with cardiopulmonary compromise/failure	
Directs initiation of bag-mask ventilation with 100% oxygen	
Directs application of cardiac monitor and pulse oximetry	
Reassesses heart rate and systemic perfusion after initiation of bag-mask ventilation	
Recognizes indications for high-quality CPR (chest compressions plus ventilation) in a bradycardic patient	
<i>If the student does not indicate the above, prompt the student with the following question: "What are the indications for high-quality CPR in a bradycardic patient?"</i>	
Directs establishment of IV or IO access	
Directs or discusses preparation for and appropriate administration and dose (0.01 mg/kg IV/IO [0.1 mL/kg of 0.1 mg/mL concentration]) of epinephrine	
Performs reassessment of patient in response to treatment	
Case Conclusion/Debriefing	
Verbalizes consideration of 3 potential causes of bradycardia in infants and children	
<i>If the student does not verbalize the above, prompt the student with the following statement: "Tell me 3 potential causes of bradycardia in infants and children."</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Cardiac Case Scenario Asystole/PEA



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Identifies cardiac arrest	
Directs immediate initiation of high-quality CPR, and ensures performance of high-quality CPR at all times	
Directs placement of pads/leads and activation of monitor/defibrillator	
Identifies asystole or PEA	
Directs establishment of IO or IV access	
Directs preparation and administration of appropriate dose of epinephrine at appropriate intervals	
Directs checking rhythm approximately every 2 minutes while minimizing interruptions in chest compressions	
Case Conclusion/Debriefing	
Verbalizes at least 3 reversible causes of PEA or asystole	
<i>If the student does not verbalize the above, prompt the student with the following statement: "Tell me at least 3 reversible causes of PEA or asystole."</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		

PALS Case Scenario Testing Checklist Cardiac Case Scenario VF/Pulseless VT



Student Name _____ Date of Test _____

Critical Performance Steps	Check if done correctly
Team Leader	
Assigns team member roles	
Uses effective communication throughout	
Patient Management	
Identifies cardiac arrest	
Directs immediate initiation of high-quality CPR, and ensures performance of high-quality CPR at all times	
Directs placement of pads/leads and activation of monitor/defibrillator	
Identifies VF or pulseless VT cardiopulmonary arrest	
Directs safe performance of attempted defibrillation at 2 J/kg	
After delivery of every shock, directs immediate resumption of CPR, beginning with chest compressions	
Directs establishment of IO or IV access	
Directs preparation and administration of appropriate dose of epinephrine at appropriate intervals	
Directs safe delivery of second shock at 4 J/kg (subsequent doses 4 to 10 J/kg, not to exceed 10 J/kg or standard adult dose for that defibrillator)	
Directs preparation and administration of appropriate dose of antiarrhythmic (amiodarone or lidocaine) at appropriate time	
Case Conclusion/Debriefing	
Verbalizes possible need for additional doses of epinephrine and antiarrhythmic (amiodarone or lidocaine), and consideration of reversible causes of arrest (H's and T's)	
<i>If the student does not verbalize the above, prompt the student with the following question: "If VF persists despite the therapies provided, what else should you administer or consider?"</i>	

STOP TEST

Instructor Notes <ul style="list-style-type: none"> Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results Circle PASS or NR to indicate pass or needs remediation:	PASS	NR
Instructor Initials _____ Instructor Number _____ Date _____		